

98. A number consists of 3 digits whose sum is 10. The middle digit is equal to the sum of the other two and the number will be increased by 99 if its digits are reversed. The number is (Hotel Management, 2003)
- (a) 145 (b) 253
(c) 370 (d) 352
99. A two-digit number becomes five-sixth of itself when its digits are reversed. The two digits differ by one. The number is
- (a) 45 (b) 54
(c) 56 (d) 65
100. If the square of a two-digit number is reduced by the square of the number formed by reversing the digits of the number, the final result is (N.M.A.T., 2008)
- (a) divisible by 11 (b) divisible by 9
(c) necessarily irrational (d) Both (a) and (b)
101. A number consists of two digits such that the digit in the ten's place is less by 2 than the digit in the unit's place. Three times the number added to $\frac{6}{7}$ times the number obtained by reversing the digits equals 108. The sum of the digits in the number is (S.S.C., 2003)
- (a) 6 (b) 7
(c) 8 (d) 9
102. The digit in the unit's place of a number is equal to the digit in the ten's place of half of that number and the digit in the ten's place of that number is less than the digit in unit's place of half of the number by 1. If the sum of the digits of the number is 7, then what is the number? (S.B.I.P.O., 2001)
- (a) 34 (b) 52
(c) 162 (d) Data inadequate
(e) None of these
103. In a two-digit number, the digit in the unit's place is more than twice the digit in ten's place by 1. If the digits in the unit's place and the ten's place are interchanged, difference between the newly formed number and the original number is less than the original number by 1. What is the original number?
- (a) 25 (b) 37
(c) 49 (d) 52
(e) 73
104. A certain number of two digits is three times the sum of its digits and if 45 be added to it, the digits are reversed. The number is (N.M.A.T. 2006; L.I.C.A.A.O., 2003)
- (a) 23 (b) 27
(c) 32 (d) 72
105. A two-digit number is such that the product of the digits is 8. When 18 is added to the number, then the digits are reversed. The number is (M.B.A., 2003)
- (a) 18 (b) 24
(c) 42 (d) 81
106. In a number of three digits, the digits in the unit's place and in the hundred's place are equal and the sum of all the digits is 8. The number of such numbers is
- (a) 3 (b) 4
(c) 5 (d) 6
107. In a three-digit number, the digit in the unit's place is 75% of the digit in the ten's place. The digit in the ten's place is greater than the digit in the hundred's place by 1. If the sum of the digits in the ten's place and the hundred's place is 15, what is the number? (Bank P.O., 2006)
- (a) 687
(b) 786
(c) 795
(d) Cannot be determined
(e) None of these
108. The product of two fractions is $\frac{14}{15}$ and their quotient is $\frac{35}{24}$. The greater fraction is (S.S.C., 2005)
- (a) $\frac{4}{5}$ (b) $\frac{7}{6}$
(c) $\frac{7}{4}$ (d) $\frac{7}{3}$
109. In a pair of fractions, fraction A is twice the fraction B and the product of two fractions is $\frac{2}{25}$. What is the value of fraction A?
- (a) $\frac{1}{5}$ (b) $\frac{1}{25}$
(c) $\frac{2}{5}$ (d) Data inadequate
110. If the difference between the reciprocal of a positive proper fraction and the fraction itself be $\frac{9}{20}$, then the fraction is (C.P.O., 2006)
- (a) $\frac{3}{5}$ (b) $\frac{4}{5}$
(c) $\frac{5}{4}$ (d) $\frac{3}{10}$
111. The sum of the numerator and denominator of a fraction is 11. If 1 is added to the numerator and 2 is subtracted from the denominator, it becomes $\frac{2}{3}$. The fraction is
- (a) $\frac{5}{6}$ (b) $\frac{6}{5}$
(c) $\frac{3}{8}$ (d) $\frac{8}{3}$