A)

CPRD Aurum

135,666 participants had undergo an AVR

91,310 excluded
- **58,021** had the record while not in observation
- **0** had missing record date
- **7,106** did not have 365 days of prior observation before their first AVR
- **26,183** had the procedure before the study period start (before 01/01/2012)

44,356 participants included in the AVR cohort

13,500 participants underwent TAVI
- **11,566** identified through the combination of AVR and transcatheter procedure codes
- **4,266** identified through a TAVI code

30,856 participants underwent SAVR

B)

DataLoch Lothian

x participants had undergo an AVR

x excluded
- **x** had the record while not in observation
- **x** had missing record date
- **x** did not have 365 days of prior observation before their first AVR
- **x** had the procedure before the study period start (before 01/01/2012)

x participants included in the AVR cohort

x participants underwent TAVI
- **x** identified through the combination of AVR and transcatheter procedure codes
- **x** identified through a TAVI code

x participants underwent SAVR

A)

CPRD Aurum

454,684 participants had a diagnosis of aortic stenosis

252,724 excluded
- **160,368** had the record while not in observation
- **0** had missing record date
- **76,878** were diagnosed before 01-01-2012
- **15,478** did not have 365 days of prior observation

201,960 participants

166,925 participants excluded because they did not undergo AVR within one year of the diagnosis

35,035 participants

4,382 participants excluded due to a diagnosis of aortic insufficiency or endocarditis on or after the initial diagnosis but before undergoing AVR

30,653 participants included in the aortic stenosis cohort

B)

DataLoch Lothian

x participants had a diagnosis of aortic stenosis

x excluded
- **x** had the record while not in observation
- **x** had missing record date
- **x** were diagnosed before 01-01-2012
- **x** did not have 365 days of prior observation

x participants

x participants excluded because they did not undergo AVR within one year of the diagnosis

x participants

x participants excluded due to a diagnosis of aortic insufficiency or endocarditis on or after the initial diagnosis but before undergoing AVR

x participants included in the aortic stenosis cohort

A)

CPRD Aurum

247,618 participants had a diagnosis of aortic insufficiency

139,439 excluded
- **95,253** had the record while not in observation
- **0** had missing record date
- **37,219** were diagnosed before 01-01-2012
- **6,967** did not have 365 days of prior observation

108,179 participants

97,795 participants excluded because they did not undergo AVR within one year of the diagnosis

10,384 participants

5,037 participants excluded due to a diagnosis of aortic stenosis or endocarditis on or after the initial diagnosis but before undergoing AVR

5,315 participants included in the aortic stenosis cohort

B)

DataLoch Lothian

x participants had a diagnosis of aortic stenosis

x excluded
- **x** had the record while not in observation
- **x** had missing record date
- **x** were diagnosed before 01-01-2012
- **x** did not have 365 days of prior observation

x participants

x participants excluded because they did not undergo AVR within one year of the diagnosis

x participants

x participants excluded due to a diagnosis of aortic stenosis or endocarditis on or after the initial diagnosis but before undergoing AVR

x participants included in the aortic stenosis cohort

A)

CPRD Aurum

61,864 participants had a diagnosis of aortic endocarditis

45,357 excluded
- **30,945** had the record while not in observation
- **0** had missing record date
- **13,011** were diagnosed before 01-01-2012
- **1,401** did not have 365 days of prior observation

16,507 participants

16,015 participants excluded because they did not undergo AVR within one year of the diagnosis

492 participants

400 participants excluded due to a diagnosis of aortic stenosis or insufficiency on or after the initial diagnosis but before undergoing AVR

92 participants included in the aortic endocarditis cohort

B)

DataLoch Lothian

x participants had a diagnosis of aortic stenosis

x excluded
- **x** had the record while not in observation
- **x** had missing record date
- **x** were diagnosed before 01-01-2012
- **x** did not have 365 days of prior observation

x participants

x participants excluded because they did not undergo AVR within one year of the diagnosis

x participants

x participants excluded due to a diagnosis of aortic stenosis or insufficiency on or after the initial diagnosis but before undergoing AVR

x participants included in the aortic stenosis cohort