



Congenital Aortic Stenosis


Baseline Summary

Baseline Summary Table

Cohort name	Age group	Sex	Variable name	Variable level	Estimate name	CDM name
						GOSH_OMOP
						group_level
Aortic Stenosis Congenital	overall	overall	Number subjects	—	N	565
			Age	—	Median [Q25 - Q75]	7 [1 - 12]
					Mean (SD)	7.12 (5.80)
					Range	0 to 17
			Age group	0 to 1	N (%)	165 (29.20%)
				2 to 12	N (%)	269 (47.61%)
				13 to 17	N (%)	131 (23.19%)
			Sex	Female	N (%)	180 (31.86%)
				Male	N (%)	385 (68.14%)
			Comorbidity cohorts flag -inf to 0	Mitral valve disorder	N (%)	79 (13.98%)
				Aortic insufficiency	N (%)	245 (43.36%)
				Atrial septal defect	N (%)	82 (14.51%)
				Cardiomegaly	N (%)	134 (23.72%)
				Aortic aneurysm	N (%)	109 (19.29%)


Cohort name	Age group	Sex	Variable name	Variable level	Estimate name	CDM name
						GOSH_OMOP
						group_level
				Patent ductus arteriosus	N (%)	74 (13.10%)

Age Group by Sex



Distribution of children with congenital aortic stenosis across age groups and sex in the index cohort at Great Ormond Street Hospital.


Top Pre-Index Conditions



Most common pre-index comorbid conditions in the congenital aortic stenosis cohort, ranked by the percentage of children with each condition.


Outcome Risk

Overall Cumulative Failure by Outcome




Kaplan-Meier cumulative failure curves for the main study outcomes in the congenital aortic stenosis cohort, shown overall for each outcome across follow-up.

Age-Stratified Cumulative Failure by Outcome



Supplementary Figures

Age-Sex Cumulative Failure



Kaplan-Meier cumulative failure curves for each study outcome stratified jointly by age group and sex. This figure is presented as supplementary because the panel layout is more detailed.