






However, we did not talk
about how to estimate the
Weights and Biases.






So let's talk about how **Backpropagation** optimizes the **Weights** and **Biases** in this and other **Neural Networks**.






So let's talk about how **Backpropagation** optimizes the **Weights** and **Biases** in this and other **Neural Networks**.






So let's talk about how **Backpropagation** optimizes the **Weights** and **Biases** in this and other **Neural Networks**.






So let's talk about how **Backpropagation** optimizes the **Weights** and **Biases** in this and other **Neural Networks**.







So let's talk about how **Backpropagation** optimizes the **Weights** and **Biases** in this and other **Neural Networks**.






NOTE: Backpropagation is relatively simple, but there are a ton of details. So I've spit it up into bite sized pieces.






In this part, we talk about the **Main Ideas of Backpropagation:**






In this part, we talk about the **Main Ideas of Backpropagation:**




1) Using the **Chain Rule** to calculate derivatives...

$$\frac{d \text{SSR}}{d \text{bias}} = \frac{d \text{SSR}}{d \text{Predicted}} \times \frac{d \text{Predicted}}{d \text{bias}}$$






In this part, we talk about the **Main Ideas of Backpropagation:**



2) Plugging the derivatives into **Gradient Descent** to optimize parameters.






In this part, we talk about the **Main Ideas of Backpropagation:**

Yes (1)
Output
No (0)

Dosage

2) Plugging the derivatives into **Gradient Descent** to optimize parameters.







In this part, we talk about the **Main Ideas of Backpropagation:**

Yes (1)
Output
No (0)




Dosage

2) Plugging the derivatives into **Gradient Descent** to optimize parameters.





In this part, we talk about the **Main Ideas of Backpropagation:**







First, so we can be clear about which specific **Weights** we are talking about, let's give each one a name.







First, so we can be clear about which specific **Weights** we are talking about, let's give each one a name.







And let's name
each **Bias**.







And let's name
each **Bias**.






...and works its way backwards to estimate all of the other parameters.






...and works its way backwards to estimate all of the other parameters.






...and works its way backwards to estimate all of the other parameters.






...and works its way backwards to estimate all of the other parameters.






However, we can discuss all of the **Main Ideas** behind **Backpropagation** by just estimating the last **Bias**, b_3 .






So, in order to start from the back, let's assume that we already have **optimal values** for all of the parameters...






So, in order to start from the back, let's assume that we already have **optimal values** for all of the parameters...

...except for the last **Bias term, b_3 .**






NOTE: Throughout this and the next **StatQuests**, I will make parameter values that have already been optimized **green**...






NOTE: Throughout this and the next **StatQuests**, I will make parameter values that have already been optimized **green**...

...and unoptimized parameters will be **red**.






ALSO NOTE: To keep the math simple, let's assume **Dosages** go from **0** (low) to **1** (high).






Now, if we run
Dosages from 0 to 1...






...to get this
green squiggle.






Now we are ready to add the final
Bias, b_3 , to the green squiggle.






Because we don't yet know the optimal value for b_3 , we have to give it an initial value...






...and because **Bias** terms are frequently initialized to **0**, we will set $b_3 = 0$.






Now, adding **0** to all of the y-axis coordinates on the **green squiggle** leaves it right where it is.






However, that means the **green squiggle** is pretty far from the data that we observed.






We can quantify how good the **green squiggle** fits the data by calculating the **Sum of the Squared Residuals.**





A **Residual** is the difference between the **Observed** and **Predicted** values.




Residual = (Observed - Predicted)






For example,
this **Residual**...





For example,
this **Residual**...

...is the **Observed**
value, 0...






For example,
this **Residual**...

...is the **Observed**
value, 0...

...minus the
Predicted value from
the **green squiggle**,

-2.6






For example,
this **Residual**...


...is the **Observed**
value, 0...

...minus the
Predicted value from
the **green squiggle**,





This Residual...




Residual = (Observed - Predicted)






This Residual...
...is the Observed
value, 1...





This Residual...
...is the Observed
value, 1...






This Residual...

...is the Observed
value, 1...

...minus the
Predicted value from
the green squiggle,
-1.61




$$\text{Residual} = (1 - \text{Predicted})$$






Lastly, this
Residual...





Lastly, this
Residual...
...is the **Observed**
value, 0...






Lastly, this
Residual...

...is the **Observed**
value, 0...

...minus the
Predicted value from
the **green squiggle**,
-2.61






Lastly, this
Residual...


...is the **Observed**
value, 0...

...minus the
Predicted value from
the **green squiggle**,
-2.61






Now we square
each **Residual**...






Now we square
each **Residual**...





...and add them all together...




A large black curved arrow points from the text "...and add them all together..." to three mathematical expressions below:

$$(0 - -2.6)^2$$
$$(1 - -1.61)^2$$
$$(0 - -2.61)^2$$





...and add them all together...




A diagram of a neural network node. It has three inputs: "Input" (represented by a white square), a bias input "b3" (represented by a red square with "+ 0.00"), and a summation input "sum" (represented by a blue square with "sum"). The output is a green square labeled "Output". A curved black arrow points from the text "...and add them all together..." to the summation input "sum".

$$(0 - -2.6)^2 + (1 - -1.61)^2 + (0 - -2.61)^2$$





...to get **20.4** for the **Sum of the Squared Residuals (SSR)**.




$$\begin{aligned} \text{SSR} = & (0 - -2.6)^2 \\ & + (1 - -1.61)^2 \\ & + (0 - -2.61)^2 = 20.4 \end{aligned}$$





So, when $b_3 = 0$, the
SSR = 20.4...




$$\begin{aligned} \text{SSR} = & (0 - -2.6)^2 \\ & + (1 - -1.61)^2 \\ & + (0 - -2.61)^2 = 20.4 \end{aligned}$$






...and that corresponds to this location on this graph...




$$\begin{aligned} \text{SSR} = & (0 - -2.6)^2 \\ & + (1 - -1.61)^2 \\ & + (0 - -2.61)^2 = 20.4 \end{aligned}$$





...that has the **SSR**
on the y-axis...




$$\begin{aligned} \text{SSR} = & (0 - -2.6)^2 \\ & + (1 - -1.61)^2 \\ & + (0 - -2.61)^2 = 20.4 \end{aligned}$$






...and the **Bias**, b_3 , on
the x-axis.





$$\begin{aligned} \text{SSR} = & (0 - -2.6)^2 \\ & + (1 - -1.61)^2 \\ & + (0 - -2.61)^2 = 20.4 \end{aligned}$$





Now, if we increase b_3 to 1...







...then we add 1 to the y-axis
coordinates on the **green**
squiggle...






...and shift the green squiggle up 1...





...and we end up with shorter
Residuals.






When we do the math...





...and that corresponds to this point on our graph.




$$\begin{aligned} \text{SSR} = & (0 - -1.6)^2 \\ & + (1 - -0.61)^2 \\ & + (0 - -1.61)^2 = 7.8 \end{aligned}$$






If we increase b_3 to 2, then
the **SSR** = 1.11...




$$\begin{aligned} \text{SSR} &= (0 - -0.6)^2 \\ &\quad + (1 - 0.39)^2 \\ &\quad + (0 - -0.61)^2 = 1.11 \end{aligned}$$





...and if we increase b_3 to 3,
then the **SSR = 0.46**.






$$\begin{aligned} \text{SSR} = & (0 - 0.4)^2 \\ & + (1 - 1.39)^2 \\ & + (0 - 0.39)^2 = 0.46 \end{aligned}$$







And if we had time to plug in tons
of values for b_3 , we would get
this **pink curve**...








...and we could find the lowest point, which corresponds to the value for b_3 that results in the lowest **SSR**, here.







...and we could find the lowest point, which corresponds to the value for b_3 that results in the lowest **SSR**, here.







However, instead of plugging in tons of values to find the lowest point in the **pink curve**, we use **Gradient Descent** to find it relatively quickly.







However, instead of plugging in tons of values to find the lowest point in the **pink curve**, we use **Gradient Descent** to find it relatively quickly.







However, instead of plugging in tons of values to find the lowest point in the **pink curve**, we use **Gradient Descent** to find it relatively quickly.







However, instead of plugging in tons of values to find the lowest point in the **pink curve**, we use **Gradient Descent** to find it relatively quickly.






And that means we need to find the derivative of the **Sum of the Squared Residuals** with respect to b_3 .






Now, remember the **Sum of the Squared Residuals** equals...




SSR =






...the first **Residual** squared...




$$SSR = (\text{Observed}_1 - \text{Predicted}_1)^2$$






...plus all of the other squared **Residuals**.



$$\text{SSR} = (\text{Observed}_1 - \text{Predicted}_1)^2 + (\text{Observed}_2 - \text{Predicted}_2)^2$$







Now, because this equation
takes up a lot of space...



$$\begin{aligned} \text{SSR} = & (\text{Observed}_1 - \text{Predicted}_1)^2 \\ & + (\text{Observed}_2 - \text{Predicted}_2)^2 \\ & + (\text{Observed}_3 - \text{Predicted}_3)^2 \end{aligned}$$







...we can make it
smaller by using
summation notation.

$$SSR = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$





$$\boxed{\begin{aligned} SSR &= (\text{Observed}_1 - \text{Predicted}_1)^2 \\ &+ (\text{Observed}_2 - \text{Predicted}_2)^2 \\ &+ (\text{Observed}_3 - \text{Predicted}_3)^2 \end{aligned}}$$





Now let's talk a little bit more about the **Predicted** values.


$$SSR = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$







Each Predicted value comes from the green squiggle...

$$SSR = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$




Predicted_i = green squiggle_i





...and the **green squiggle** comes from the last part of the **Neural Network**.


$$SSR = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$





In other words, the **green squiggle** is the **sum** of the **blue** and **orange** curves...


$$SSR = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$






...plus b_3 .

$$SSR = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$






Now remember, we want to use
Gradient Descent to optimize b_3 ...



$$SSR = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$



Predicted_i = **green squiggle** = **blue** + **orange** + **b_3**





$$\frac{d \text{SSR}}{d b_3}$$

...and that means we need to take the derivative of the **SSR** with respect to b_3 .



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$


$$\text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$






$$\frac{d \text{SSR}}{d b_3}$$

...and that means we need to take the derivative of the **SSR** with respect to b_3 .



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$


$$\text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$






$$\frac{d \text{SSR}}{d b_3}$$

...and that means we need to take the derivative of the **SSR** with respect to b_3 .



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$


$$\text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$






$$\frac{d \text{SSR}}{d b_3}$$

...and that means we need to take the derivative of the **SSR** with respect to b_3 .



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$


$$\text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$






$$\frac{d \text{SSR}}{d b_3}$$

And because the **SSR** are linked to b_3 ...



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$


Predicted_i = **green squiggle_i** = **blue** + **orange** + **b_3**






$$\frac{d \text{SSR}}{d b_3}$$

...by the **Predicted** values...



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$


$$\text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$






$$\frac{d \text{SSR}}{d b_3}$$

...we can use
The Chain Rule...



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \boxed{\text{Predicted}_i})^2$$



Predicted_i = green squiggle_i = blue + orange + b₃





$$\frac{d \text{SSR}}{d b_3}$$

...to solve for the derivative of the **SSR** with respect to b_3 .



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$


Predicted_i = **green squiggle_i** = **blue** + **orange** + **b_3**






$$\frac{d \text{SSR}}{d b_3} = \leftarrow$$

The Chain Rule says that
the derivative of the **SSR**
with respect to b_3 ...



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \boxed{\text{Predicted}_i})^2$$


$$\boxed{\text{Predicted}_i} = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$






$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}}$$

...is the derivative of the **SSR** with respect to the **Predicted** values...



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$


$\text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$






$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

...times the derivative of the **Predicted** values with respect to b_3 .



$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$

Predicted_i = green squiggle_i = blue + orange + **b_3**





$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}} \times \frac{d \text{Predicted}}{d b_3}$$

$$\frac{d \text{SSR}}{d \text{Predicted}} =$$

Now, we can solve for the derivative of the **SSR** with respect to the **Predicted** values by first substituting in the equation...

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}}$$

Now, we can solve for the derivative of the **SSR** with respect to the **Predicted** values by first substituting in the equation...

$$\frac{d \text{SSR}}{d \text{Predicted}} = \frac{d}{d \text{Predicted}} \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$



$$\text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

...and then use
The Chain Rule...

$$\frac{d \text{ SSR}}{d \text{ Predicted}} = \frac{d}{d \text{ Predicted}} \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}} \times \frac{d \text{Predicted}}{d b_3}$$

...and we have the derivative
of the **SSR** with respect to
the **Predicted** values.

$$\frac{d \text{SSR}}{d \text{Predicted}} = \frac{d}{d \text{Predicted}} \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$

$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$


$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

So let's move
that up here...

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$



$$\frac{d \text{ SSR}}{d \text{ Predicted}} = \frac{d}{d \text{ Predicted}} \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times$$



$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

...and now we are done
with the first part.

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}} \times \frac{d \text{Predicted}}{d b_3}$$

Now let's solve for the second part, the derivative of the **Predicted** values with respect to b_3 .

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

We start by plugging in the equation for the **Predicted** values.

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

$$\frac{d \text{ Predicted}}{d b_3}$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$

$$\text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

We start by plugging in the equation for the **Predicted** values.

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle}$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$

Predicted_i = **green squiggle_i** = **blue** + **orange** + **b₃**



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

We start by plugging in the equation for the **Predicted** values.

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle} = \frac{d}{d b_3} (\text{blue} + \text{orange} + b_3)$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$

Predicted_i = green squiggle_i = blue + orange + b₃




$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

Remember, the **blue** and **orange** curves were created before we got to b_3 ...

$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle} = \frac{d}{d b_3} (\text{blue} + \text{orange} + b_3)$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2$$






$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

...so the derivative of the **blue curve** with respect to b_3 is 0, because the **blue curve** is independent of b_3 ...



$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle} = \frac{d}{d b_3} (\text{blue} + \text{orange} + b_3) = 0$$


$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

...and the derivative of the
orange curve with respect
to b_3 is also 0.

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$



$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle} = \frac{d}{d b_3} (\text{blue} + \text{orange} + b_3) = 0 + 0$$


$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$

Lastly, the derivative of b_3 with respect to b_3 is 1.



$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle} = \frac{d}{d b_3} (\text{blue} + \text{orange} + b_3) = 0 + 0 + 1$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

Now we just add everything up...

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$



$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle} = \frac{d}{d b_3} (\text{blue} + \text{orange} + b_3) = 0 + 0 + 1$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

...and the derivative of the **Predicted** values with respect to b_3 is 1.

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i)$$



$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle} = \frac{d}{d b_3} (\text{blue} + \text{orange} + b_3) = 1$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \boxed{x \ 1}$$

So we multiply the derivative of the **SSR** with respect to the **Predicted** values by **1**.

$$\frac{d \text{ Predicted}}{d b_3} = \frac{d}{d b_3} \text{ green squiggle} = \frac{d}{d b_3} (\text{blue} + \text{orange} + b_3) = 1$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \boxed{\times 1}$$

NOTE: This “times 1” part in the equation doesn’t do anything...


$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}} \times \frac{d \text{Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$

...but I'm leaving it in to remind us that the derivative of the **SSR** with respect to **b_3** consists of two parts.


$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$
$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$

The derivative of the **SSR** with respect to the **Predicted** values...

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}} \times \boxed{\frac{d \text{Predicted}}{d b_3}}$$

...and the derivative of the Predicted values with respect to b_3 .


$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \boxed{x \ 1}$$

$$\text{SSR} = \sum_{i=1}^{n=3} (\text{Observed}_i - \text{Predicted}_i)^2 \quad \text{Predicted}_i = \text{green squiggle}_i = \text{blue} + \text{orange} + b_3$$



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$




And at long last, we have
the derivative of the **SSR**
with respect to b_3 .



$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$

And that means we can plug this derivative into **Gradient Descent** to find the optimal value for b_3 .






$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$

And that means we can plug this derivative into **Gradient Descent** to find the optimal value for b_3 .






$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}} \times \frac{d \text{Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$

And that means we can plug this derivative into **Gradient Descent** to find the optimal value for b_3 .






$$\frac{d \text{SSR}}{d b_3} = \frac{d \text{SSR}}{d \text{Predicted}} \times \frac{d \text{Predicted}}{d b_3}$$

$$= \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$


And that means we can plug this derivative into **Gradient Descent** to find the optimal value for b_3 .





$$\frac{d \text{SSR}}{d b_3} = \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$

Anyway, first, we expand the summation.






$$\frac{d \text{SSR}}{d b_3} = -2 \times (\text{Observed}_1 - \text{Predicted}_1) \times 1 \\ + -2 \times (\text{Observed}_2 - \text{Predicted}_2) \times 1 \\ + -2 \times (\text{Observed}_3 - \text{Predicted}_3) \times 1$$




Anyway, first, we expand
the summation.






$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - \text{Predicted}_1) \times 1 \\ + -2 \times (1 - \text{Predicted}_2) \times 1 \\ + -2 \times (0 - \text{Predicted}_3) \times 1$$




Then we plug in the
Observed values....






$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - \text{Predicted}_1) \times 1 \\ + -2 \times (1 - \text{Predicted}_2) \times 1 \\ + -2 \times (0 - \text{Predicted}_3) \times 1$$





...and the values **Predicted** by
the **green squiggle**.






$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - \text{Predicted}_1) \times 1 \\ + -2 \times (1 - \text{Predicted}_2) \times 1 \\ + -2 \times (0 - \text{Predicted}_3) \times 1$$



Remember, we get the **Predicted** values on the **green squiggle** by running the **Dosages** through the **Neural Network**.



$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 \\ + -2 \times (1 - -1.6) \times 1 \\ + -2 \times (0 - -2.61) \times 1 = -15.7$$




Now we just do the
math and get **-15.7...**







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 \\ + -2 \times (1 - -1.6) \times 1 \\ + -2 \times (0 - -2.61) \times 1 = -15.7$$




...and that corresponds to the slope for when $b_3 = 0$.







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$



Step Size = Slope \times Learning Rate




Now we plug the slope into the **Gradient Descent** equation for **Step Size**...







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$



Step Size = -15.7 × Learning Rate


Now we plug the slope into the **Gradient Descent** equation for **Step Size**...







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$

...and in this example, we'll set the **Learning Rate** to **0.1...**





Step Size = $-15.7 \times \text{Learning Rate}$



SQ!
double
BAM!!

$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$

...and that means the
Step Size is **-1.57**.




$$\text{Step Size} = -15.7 \times 0.1 = -1.57$$







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$



$$\text{Step Size} = -15.7 \times 0.1 = -1.57$$

$$\text{New } b_3 = \text{Old } b_3 - \text{Step Size}$$


Now we use the **Step Size** to calculate the new value for b_3 ...







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$

...by plugging in the current value for b_3 , 0...



$$\text{Step Size} = -15.7 \times 0.1 = -1.57$$


$$\text{New } b_3 = \boxed{\text{Old } b_3} - \text{Step Size}$$






$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$

...by plugging in the current value for b_3 , 0...



$$\text{Step Size} = -15.7 \times 0.1 = -1.57$$


$$\text{New } b_3 = 0 - \text{Step Size}$$







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$

...and the **Step Size**,
-1.57.



$$\text{Step Size} = -15.7 \times 0.1 = -1.57$$


$$\text{New } b_3 = 0 - \text{Step Size}$$







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$

...and the **Step Size**,
-1.57.



$$\text{Step Size} = -15.7 \times 0.1 = -1.57$$


$$\text{New } b_3 = 0 - (-1.57)$$







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -2.6) \times 1 + -2 \times (1 - -1.6) \times 1 + -2 \times (0 - -2.61) \times 1 = -15.7$$

And the new value for b_3 is **1.57**.





$$\text{Step Size} = -15.7 \times 0.1 = -1.57$$

$$\text{New } b_3 = 0 - (-1.57) = 1.57$$






$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - \text{Predicted}_1) \times 1 \\ + -2 \times (1 - \text{Predicted}_2) \times 1 \\ + -2 \times (0 - \text{Predicted}_3) \times 1$$




Changing b_3 to 1.57 shifts the **green squiggle** up...




SQ!
double
BAM!!

$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - \text{Predicted}_1) \times 1 + -2 \times (1 - \text{Predicted}_2) \times 1 + -2 \times (0 - \text{Predicted}_3) \times 1$$




Changing b_3 to 1.57 shifts the **green squiggle** up...



SQ!
double
BAM!!

$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - \text{Predicted}_1) \times 1 + -2 \times (1 - \text{Predicted}_2) \times 1 + -2 \times (0 - \text{Predicted}_3) \times 1$$


...and that shrinks the
Residuals.





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$



Now, plugging in the new Predicted values and doing the math gives us **-6.26...**





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$


...which corresponds to the slope when $b_3 = 1.57$.







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$

Then we calculate
the **Step Size**...




Step Size = Slope × Learning Rate







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$

Then we calculate
the **Step Size**...




Step Size = Slope \times Learning Rate







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$



Step Size = -6.26 × Learning Rate




Then we calculate the **Step Size**...







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$



Step Size = -6.26×0.1




Then we calculate
the **Step Size**...







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$

Then we calculate
the **Step Size**...




$$\text{Step Size} = -6.26 \times 0.1 = -0.626$$





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$

...and the new value for b_3 ...





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$

...and the new value for b_3 ...





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$



...and the new value for b_3 ...





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$



...and the new value for b_3 ...





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$




...and the new value for b_3 ...





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$


...and the new value for b_3 ...







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$

...which is
2.19.



$$\text{Step Size} = -6.26 \times 0.1 = -0.626$$


$$\text{New } b_3 = 1.57 - (-0.626) = 2.19$$



SQ!


$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 \\ + -2 \times (1 - -0.03) \times 1 \\ + -2 \times (0 - -1.04) \times 1 = -6.26$$

Changing b_3 to **2.19** shifts the **green squiggle** up further...



SQ!
double
BAM!!


$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - -1.03) \times 1 + -2 \times (1 - -0.03) \times 1 + -2 \times (0 - -1.04) \times 1 = -6.26$$




Changing b_3 to 2.19 shifts the **green squiggle** up further...

SQ!
double
BAM!!

$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - \text{Predicted}_1) \times 1 + -2 \times (1 - \text{Predicted}_2) \times 1 + -2 \times (0 - \text{Predicted}_3) \times 1$$



...and that shrinks the **Residuals** even more.







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - \text{Predicted}_1) \times 1 + -2 \times (1 - \text{Predicted}_2) \times 1 + -2 \times (0 - \text{Predicted}_3) \times 1$$




Now we just keep taking steps...







$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - 0.01) + -2 \times (1 - 1.01) + -2 \times (0 - 0) = \text{close to } 0$$

...until the **Step Size** is close to **0**.




$$\text{Step Size} = \text{close to } 0 \times 0.1 = \text{close to } 0$$

$$\text{New } b_3 = \text{Old } b_3 - \text{Step Size}$$






$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - 0.01) + -2 \times (1 - 1.01) + -2 \times (0 - 0) = \text{close to } 0$$




Step Size = close to 0 \times 0.1 = close to 0

New b_3 = Old b_3 - Step Size




And because the **Step Size** is close to 0 when $b_3 = 2.61\dots$





$$\frac{d \text{SSR}}{d b_3} = -2 \times (0 - 0.01) + -2 \times (1 - 1.01) + -2 \times (0 - 0) = \text{close to } 0$$





Step Size = close to 0 \times 0.1 = close to 0

New b_3 = Old b_3 - Step Size




...we decide that
2.61 is the optimal
value for b_3 .






So the **Main Ideas** for
Backpropagation are that
when a parameter is
unknown, like b_3 ...





...we use **The Chain Rule** to calculate the derivative of the **Sum of the Squared Residuals (SSR)**...






$$\frac{d \text{ SSR}}{d b_3}$$

Yes (1)
Output
No (0)


Dosage

...with respect to the unknown parameter, which in this case was b_3 .






$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}}$$






$$\frac{d \text{ SSR}}{d b_3} = \frac{d \text{ SSR}}{d \text{ Predicted}} \times \frac{d \text{ Predicted}}{d b_3}$$




...with respect to the unknown parameter, which in this case was b_3 .






$$\frac{d \text{SSR}}{d b_3} = \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$




Then we initialize the unknown parameter with a number, and in this case we set $b_3 = 0$...







$$\frac{d \text{SSR}}{d b_3} = \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$




...and used **Gradient Descent** to optimize the unknown parameter.






$$\frac{d \text{SSR}}{d b_3} = \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$






$$\frac{d \text{SSR}}{d b_3} = \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$





$$\frac{d \text{SSR}}{d b_3} = \sum_{i=1}^{n=3} -2 \times (\text{Observed}_i - \text{Predicted}_i) \times 1$$



...and used **Gradient Descent** to optimize the unknown parameter.

