

ELE 535: Machine Learning and Pattern  
Recognition  
Homework 9

Zachary Hervieux-Moore

Friday 21<sup>st</sup> December, 2018

**Exercise 1: Kernels:**

- a) Let  $\mathcal{A}$  be a finite set and for each subset  $\mathcal{U} \subseteq \mathcal{A}$  let  $|\mathcal{U}|$  denote the number of element in  $\mathcal{U}$ . For  $\mathcal{U}, \mathcal{V} \subset \mathcal{A}$ , let  $k(\mathcal{U}, \mathcal{V}) = |\mathcal{U} \cap \mathcal{V}|$ . By finding a suitable feature map, show that  $k(\cdot, \cdot)$  is a kernel on the power set  $\mathcal{P}(\mathcal{A})$  of all subsets of  $\mathcal{A}$ .
- b) Show that  $k(x, z) = \sum_{i=1}^n \cos^2(x_i - z_i)$  is a kernel on  $\mathbb{R}^n$ .
- c) Let  $P \in \mathbb{R}^n \times \mathbb{R}^n$  be a symmetrix PSD. Show that  $k(x, z) = e^{-\frac{1}{2}(x-z)^T P (x-z)}$  is a kernel on  $\mathbb{R}^n$ .
- d)  $k(x, y) = h_t(Ax)^T h_t(Ay)$  where  $A \in \mathbb{R}^{n \times n}$  and  $h_t$  is a thresholding function that maps  $z = [z_i]$  to  $h_t(z) = [\tilde{z}_i]$  with  $\tilde{z}_i = z_i$  if  $|z_i| > t$  and 0 otherwise.

**Answer:**

**Exercise 2:** Let  $k_j$  be a kernel on  $\mathcal{X}$  with feature map  $\phi_j : \mathcal{X} \rightarrow \mathbb{R}^q$ ,  $j = 1, 2$ . In each part below, find a simple feature map for the kernel  $k$  in terms of feature maps for the kernels  $k_j$ . By this means, give an interpretation for the new kernel  $k$ .

a)  $k(x, z) = k_1(x, z) + k_2(x, z)$

b)  $k(x, z) = k_1(x, z)k_2(x, z)$

c)  $k(x, z) = k_1(x, z) / \sqrt{k_1(x, x)k_2(z, z)}$

**Answer:**

**Exercise 3:** A binary labelled set of data in  $\mathbb{R}^2$  is used to learn a SVM using the homogeneous quadratic kernel. By writing the equation for the decision boundary in terms of a quadratic form, reason about the types of decision boundaries that are possible in  $\mathbb{R}^2$ . In each case, give a neat sketch.

**Answer:**

**Exercise 4:** Consider the definite integral

$$\int_0^\infty e^{-(ax^2 + \frac{b}{x^2})} dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} e^{-2\sqrt{ab}}$$

Show that it can be rewritten as

$$\frac{\alpha}{\sqrt{\pi}} \int_0^\infty e^{-\frac{s}{t^2}} e^{-(\frac{\alpha t}{2})^2} dt = e^{-\alpha\sqrt{s}}, \quad \alpha > 0$$

Now use the above results to show that  $k(x, z) = e^{-\alpha\|x-z\|_2}$  is a kernel.

**Answer:**

**Exercise 5:** Let  $a > 0$  and  $L_2[0, a]$  denote the set of real valued square integrable functions on the interval  $[0, a]$ .  $L_2[0, a]$  is a Hilbert space under the inner product  $\langle g, h \rangle = \int_0^a g(s)h(s)ds$ . For  $f \in L_2[0, a]$ , let  $g(t) = \int_0^t f^2(s)ds$  and  $h(t) = \int_t^a f^2(s)ds$ , where  $t \in [0, a]$ . Show that for each  $a > 0$ ,

- a)  $k(x, z) = \min(x, z)$  is a kernel on  $[0, a]$ .
- b)  $k(x, z) = a - \max(x, z)$  is a kernel on  $[0, a]$ .
- c)  $k(x, z) = e^{-(\max(x, z) - \min(x, z))}$  is a kernel on  $[0, a]$ . Plot the function  $\max(x, z) - \min(x, z)$  and use this to simplify the result further.
- d) for each  $a > 0$ , and  $\gamma \geq 0$ ,  $k(x, z) = e^{-\gamma|x-z|}$  is a kernel on  $[-a, a]$ .
- e) for each  $\gamma \geq 0$ ,  $k(x, z) = e^{-\gamma|x-z|}$  is a kernel on  $\mathbb{R}$ .

**Answer:**