

ELE 535: Machine Learning and Pattern
Recognition
Homework 8

Zachary Hervieux-Moore

Friday 7th December, 2018

Exercise 1: Let $\{(x_i, y_i)\}_{i=1}^m$ with $x_i \in \mathbb{R}^n$ and $y_i \in \{\pm 1\}$, $i \in [1 : m]$, be a linearly separable set of training data. Show that if C is sufficiently large, the solution of the primal SVM problem will give the unique maximum margin separating hyperplane. How large does C need to be?

Answer:

Exercise 2: Let $\{(x_i, y_i)\}_{i=1}^m$ with $x_i \in \mathbb{R}^n$ and $y_i \in \{\pm 1\}$, $i \in [1 : m]$ be a training dataset. For a fixed value of C , let the corresponding SVM classifier have parameters w^* , b^* .

- a) Let $h \in \mathbb{R}^n$ and $Q \in \mathcal{O}_n$, and form the second training set: $\{(Q(x_i - h), y_i)\}_{i=1}^m$. Show that the SVM classifier for this second dataset using the same value of C has parameters Qw^* , $w^{*T}h + b^*$.
- b) If we first center the training examples, how does this change the SVM classifier?

Answer:

Exercise 3: Give a clear and concise derivation of the dual of the primal linear SVM problem shown below and explain the origin of each of the constraints in the dual problem.

$$\begin{aligned} \min_{w \in \mathbb{R}^n, b \in \mathbb{R}, s \in \mathbb{R}^m} \quad & \frac{1}{2} w^T w + C \mathbf{1}^T s \\ \text{s.t.} \quad & Z^T w + b y + s - \mathbf{1} \geq \mathbf{0} \\ & s \geq \mathbf{0} \end{aligned}$$

Answer:

Exercise 4: Suppose that instead of using $C \sum_{i=1}^m s_i$ as the penalty term in the objective of the primal SVM problem we use the quadratic penalty $\frac{1}{2}C \sum_{i=1}^m s_i^2$, while maintaining the constraint $s_i \geq 0$.

- a) Formulate the new primal problem in vector form. When is the primal problem feasible?
- b) Does strong duality hold for this problem? Justify your answer.
- c) Write down the KKT conditions.
- d) Find the dual problem.

Answer:

Exercise 5: You are provided with $m > 1$ data points $\{x_j \in \mathbb{R}^n\}_{j=1}^m$ of which at least d , with $1 < d \leq m$ are distinct. Let $X = [x_1, \dots, x_m]$ and consider the one class SVM problem:

$$\begin{aligned} \min_{R \in \mathbb{R}, a \in \mathbb{R}^n, s \in \mathbb{R}^m} \quad & R^2 + C \mathbf{1}^T s \\ \text{s.t.} \quad & \|x_i - a\|_2^2 \leq R^2 + s_i, \quad i = 1, \dots, m \\ & s \geq \mathbf{0} \end{aligned}$$

- a) Show that this is a feasible convex program and that strong duality holds. [Hint: let $r = R^2$]
- b) Write down the KKT conditions.
- c) Show that $\alpha^* \neq \mathbf{0}$ and that if $C > 1/(d-1)$ then $(R^2)^* > 0$ (harder).
- d) What are the support vectors for this problem?
- e) Derive the dual problem.
- f) Assume $C > 1/(d-1)$. Given the dual solution, how should a and R^2 be selected?

Answer: