

Epigenetic drugs: reducing the transgenerational effect of maternal dietary exposures on offspring mammary cancer risk

Helen S. Jung¹, Nguyen Nguyen², Sonia de Assis², Carlos Benitez², Idalia Cruz², and Leena Hilakivi-Clarke²
Haverford College¹; and Lombardi Comprehensive Cancer Center, Georgetown University²

ABSTRACT

Maternal dietary exposures to high levels of polyunsaturated fatty acids (PUFA) adversely change the epigenome of subsequent generations of offspring. Previous studies in our laboratory discovered that due to their in utero high-fat exposure, offspring have higher risks and more aggressive cases of mammary cancer, **as the diet increases** both F1 and F3 generations' tumor incidence (F1: $p < 0.0382$), burden (F1: $p < 0.0001$, F3: $p < 0.0001$) and malignancy; **we** hypothesized that the observed results were **regulated** by epigenetic mechanisms. The apparent permanence of the transgenerational effects of the diet, however, can be reversed through the use of valproic acid, a histone deacetylase (HDAC) (1.2 g/kg), and hydralazine a DNA methyltransferase (DNMT) (5 mg/kg) inhibitors, given via drinking water. F1 and F3 lineage that were in utero-exposed to F0 maternal high-fat diet were treated with these inhibitors. While **we** support **earlier studies** that presented **a worse** mammary cancer prognosis among control-diet mice following the use of epigenetic drugs, our findings reveal that offspring of high-fat female mice significantly benefited from daily epigenetic drug treatments that began before mammary tumorigenesis (F1 incidence: $p < 0.1024$, burden: $p < 0.0082$). Such encouraging prognoses may ultimately lead to the translation of the drug combination into **clinical** cancer prevention.

BACKGROUND

Breast cancer is the second leading cause of cancer death in American women, and about 1 in 8 will develop invasive breast cancer during their lifetime¹. While effective cancer therapies are on the rise, currently there are endeavors to progress cancer prevention. Environmental and lifestyle factors, such as everyday exposures to chemicals, physical activity, and diet, have been discovered to strongly correlate with cancer incidence and **aggression**.

It was previously studied that maternal dietary exposures influence the risk of breast cancer in subsequent generations. In particular, there is significant evidence that a maternal **PUFA-rich diet, which henceforth will be identified as generation F0**, even increases breast cancer risk for F3 offspring², alluding to a certain permanence of the diet's effects. These findings thus concluded that **the high-fat diet** affected the offspring epigenome through changes in methylation and acetylation patterns. Considering these epigenetic changes, we investigated possible measures to alleviate the adverse effects.

Epigenetic therapy has been recently recognized as an effective method of treatment for various cancers. Combinations of epigenetic-targeted drugs, such as the HDAC inhibitor Valproic acid and DNMT inhibitor Hydralazine³, are traditionally used as a **final** treatment option for cancer patients⁴. However, our study demonstrates that HDAC and DNMT inhibitors may assume roles beyond curative measures. In the form of daily preventive medication, epigenetic drugs may be the means to reverse maternal transgenerational effects on offspring mammary cancer risk.

Valproate	Side effects
Options: Show MedDRA Preferred Terms	Options: Show MedDRA Preferred Terms
Side effect	Data for drug
Athralgia	common
Joint swelling	common
Constipation	common
Parosmia	very rare
Myalgia	common
Dysphagia	common
Palpitations	common, very common
Anorexia	common, common
Angina Pectoris	common
Nausea	common
Vomiting	common
Flushing	common
Abnormal digestive tract function	common
Hypotension	common
Tachycardia	common, very common

Side effect	Data for drug
Nausea	8% - 48%
Headache	21% - 39%
Asthenia	postmarketing, 0% - 27%
Tremor	2% - 57%
Somnolence	1% - 31%
Fatigue	common, 4% - 19%
Diplopia	16%
Vomiting	5% - 27%
Communicable Diseases	12% - 20%
Diarrhea	7% - 23%
Dizziness	12% - 25%
Thrombocytopenia	1% - 24%
Influenza	12%
Sleeplessness	9% - 15%
Pain	common, 1% - 23%

Color scheme: standard - conservative

Frequency (1% to 100%)

Infrequent (0.1% to 1%)

Rare (< 0.1%)

Postmarketing

No frequency information

Not found on label

Most common side effects of HDAC inhibitor valproic acid (valproate) and DNMT inhibitor hydralazine.


METHODOLOGY

Maternal Exposure to High-Fat Diet

On gestation day (GD) 10, while one F0 group of pregnant C57BL/6NTac mice continued to be fed the control diet (6% corn oil), the F0 high-fat (HF) group was fed an isocaloric high n-6 polyunsaturated fat diet (18% corn oil) until GD 20. The HF lineage were only exposed to a dietary change during this time period.

Epigenetic Drug Treatment

On post-natal day (PND) 75, both control and HF mice were given valproic acid and hydralazine at a low dosage in their drinking water. They received constant exposure to the drug solution for about twenty weeks or until euthanasia is necessary. Each mouse consumed 1.2 g/kg of valproic acid and 5 mg/kg of hydralazine per day.



RESULTS





Figure 1. Maternal HF exposure increases tumor incidence and burden up to F3 generation. [A] F1 HF *in utero*-exposed female offspring had significantly higher mammary tumor incidence than F1 CON female offspring (log-rank test: $p < 0.0382$). [B] F3 groups began to have a **significant difference in tumor incidence** from week 12 until the end of the study period. [C],[D] F1 and F3 HF mice exhibited significantly greater tumor burden than control mice (paired t-test: F1 $p < 0.0001$, F3 $p < 0.0001$).

Figure 2. Maternal HF dietary exposure increases tumor malignancy for F1 – F3 offspring. [A] Using stained histopathology slides, 70% of tumors collected from F1 HF *in utero*-exposed offspring were identified as malignant, in comparison to 43% of CON offspring. [B] From F3 HF offspring, 69% of tumors collected are considered as malignant, in comparison to 44% of CON offspring.



RESULTS (cont.)




Figure 3. [A] Earlier studies indicated that in control mice, HDACi + DNMTi drugs do not inhibit tumor growth when compared to the vehicle, which was the mice group on the control diet. [B] Yet tumors of *in utero* estradiol-exposed mice significantly reduced in size after epigenetic drug treatment.




Figure 4. HDAC and DNMT inhibitors curb tumor incidence and burden for HF *in utero*-exposed offspring. For F1 HF offspring, [A] tumor incidence decreases ($p < 0.1024$) and [B] tumor burden also decreases ($p < 0.0082$).

DISCUSSION / CONCLUSION

- Our findings support the hypothesis that *in utero* exposure to a high-fat diet during critical periods of embryonic development can increase breast cancer risk transgenerationally, as tumors with a greater level of **aggression proliferated in numbers**.
- In pre-clinical rodent models, mammary cancer risk in *in utero*-exposed offspring decreases to normal levels when the mice consume epigenetic drugs as preventive medication. Valproic acid and hydralazine effectively reversed this transgenerational change.
- Because the two drugs in solution are low in toxicity, are FDA-approved—thus widely accessible—and cause a few side effects, they have a promising future in breast cancer prevention.
- For the successful transition of epigenetic drugs into clinical preventive practice, we must identify and characterize biomarkers that would appropriately indicate at-risk individuals. Following these studies can proper, timely methods of breast cancer prevention be administered.

ACKNOWLEDGEMENTS

This work was supported by the National Cancer Institute R01 grant. I thank Dr. Leena Hilakivi-Clarke, Ph.D. candidate Nguyen Nguyen, and the rest of the Clarke labs for their guidance and support. I also thank the Haverford College Koshland Integrated Science Center for my summer research funding.

REFERENCES

- American Cancer Society. What are the key statistics about breast cancer? <http://www.cancer.org/cancer/breastcancer/detailedguide/breast-cancer-key-statistics> (accessed Aug 12, 2015).
- de Assis, S.; Hilakivi-Clarke, L. *et al.* High-fat or ethinyl-oestradiol intake during pregnancy increases mammary cancer risk in several generations offspring. *Nat Commun.* **2012**, *3*, 1-9.
- SIDER Side Effect Resource. Side effect information for Valproate and Hydralazine. <http://sideeffects.embl.de/drugs/3121/> (accessed Aug 12, 2015).
- Bauman, J. *et al.* A Phase I Protocol of Hydralazine and Valproic Acid in Advanced, Previously Treated Solid Cancers. *Trans. Oncol.* **2014**, *7*, 349-354.