

Theorema 2.0: A First Tour

NB reached List reached2 CellGroupData reached List reached2 NullCell
reached

We consider “proving”, “computing”, and “solving” as the three basic mathematical activities.

CellGroupData reached List reached2

1 Proving

We want to prove

$$(\forall x, (P(x) \vee Q(x))) \wedge (\forall y, (P(y) \Rightarrow Q(y))) \Leftrightarrow (\forall x, Q(x)).$$

To prove a formula like the above, we need to enter it in the context of a Theorema environment.

Cell reached CellGroupData reached List reached2 Cell reached Cell reached
Cell reached Cell reached CellGroupData reached List reached2 Cell reached
Cell reached Cell reached Cell reached Cell reached Cell reached Cell reached
Cell reached Cell reached Cell reached Cell reached CellGroupData reached List
reached2

2 Computing

Cell reached CellGroupData reached List reached2 Cell reached Cell reached
Cell reached Cell reached Cell reached Cell reached CellGroupData reached
List reached2 Cell reached Cell reached CellGroupData reached List reached2
Cell reached Cell reached Cell reached Cell reached Cell reached Cell reached
Cell reached Cell reached Cell reached Cell reached CellGroupData reached List
reached2

3 Set Theory

Cell reached CellGroupData reached List reached2 Cell reached Cell reached
Cell reached Cell reached Cell reached Cell reached CellGroupData reached List
reached2 Cell reached Cell reached Cell reached Cell reached CellGroupData
reached List reached2 Cell reached Cell reached CellGroupData reached List
reached2 Cell reached Cell reached