On Ordowski's characterization of odd primes by the number of squares modulo n (OEIS A000224): even numbers, prime powers and semiprimes, formalized in Lean 4

Hetark Shah, AI-assisted by Grok

October 12, 2026

Abstract

Let $a(n)$ be the number of squares modulo n (OEIS A000224 [1]). T. Ordowski conjectured that for $n > 1$, $n^2 \equiv 1 \pmod{a(n)(a(n)-1)}$ holds if and only if n is an odd prime. **The full conjecture remains open.** We prove it for every $n > 1$ that is even, a prime power p^k ($k \geq 1$), or a product pq of two distinct odd primes. The prime-power case uses Stangl's formula for $a(p^k)$ [3], which we also formalize. The semiprime case combines a size bound with a divisibility bound and a finite check. Everything is formalized in Lean 4 with Mathlib against the exact definition of a in Google DeepMind's `formal-conjectures` repository [2]. The development uses no `sorry` and no `native_decide`, and only the standard axioms.

1 Introduction

For $n \geq 1$ let $a(n) = \#\{k^2 \bmod n : 0 \leq k < n\}$.

Conjecture 1 (Ordowski, 2025)**.** *For $n > 1$, n is an odd prime if and only if $n^2 \equiv 1 \pmod{a(n)(a(n)-1)}$.*

The conjecture appears as a comment on A000224 [1], together with M. Marcus's remark that it holds up to $n = 10^8$. It is formalized as `OeisA224.conjecture` in `FormalConjectures/OEIS/224.lean` of `formal-conjectures` [2], where it is tagged `research open`. The forward direction is immediate from $a(p) = (p+1)/2$. The open part is to show that no composite n satisfies the congruence.

2 Related work

The count $a(n)$ is multiplicative, and Stangl [3] gives its value at prime powers; see also Finch and Sebah [4]. On October 12, 2026 we searched OEIS, the web, MathOverflow and the issues and pull requests of `formal-conjectures` (upstream commit `34ee0e3`). We found no proof of Conjecture 1, partial or complete. The cases treated here therefore appear to be new, although the arguments are elementary. The formalization uses Lean 4 [6] and Mathlib [5].

3 Statement

Theorem 2. *Conjecture 1 holds for every $n > 1$ that is (i) even, (ii) a prime power p^k with $k \geq 1$, or (iii) a product pq of distinct odd primes p, q.*

Cases not covered include squarefree n with three or more odd prime factors, and odd n such as p^2q that have a repeated prime factor and at least two distinct primes.

4 Proof overview

4.1 Preliminaries

Lemma 3. *$a(n) = \#\{x^2 : x \in \mathbb{Z}/n\}$; a is multiplicative; and $a(p) = (p+1)/2$ for odd primes p.*

Proof. The first claim holds because reduction mod n is a bijection $\{0, \dots, n-1\} \to \mathbb{Z}/n$. If $\gcd(m, n) = 1$, the CRT ring isomorphism $\mathbb{Z}/mn \cong \mathbb{Z}/m \times \mathbb{Z}/n$ commutes with squaring, so it maps the set of squares onto the product of the two sets of squares. For an odd prime p, the squares are the values k^2 with $0 \le k \le (p-1)/2$, since $x^2 = (-x)^2$. These values are distinct: $k^2 = l^2$ forces $p \mid (k-l)(k+l)$, and $0 \le k + l < p$. □

4.2 The even case

If n is even, then $n^2 - 1$ is odd, while $a(n)(a(n) - 1)$ is a product of two consecutive integers and hence even. So $n^2 \not\equiv 1$. (If $a(n)(a(n)-1) = 0$, the congruence would say $n^2 = 1$, which is impossible for $n > 1$.)

4.3 The semiprime case

Theorem 4. *Let $p < q$ be odd integers with $p \ge 3$ prime, and let $s = (p+1)(q+1)/4$. Then $s(s-1) \nmid (pq)^2 - 1$.*

Primality of q is not needed. By Lemma 3, $a(pq) = s$ for distinct odd primes, so Theorem 4 gives the semiprime case of Theorem 2.

Proof. Put $P = (p+1)(q+1)$, so that $16\, s(s-1) = P(P-4)$. Suppose $Q\, s(s-1) = (pq)^2 - 1$, so $Q\, P(P-4) = 16((pq)^2 - 1)$.

Step 1: $Q \le 15$. Expanding, $P(P-4) - ((pq)^2 - 1) = 2pq(p+q+1) - 4pq + (p+q+1)^2 - 4(p+q+1) + 1 > 0$. Hence $Q < 16$.

Step 2: $p \le 61$. By Step 1, $16((pq)^2 - 1) \le 15P(P-4) < 15P^2$. Suppose $p \ge 62$. Since $q > p$ we have $(q+1)p \le (p+1)q$, so $P^2p^2 \le (p+1)^4q^2$. Also $15(p+1)^4 \le 16p^4 - 16$ for $p \ge 62$. Together with $q \ge p$ this gives $15P^2 \le 16((pq)^2 - 1)$, a contradiction.

Step 3: $q+1 \mid 16(p^2-1)$. Since $q+1 \mid P$, we get $q+1 \mid 16((pq)^2 - 1)$. Also $16((pq)^2 - 1) - 16(p^2-1) = 16p^2(q^2-1)$ is divisible by $q+1$.

Step 4: finite check. For each prime $p \in [3, 61]$ and each divisor $d = q+1$ of $16(p^2-1)$ with $q > p$ odd, direct computation shows $s(s-1) \nmid (pq)^2 - 1$. There are only finitely many such pairs. □

4.4 Odd prime powers

Lemma 5 (Stangl [3])**.** *For an odd prime p and $k \ge 2$, $a(p^k) = a(p^{k-2}) + \frac{1}{2}p^{k-1}(p-1)$. Consequently $2(p+1)\, a(p^k) = p^{k+1} + p + 2$ if k is even and $p^{k+1} + 2p + 1$ if k is odd.*

Proof. Let $N = p^k$. A square x^2 mod N is divisible by p iff $p \mid x$. Writing $x = pt$ gives $(pt)^2$ mod $p^2M = p^2(t^2 \bmod M)$ with $M = p^{k-2}$, so the squares divisible by p are exactly p^2 times the squares modulo M. For the squares prime to p: if $x, y \in [0, N/2)$ are prime to p and $N \mid (y-x)(y+x)$, then p cannot divide both factors (as p is odd and $p \nmid y$), so N divides one factor, forcing $x = y$.

Replacing x by $N-x$, every unit square has such a representative. Hence the unit squares number $\varphi(N)/2$. The closed form follows by induction from $a(1)=1$ and $a(p)=(p+1)/2$. □

Theorem 6. *If p is an odd prime, $k \geq 2$, and s satisfies the identity of Lemma 5, then $s(s-1) \nmid p^{2k}-1$. Consequently, $(p^k)^2 \not\equiv 1 \pmod{a(p^k)(a(p^k)-1)}$, and Conjecture 1 holds for all odd prime powers.*

Sketch. Let $X = p^k$ and $M = 2(p+1)$, and write $Q\,s(s-1) = X^2-1$. Multiplying by M^2 gives $Q(Ms)(Ms-M) = M^2(X^2-1)$. For even k, reducing modulo p gives $p \mid 4(p+1)^2$, which is impossible. For odd $k \geq 3$, a size estimate gives $1 \leq Q \leq 7$, and reducing modulo p gives $p \mid Q-4$. If $Q=4$, the equation becomes $(2p+1)X^2 = 2p^2X+p^2$, which is impossible since $X > p$. Otherwise $p=3$ and $Q \in \{1,7\}$, and both cases are ruled out by elementary inequalities. □

5 Formalization

All declarations are in namespace `OeisA224`. The definition `a` in `Defs.lean` is copied verbatim from `formal-conjectures`. The theorem `conjecture_main` concludes exactly `(n.Prime ∧ n ≠ 2) ↔ n ^ 2 ≡ 1 [MOD a n * (a n - 1)]` under its case hypothesis.

Lean name (file)	Statement
`a_eq_aZ`, `aZ_mul`, `aZ_prime` (`Count.lean`)	Lemma 3
`even_case`, `prime_direction` (`Basic.lean`)	even case; odd prime ⇒ congruence
`finCheck_true` (`FiniteCheck.lean`)	Step 4, checked with `decide +kernel`
`pq_reduction`, `pq_case`, `semiprime_case` (`Semiprime.lean`)	Theorem 4 and case (iii)
`pp_even`, `pp_odd` (`PrimePowerCore.lean`)	arithmetic core of Theorem 6
`stangl`, `prime_power_case` (`PrimePower.lean`)	Lemma 5; Theorem 6
`conjecture_of_even_or_prime_or_semiprime`, `conjecture_main` (`Main.lean`)	Theorem 2

Table 1: Theorems and their Lean names.

6 Reproducibility

- Source code: `https://github.com/het4rk/ordowski-a224`.
- Toolchain: `leanprover/lean4:v4.35.0-rc4`. Mathlib: tag `v4.35.0-rc4`, commit `1f414401f69059aa7eead47b53ee40bd38455eeb`, pinned in `lake-manifest.json`.
- Build:

```
lake exe cache get
lake build
lake env lean Axioms.lean
```

`FiniteCheck.lean` takes a few minutes and several GB of RAM.

- Axioms: all 12 declarations listed in `Axioms.lean`, including `conjecture_main`, depend only on `propext`, `Classical.choice` and `Quot.sound`.
- Attestation: `verification/` contains the fresh-clone build log, the axiom output and `VERIFICATION.md`, which records the commit, the toolchain and SHA-256 hashes of the Lean sources. CI (`https://github.com/het4rk/ordowski-a224/actions/workflows/lean.yml`) rebuilds the project on every push. It rejects `sorry` and non-whitelisted axioms, and uploads the build log and axiom report as artifacts.

7 Limitations

The full conjecture remains open. We do not treat squarefree n with three or more prime factors, or general odd composites with a repeated factor and two or more distinct primes (for example p^2q). One part of the method extends but is not formalized: the argument of Step 3 shows that if $n = mq$ satisfies the congruence, with q prime and $q \nmid m$, then $q + 1 \mid 4(m^2 - 1)$. The size argument of Step 2, however, does not bound the second-smallest prime once there are three or more factors, so the method does not directly extend. The independent computational checks in `scripts/` are not part of the formal proof.

References

[1] OEIS Foundation Inc., *A000224: Number of squares mod n*, `https://oeis.org/A000224`.

[2] Google DeepMind, *formal-conjectures*, `https://github.com/google-deepmind/formal-conjectures`, file `FormalConjectures/OEIS/224.lean`.

[3] W. D. Stangl, Counting squares in $\mathbb{Z}_n$, *Math. Mag.* **69** (1996), 285–289.

[4] S. R. Finch, P. Sebah, *Squares and cubes modulo n*, arXiv:math/0604465.

[5] The mathlib Community, *The Lean mathematical library*, in Proc. CPP 2020, ACM, 367–381.

[6] L. de Moura, S. Ullrich, *The Lean 4 theorem prover and programming language*, in CADE-28, LNCS 12699, Springer, 2021, 625–635.