

## UNIT 3

### Boolean Algebra (continued)

***This chapter includes:***

- 3.1 Multiplying out and Factoring Expressions
- 3.2 Exclusive-OR and Equivalence Operations
- 3.3 The Consensus Theorem
- 3.4 Algebraic Simplification of Switching Expressions
- 3.5 Proving Validity of an Equation

# Learning Objectives

1. Apply the laws and theorems of Boolean algebra to the manipulation of algebraic expressions to simplifying an expression, finding the complement of an expression and multiplying out and factoring an expression.
2. Prove any of the theorems using a truth table or give an algebraic proof.
3. Define the exclusive-OR and equivalence operations. State, prove, and use the basic theorems that concern these operations.
4. Use the consensus theorem to delete terms from and add terms to a switching algebra expression.
5. Given an equation, prove algebraically that it is valid or show that it is not valid.

# Multiplying out and Factoring Expressions

Given an expression in product-of-sums form, the corresponding sum-of-products expression can be obtained by multiplying out, using the two distributive laws:

$$X(Y + Z) = XY + XZ \quad (3-1)$$

$$(X + Y)(X + Z) = X + YZ \quad (3-2)$$

In addition, the following theorem is very useful for factoring and multiplying out:

$$(X + \overline{Y})(X' + Z) = XZ + X'Y \quad (3-3)$$

# Multiplying out and Factoring Expressions

- ❖ In general, when we multiply out an expression, we should use (3-3) along with (3-1) and (3-2).
- ❖ To avoid generating unnecessary terms when multiplying out, (3-2) and (3-3) should generally be applied before (3-1), and terms should be grouped to expedite their application.

# Multiplying out and Factoring Expressions

## Example 1:

### Example

$$\begin{aligned}
 & (A + B + C')(A + B + D)(A + B + E)(\overline{A + D' + E})(A' + C) \\
 &= (\overbrace{A + B + C'D}^{\downarrow})(\overbrace{A + B + E})[AC + A'(D' + E)] \\
 &= (\overbrace{A + B + C'DE}^{\downarrow})(AC + A'D' + A'E) \\
 &= AC + \cancel{ABC} + A'BD' + A'BE + A'C'DE \qquad (3-4)
 \end{aligned}$$

What theorem was used to eliminate  $ABC$ ? (*Hint: let  $X = AC$ .*)

In this example, if the ordinary distributive law (3-1) had been used to multiply out the expression by brute force, 162 terms would have been generated, and 158 of these terms would then have to be eliminated.

# Multiplying out and Factoring Expressions

## Example 2:

❖ By repeatedly applying (3-1), (3-2), and (3-3), any expression can be converted to a product-of-sums form.

### Example of Factoring

$$\begin{aligned}
 & AC + A'BD' + A'BE + A'C'DE \\
 &= \underbrace{AC}_{XZ} + A' \underbrace{(BD' + BE + C'DE)}_{Y'} \\
 &= (A + BD' + BE + C'DE)(A' + C) \\
 &= [\underbrace{A + C'DE}_X + B \underbrace{(D' + E)}_{YZ}](A' + C)
 \end{aligned}$$

# Multiplying out and Factoring Expressions

## Example 2 (continued):

$$\begin{aligned}
 &= (A + B + C'DE)(A + \cancel{C'DE} + D' + E)(A' + C) \\
 &= (A + B + C')(A + B + D)(A + B + E)(A + D' + E)(A' + C) \quad (3-5)
 \end{aligned}$$

This is the same expression we started with in (3-4).



# Exclusive-OR and Equivalence Operations

## Exclusive-OR (XOR) Operation:

(a)

$X$	$Y$	$X \oplus Y$
0	0	0
0	1	1
1	0	1
1	1	0

(b)

$$\begin{array}{ll}
 0 \oplus 0 = 0 & 0 \oplus 1 = 1 \\
 1 \oplus 0 = 1 & 1 \oplus 1 = 0
 \end{array}$$



From left to right, (a) XOR truth table, (b) XOR definition, (c) XOR logic symbol.

# Exclusive-OR and Equivalence Operations

- ❖ For **exclusive-OR**,  $X \oplus Y = 1$  if and only if (iff)  $X = 1$  or  $Y = 1$ , but not both.
- ❖ The ordinary OR operation is sometimes called **inclusive-OR** because  $X + Y = 1$  iff  $X = 1$  or  $Y = 1$ , or both.
- ❖ Exclusive OR can be expressed in terms of AND and OR. Because  $X \oplus Y = 1$  iff  $X$  is 0 and  $Y$  is 1 or  $X$  is 1 and  $Y$  is 0, we can write
$$X \oplus Y = XY + X\bar{Y} \quad (3-6)$$
- ❖ Derivation of (3-6) is on page 68.

# Exclusive-OR and Equivalence Operations

## Theorems for Exclusive-OR:

$$X \oplus 0 = X \quad (3-8)$$

$$X \oplus 1 = X' \quad (3-9)$$

$$X \oplus X = 0 \quad (3-10)$$

$$X \oplus X' = 1 \quad (3-11)$$

$$X \oplus Y = Y \oplus X \text{ (commutative law)} \quad (3-12)$$

$$(X \oplus Y) \oplus Z = X \oplus (Y \oplus Z) = X \oplus Y \oplus Z \text{ (associative law)} \quad (3-13)$$

$$X(Y \oplus Z) = XY \oplus XZ \text{ (distributive law)} \quad (3-14)$$



$$(X \oplus Y)' = X \oplus Y' = X' \oplus Y = XY + X'Y' \quad (3-15)$$

# Exclusive-OR and Equivalence Operations

## Equivalence Operation:

(a)

$X$	$Y$	$X \equiv Y$
0	0	1
0	1	0
1	0	0
1	1	1



(b)

$$\begin{array}{ll} (0 \equiv 0) = 1 & (0 \equiv 1) = 0 \\ (1 \equiv 0) = 0 & (1 \equiv 1) = 1 \end{array}$$

From left to right, (a) Equivalence operation truth table, (b) Definition, (c) Logic symbols, equivalence and exclusive-NOR gates

# Exclusive-OR and Equivalence Operations

## Equivalence Operations:

❖ For this operation, the output will be 1 iff X and Y have the same value. So,

$$(X \equiv Y) = XY + X'Y' \quad (3-17)$$

❖ Equivalence is the complement of the exclusive-OR operation.

$$\begin{aligned} (X \oplus Y)' &= (X'Y + XY')' = (X + Y')(X' + Y) \\ &= XY + X'Y' = (X \equiv Y) \end{aligned} \quad (3-18)$$

# The Consensus Theorem

## Consensus Theorem:

❖ The **consensus theorem** is used to cancel out redundant terms in an expression and is stated below:

$$XY + X'Z + YZ = XY + X'Z \quad (3-20)$$

❖ The term that is eliminated can be called the **consensus term**.

❖ The dual form of the consensus theorem is:

$$(X + Y)(X' + Z)(Y + Z) = (X + Y)(X' + Z) \quad (3-21)$$

# The Consensus Theorem

## Example 3:

### Example

$$A'C'D + A'BD + \cancel{BCD} + ABC + ACD' \quad (3-22)$$

First, we eliminate  $BCD$  as shown. (Why can it be eliminated?)

Now that  $BCD$  has been eliminated, it is no longer there, and it *cannot* be used to eliminate another term. Checking all pairs of terms shows that no additional terms can be eliminated by the consensus theorem.

Now we start over again:

$$A'C'D + \cancel{A'BD} + BCD + \cancel{ABC} + ACD' \quad (3-23)$$

This time, we do not eliminate  $BCD$ ; instead we eliminate two other terms by the consensus theorem. After doing this, observe that  $BCD$  can no longer be eliminated. Note that the expression reduces to four terms if  $BCD$  is eliminated first, but that it can be reduced to three terms if  $BCD$  is not eliminated.

# The Consensus Theorem

## Example 3 (continued):

Sometimes it is impossible to directly reduce an expression to a minimum number of terms by simply eliminating terms. It may be necessary to first add a term using the consensus theorem and *then* use the added term to eliminate other terms. For example, consider the expression

$$F = ABCD + B'CDE + A'B' + BCE'$$

If we compare every pair of terms to see if a consensus term can be formed, we find that the only consensus terms are  $ACDE$  (from  $ABCD$  and  $B'CDE$ ) and  $A'CE'$



# The Consensus Theorem

## Example 3 (continued):

(from  $A'B'$  and  $BCE'$ ). Because neither of these consensus terms appears in the original expression, we cannot directly eliminate any terms using the consensus theorem. However, if we first add the consensus term  $ACDE$  to  $F$ , we get

$$F = \overbrace{ABCD + B'CDE} + A'B' + BCE' + ACDE$$

Then, we can eliminate  $ABCD$  and  $B'CDE$  using the consensus theorem, and  $F$  reduces to

$$F = A'B' + BCE' + ACDE$$

The term  $ACDE$  is no longer redundant and cannot be eliminated from the final expression.

# Algebraic Simplification of Switching Expressions

## **Ways to simplify switching expressions:**

❖ In addition to multiplying out and factoring, three basic ways of simplifying switching functions are:

- combining terms
- eliminating terms
- eliminating literals

# Algebraic Simplification of Switching Expressions

## Combining Terms:

1. *Combining terms.* Use the theorem  $XY + XY' = X$  to combine two terms. For example,

$$abc'd' + abcd' = abd' \quad [X = abd', Y = c] \quad (3-24)$$

When combining terms by this theorem, the two terms to be combined should contain exactly the same variables, and exactly one of the variables should appear complemented in one term and not in the other. Because  $X + X = X$ , a given term may be duplicated and combined with two or more other terms. For example,

$$ab'c + abc + a'bc = ab'c + abc + abc + a'bc = ac + bc$$

The theorem still can be used, of course, when  $X$  and  $Y$  are replaced with more complicated expressions. For example,

$$(a + bc)(d + e') + a'(b' + c')(d + e') = d + e'$$
$$[X = d + e', Y = a + bc, Y' = a'(b' + c')]$$

# Algebraic Simplification of Switching Expressions

## Eliminating Terms and Literals:

2. *Eliminating terms.* Use the theorem  $X + XY = X$  to eliminate redundant terms if possible; then try to apply the consensus theorem ( $XY + X'Z + YZ = XY + X'Z$ ) to eliminate any consensus terms. For example,

$$\begin{aligned} a'b + a'bc &= a'b & [X = a'b] \\ a'bc' + bcd + a'bd &= a'bc' + bcd & [X = c, Y = bd, Z = a'b] \end{aligned} \quad (3-25)$$

3. *Eliminating literals.* Use the theorem  $X + X'Y = X + Y$  to eliminate redundant literals. Simple factoring may be necessary before the theorem is applied.

# Algebraic Simplification of Switching Expressions

## Example 4:

Example

$$\begin{aligned}
 A'B + A'B'C'D' + ABCD' &= A'(B + B'C'D') + ABCD' \\
 &= A'(B + C'D') + ABCD' \\
 &= B(A' + ACD') + A'C'D' \\
 &= B(A' + CD') + A'C'D' \\
 &= A'B + BCD' + A'C'D' \quad (3-26)
 \end{aligned}$$

## Example 5- Adding Redundant Terms:

Example

$$\begin{aligned}
 WX + XY + X'Z' + WY'Z' & \quad (\text{add } WZ' \text{ by consensus theorem}) \\
 = WX + XY + X'Z' + WY'Z' + WZ' & \quad (\text{eliminate } WY'Z') \\
 = WX + XY + X'Z' + WZ' & \quad (\text{eliminate } WZ') \\
 = WX + XY + X'Z' & \quad (3-27)
 \end{aligned}$$

# Algebraic Simplification of Switching Expressions

## Example 6:

### Example

$$\begin{aligned}
 & \underbrace{A'B'C'D' + A'BC'D'}_{\textcircled{1} A'C'D'} + A'BD + \underbrace{A'BCD}_{\textcircled{2}} + ABCD + ACD' + B'CD' \\
 &= A'C'D' + BD(A' + AC) + ACD' + B'CD' \\
 & \quad \textcircled{3} \\
 &= A'C'D' + A'BD + \underbrace{BCD + ACD'}_{+ ABC \textcircled{4}} + B'CD'
 \end{aligned}$$

$$\begin{aligned}
 &= A'C'D' + \underbrace{A'BD + BCD + ACD' + B'CD' + ABC}_{\text{consensus } BCD} \\
 & \quad \text{consensus } ACD' \\
 &= A'C'D' + A'BD + B'CD' + ABC \quad (3-28)
 \end{aligned}$$

What theorems were used in steps 1, 2, 3, and 4?

# Algebraic Simplification of Switching Expressions

## Example 7:

❖ For a product-of-sums form instead of a sum-of-products form, the duals of the preceding theorems should be applied.

### Example

$$\begin{aligned}
 & \underbrace{(A' + B' + \cancel{C})(A' + B' + C)}_{\textcircled{1} (A' + B')} (B' + C)(A + C) \underbrace{(A + \cancel{B} + C)}_{\textcircled{2}} \\
 &= (A' + B') \underbrace{(B' + \cancel{C})}_{\textcircled{3}} (A + C) = (A' + B')(A + C) \quad (3-29)
 \end{aligned}$$

What theorems were used in steps 1, 2, and 3?

# Proving the Validity of an Equation

## **Proving an equation is valid:**

To determine if an equation is valid, meaning valid for all combinations of values of the variables, several methods can be used:

1. Construct a truth table and evaluate both sides of the equation for all combinations of values of the variables.
2. Manipulate one side of the equation by applying various theorems until it is identical with the other side.



# Proving the Validity of an Equation

## **Proving an Equation is valid (continued):**

3. Reduce both sides of the equation independently to the same expression.
4. It is permissible to perform the same operation on both sides of the equation provided that the operation is reversible.

# Proving the Validity of an Equation

## To prove an equation is **NOT** valid:

❖ To prove that an equation is *not valid*, it is sufficient to show *one combination* of values of the variables for which the two sides of the equation have different values.

1. First reduce both sides to a sum of products (or a product of sums).
2. Compare the two sides of the equation to see how they differ.
3. Then try to add terms to one side of the equation that are present on the other side.
4. Finally try to eliminate terms from one side that are not present on the other.

❖ Whatever method is used, frequently compare both sides of the equation and let the difference between them serve as a guide for what steps to take next.

# Proving the Validity of an Equation

## Example 8:

Show that

$$A'BD' + BCD + ABC' + AB'D = BC'D' + AD + A'BC$$

Starting with the left side, we first add consensus terms, then combine terms, and finally eliminate terms by the consensus theorem.

$$\begin{aligned}
 &A'BD' + BCD + ABC' + AB'D \\
 &= A'BD' + BCD + ABC' + AB'D + BC'D + A'BC + ABD \\
 &\quad \text{(add consensus of } A'BD' \text{ and } ABC') \quad \nearrow \\
 &\quad \text{(add consensus of } A'BD' \text{ and } BCD) \quad \nearrow \\
 &\quad \text{(add consensus of } BCD \text{ and } ABC') \quad \nearrow \\
 &= AD + A'BD' + BCD + ABC' + BC'D' + A'BC = BC'D' + AD + A'BC \\
 &\quad \nearrow \quad \nearrow \quad \nearrow \quad \text{(eliminate consensus of } BC'D' \text{ and } AD) \\
 &\quad \text{(eliminate consensus of } AD \text{ and } A'BC) \\
 &\quad \text{(eliminate consensus of } BC'D' \text{ and } A'BC) \quad (3-30)
 \end{aligned}$$

# Proving the Validity of an Equation

## Example 9:

Show that the following equation is valid:

$$\begin{aligned} A'BC'D + (A' + BC)(A + C'D') + BC'D + A'BC' \\ = ABCD + A'C'D' + ABD + ABCD' + BC'D \end{aligned}$$

First, we will reduce the left side:

$$\begin{aligned} & A'BC'D + (A' + BC)(A + C'D') + BC'D + A'BC' \\ & \hspace{20em} \text{(eliminate } A'BC'D \text{ using absorption)} \\ & = (A' + BC)(A + C'D') + BC'D + A'BC' \\ & \hspace{20em} \text{(multiply out using (3-3))} \\ & = ABC + A'C'D' + BC'D + A'BC' \\ & \hspace{20em} \text{(eliminate } A'BC' \text{ by consensus)} \\ & = ABC + A'C'D' + BC'D \end{aligned}$$

# Proving the Validity of an Equation

## Example 9 (continued):

Now we will reduce the right side:

$$= ABCD + A'C'D' + ABD + ABCD' + BC'D$$

(combine  $ABCD$  and  $ABCD'$ )

$$= ABC + A'C'D' + ABD + BC'D$$

(eliminate  $ABD$  by consensus)

$$= ABC + A'C'D' + BC'D$$

Because both sides of the original equation were independently reduced to the same expression, the original equation is valid.

# Proving the Validity of an Equation

## Cancellation Laws in Boolean Algebra:

The cancellation law for ordinary algebra, where:

$$\text{If } x+y=x+z \text{ then } y=z$$

Does not hold for Boolean algebra.

The cancellation law for multiplication, where:

$$\text{If } xy=xz \text{ then } y=z$$

Does not hold for Boolean algebra.

However, the converses of these hold true:

$$\text{If } y = z, \quad \text{then} \quad x + y = x + z \quad (3-33)$$

$$\text{If } y = z, \quad \text{then} \quad xy = xz \quad (3-34)$$