

Math 135 – Differential Equations

University of California, Los Angeles

Duc Vu

Fall 2021

This is math 135, officially known as Ordinary Differential Equations though we also delve into partial differential equations. It's taught by Professor Hester. We meet weekly on MWF from 12:00 pm to 12:50 pm for lecture. The main textbook used for the class is *Differential Equations with Applications and Historical Notes* 3rd by *Simmons*. Other course notes can be found at my [blog site](#). Please let me know through my [email](#) if you spot any concerning typos in the note.

Contents

1 Lec 1: Sep 27, 2021	3
1.1 Laplace Transforms	3
2 Lec 2: Sep 29, 2021	5
2.1 Laplace Transform (Cont'd)	5
3 Lec 3: Oct 1, 2021	7
3.1 Existence of Laplace Transform	7
4 Lec 4: Oct 4, 2021	9
4.1 Convolution	9
4.2 Application of Laplace Transform – Integral Equation	9
5 Lec 5: Oct 6, 2021	11
5.1 Dirac Delta “Function”	11
6 Lec 6: Oct 08, 2021	13
6.1 Existence & Uniqueness of ODE Solutions	13
7 Lec 7: Oct 11, 2021	14
7.1 Picard Iteration	14
8 Lec 8: Oct 13, 2021	16
8.1 Continuity	16
9 Lec 9: Oct 15, 2021	18
9.1 Picard’s Theorem	18
10 Lec 10: Oct 18, 2021	22
10.1 Fourier Series	22
11 Lec 11: Oct 20, 2021	24
11.1 Coefficients of Fourier Series	24

12 Lec 12: Oct 22, 2021	27
12.1 Convergence of Fourier Series	27
13 Lec 13: Oct 27, 2021	29
13.1 Complex Fourier Series	29
14 Lec 14: Oct 29, 2021	31
14.1 Rescaling Intervals of Fourier Series	31
15 Lec 15: Nov 1, 2021	32
15.1 The Relationship between Smoothness and Fourier Coefficients	32
16 Lec 16: Nov 3, 2021	34
16.1 Hilbert Spaces & Convergence in Norm	34
17 Lec 17: Nov 5, 2021	36
17.1 Pointwise Convergence of Fourier Series	36
18 Lec 18: Nov 8, 2021	38
18.1 Heat Equation	38
19 Lec 19: Nov 10, 2021	41
19.1 Wave Equation	41
20 Lec 20: Nov 12, 2021	43
20.1 Midterm 2	43
20.2 Laplace Equations	43

List of Theorems

4.1 Convolution	9
9.1 Picard	18
12.1 Fourier Convergence	27

List of Definitions

8.3 Uniform Continuity	16
10.1 Fourier Series	22
16.1 L^2 integrable	34
16.2 Hilbert Space	34
16.4 Convergence in Norm	34
16.5 Orthonormal System	34
16.6 General Fourier Series	35

§1 | Lec 1: Sep 27, 2021

§1.1 Laplace Transforms

Consider the following questions

1. What is a transform?
2. What is a Laplace transform?
3. What are some examples?
4. What are some general properties?
5. Why are they useful for differential equations?

Let's tackle these questions.

1. Notice that functions: sets \rightarrow sets. Transform is in higher hierarchy, i.e.,

Transform/Operator: functions \rightarrow functions

Example 1.1 • differentiation: $\frac{d}{dx} : f \mapsto f'$

- integration: $\int^x dx : f \mapsto \int^x f'(x)dx$
- multiplication by $g(x)$: $f(x) \rightarrow g(x)f(x)$
- shifting: $f(x) \rightarrow f(x - a)$

2. Laplace transform \mathcal{L}

$$\mathcal{L} : f(t) \mapsto F(s) = \int_0^\infty f(t)e^{-st} dt$$

where $f : [0, \infty) \rightarrow \mathbb{R}$ and $F : \mathbb{C} \rightarrow \mathbb{C}$

3. Examples:

Example 1.2 • $f(t) : t \mapsto 0 \implies \mathcal{L}[0] = 0$

- $f(t) = 1$

$$\begin{aligned} \mathcal{L}[1] &= \lim_{t \rightarrow \infty} \int_0^t e^{-st} dt \\ &= \lim_{t \rightarrow \infty} \left[\frac{e^{-st}}{-s} \right]_0^t \\ &= \lim_{t \rightarrow \infty} \left(\frac{e^{-st}}{-s} + \frac{1}{s} \right) \\ &= \frac{1}{s} \text{ if } \operatorname{Re}(s) > 0 \end{aligned}$$

Example 1.3 • Consider

$$\begin{aligned}\mathcal{L}[t] &= \int_0^\infty t e^{-st} dt \\ &= \left[\frac{t e^{-st}}{-s} \right]_0^\infty + \frac{1}{s} \int_0^\infty e^{-st} dt \\ &= \frac{1}{s^2} \text{ if } \operatorname{Re}(s) > 0\end{aligned}$$

We can generalize this as

$$\mathcal{L}[t^n] = \frac{1}{s^{n+1}}, \quad \operatorname{Re}(s) > 0, \quad n \in \mathbb{N}$$

In addition,

$$\begin{aligned}\mathcal{L}[e^{at}] &= \int_0^\infty e^{-(s-a)t} dt \\ &= \frac{1}{s-a}, \quad \operatorname{Re}(s) > a \\ \mathcal{L}[\cos \omega t] &= \frac{s}{s^2 + \omega^2} \\ \mathcal{L}[\sin \omega t] &= \frac{\omega}{s^2 + \omega^2}\end{aligned}$$

4. Properties:

a) Linear!

$$\begin{aligned}\mathcal{L}[f+g] &= \mathcal{L}[f] + \mathcal{L}[g] \\ \mathcal{L}[af] &= a\mathcal{L}[f]\end{aligned}$$

b) Consider:

$$\begin{aligned}\mathcal{L}[e^{at}f(t)] &= \int_0^\infty f(t)e^{-(s-a)t} dt \\ &= F(s-a) \quad \text{if } \operatorname{Re}(s-a) > 0\end{aligned}$$

Multiply an exponential in t -space $\xrightarrow{\mathcal{L}}$ shift in s -space.

5. In reverse,

$$\mathcal{L}[f(t-a)] = \int_0^\infty f(t-a)e^{-st} dt = \int_0^\infty f(t')e^{-st'} dt' e^{-sa}$$

where $t' = t - a$. So

$$\mathcal{L}[f(t-a)] = F(s)e^{-sa}$$

Thus, a shift in t -space $\xrightarrow{\mathcal{L}}$ multiply an exponential in s -space.

6. Differentiation:

$$\begin{aligned}\mathcal{L}[f'] &= \int_0^\infty f'(t)e^{-st} dt \\ &= [f e^{-st}]_0^\infty + \int_0^\infty f(t) s e^{-st} dt \\ &= sF(s) - f(0)\end{aligned}$$

§ 2 | Lec 2: Sep 29, 2021

§ 2.1 Laplace Transform (Cont'd)

Recap: $\mathcal{L} : f \rightarrow F$

$$\mathcal{L}[f(t)] = \int_0^{\infty} f(t)e^{-st} dt$$

where $t > 0$ and $s \in \mathbb{C}$.

Example 2.1 • $\mathcal{L}[t^n] = \frac{1}{s^{n+1}}, n \in \mathbb{N}$

• $\mathcal{L}[e^{at}] = \frac{1}{s-a}$

General properties of Laplace transform:

- linear
- shifting \leftrightarrow multiplying by exponential
- $\mathcal{L}[f'] = s\mathcal{L}[f] - f(0)$

Let's now use Laplace transform to solve the following ODE

$$f'' + af' + bf = g(t), \quad f(0) = f_0, \quad f'(0) = f'_0$$

Apply \mathcal{L} ,

$$\begin{aligned} \mathcal{L}[f'' + af' + bf] &= \mathcal{L}[g] \\ \mathcal{L}[f''] + a\mathcal{L}[f'] + b\mathcal{L}[f] &= G(s) \end{aligned}$$

Notice that

$$\mathcal{L}[f''] = s^2F - sf(0) - f'(0)$$

So

$$\begin{aligned} (s^2 + as + b)F(s) &= G(s) + (s + a)f_0 + f'_0 \\ F(s) &= \frac{G(s) + (s + a)f_0 + f'_0}{s^2 + as + b} \end{aligned}$$

To get $f(t)$ we need to invert \mathcal{L} .

Example 2.2

Consider:

$$f'' + 4f = 4t, \quad f(0) = 1, \quad f'(0) = 5$$

Apply \mathcal{L} , we get

$$\begin{aligned} (s^2 + 4)F(s) &= \frac{4}{s^2} + s + 5 \\ F(s) &= \frac{\frac{4}{s^2} + s + 5}{s^2 + 4} \\ &= \frac{4}{s^2(s^2 + 4)} + \frac{s}{s^2 + 4} + \frac{5}{s^2 + 4} \end{aligned}$$

Notice that we need to use partial fractions to decompose the first term.

$$\begin{aligned} \frac{4}{s^2(s^2 + 4)} &= \frac{A}{s^2} + \frac{B}{s^2 + 4} \\ 4 &= A(s^2 + 4) + Bs^2 \\ &= (A + B)s^2 + 4A \end{aligned}$$

So, $A = 1$, $B = -1$. Then,

$$\begin{aligned} F(s) &= \frac{1}{s^2} - \frac{1}{s^2 + 4} + \frac{s}{s^2 + 4} + \frac{5}{s^2 + 4} \\ &= \frac{1}{s^2} + \frac{4}{s^2 + 4} + \frac{s}{s^2 + 4} \\ \mathcal{L}[f] &= \mathcal{L}[t + 2\sin 2t + \cos 2t] \\ \implies f &= t + 2\sin 2t + \cos 2t \end{aligned}$$

§3 | Lec 3: Oct 1, 2021

§3.1 Existence of Laplace Transform

Question 3.1. When is Laplace transform is allowed? When does Laplace transform exist?

$$\mathcal{L}[f] = \int_0^{\infty} f(t)e^{-st} dt$$

Note: Beware of ∞ – only trust limits.

$$\mathcal{L}[f] = \lim_{\tau \rightarrow \infty} \int_0^{\tau} f(t)e^{-st} dt$$

Laplace transform exists when this limit exists?

$\lim_{\tau \rightarrow \infty} f^*(\tau)$ converges to $f_{\infty} \in \mathbb{R}$ if $\forall \varepsilon > 0, \exists M > 0$ s.t.

$$|f^*(\tau) - f_{\infty}| < \varepsilon \quad \text{for all } \tau > M$$

Convergence test for integrals:

$$\lim_{\tau \rightarrow \infty} \int_0^{\tau} f(t) dt$$

Comparison Test: If $|f(t)| < g(t)$ and $\int_0^{\infty} g(t) < \infty$ (converges) then

$$\int_0^{\infty} f(t) dt \leq \int_0^{\infty} |f(t)| dt \leq \int_0^{\infty} g(t) dt < \infty$$

i.e., $\int_0^{\infty} f(t) dt$ converges. Now, back to the Laplace transform

$$\mathcal{L}[f] = \int_0^{\infty} f(t)e^{-st} dt$$

What could break this integral?

1. fe^{-st} diverges/unbounded ($\lim_{t \rightarrow t^*} f(t) = \infty$)
2. fe^{-st} doesn't decay fast enough as $t \rightarrow \infty$.

What could prevent these issues?

1. Piecewise continuous: $\lim_{t \rightarrow t^-} f(t)$ and $\lim_{t \rightarrow t^+} f(t)$ exist.
2. Exponential order

$$|f(t)| < Me^{ct} \text{ for some } M > 0 \text{ \& } c$$

Have

$$\begin{aligned} c^{-t} &\leq 1 \cdot e^{-t} & \forall t > 0 \\ 1 &\leq 1 \cdot e^{0t} & \forall t > 0 \\ t &\leq 1 \cdot e^t & \forall t > 0 \end{aligned}$$

Theorem 3.1

If f is piecewise continuous and of exponential order c then $\mathcal{L}[f]$ exists for $s \in \mathbb{C}$ with $\operatorname{Re}(s) > c$.

Proof. Have

$$\begin{aligned}
 \mathcal{L}[f](s) &= \int_0^\infty f(t)e^{-st} dt \\
 \lim_{\tau \rightarrow \infty} \int_0^\tau f(t)e^{-st} dt &\leq \lim_{\tau \rightarrow \infty} \int_0^\tau |f(t)e^{-st}| dt \\
 &= \lim_{\tau \rightarrow \infty} \int_0^\tau |f(t)| e^{-s_r t} dt \\
 &\leq \lim_{\tau \rightarrow \infty} \int_0^\tau M e^{ct} \cdot e^{-s_r t} dt \\
 &= \lim_{\tau \rightarrow \infty} M \left[\frac{e^{(c-s_r)t}}{-(c-s_r)} \right]_0^\tau \\
 &= \frac{1}{s_r - c} \quad \text{if } s_r > c \\
 &< \infty
 \end{aligned}$$

Thus, $\mathcal{L}[f]$ exists (for $\text{Re}(s) > c$) by comparison test. □

This is a sufficient condition but not necessary.

Example 3.2

Consider the function $f(t) = \frac{1}{\sqrt{t}}$

$$\begin{aligned}
 \mathcal{L}\left[\frac{1}{t^{\frac{1}{2}}}\right] &= \int_0^\infty t^{-\frac{1}{2}} e^{-st} dt \\
 &= s^{-\frac{1}{2}} \int_0^\infty x^{-\frac{1}{2}} e^{-x} dx \\
 &= s^{-\frac{1}{2}} 2 \int_0^\infty e^{-z^2} dz \\
 &= \sqrt{\frac{\pi}{s}}
 \end{aligned}$$

However, we can see that $\frac{1}{t^{\frac{1}{2}}}$ isn't continuous on $[0, \infty)$.

§4 | Lec 4: Oct 4, 2021


§4.1 Convolution

Question 4.1. Can we invert $\mathcal{L}[f] \cdot \mathcal{L}[g]$?

We have

$$\begin{aligned} F(s)G(s) &= \int_0^\infty f(t)e^{-st} dt \int_0^\infty g(t')e^{-st'} dt' \\ &= \int_0^\infty \int_0^\infty f(t)g(t')e^{-s(t+t')} dt' dt \end{aligned}$$

Let's define $\tau = t + t' \implies d\tau = dt'$



$$\begin{aligned} F(s)G(s) &= \int_0^\infty \int_0^\infty f(t)g(t')e^{-s(t+t')} dt' dt \\ &= \int_0^\infty \int_0^\infty f(t)g(\tau - t)e^{-s\tau} d\tau dt \\ &= \int_0^\infty \left(\int_0^\tau f(t)g(\tau - t)e^{-s\tau} dt \right) d\tau \\ &= \int_0^\infty \left(\int_0^\tau f(t)g(\tau - t) dt \right) e^{-s\tau} d\tau \\ &= \mathcal{L} \left[\int_0^\tau f(t)g(\tau - t) dt \right] \end{aligned}$$

Theorem 4.1 (Convolution)

We have

$$\begin{aligned} (f * g)(\tau) &= \int_0^\tau f(t)g(\tau - t) dt \\ \mathcal{L}[f * g] &= \mathcal{L}[f] \cdot \mathcal{L}[g] \end{aligned}$$

§4.2 Application of Laplace Transform – Integral Equation

Consider:

$$f(\tau) = g(\tau) + \int_0^\tau k(\tau - t)f(t) dt$$

Notice

$$\begin{aligned}\mathbf{f} &= \mathbf{g} + K \cdot \mathbf{f} \\ f(\tau) &\approx f_i \\ g(\tau) &\approx g_i \\ k(\tau - t) &\approx K_{ij}\end{aligned}$$

Have

$$f = g + k * f$$

and we use Laplace

$$\begin{aligned}\mathcal{L}[f] &= \mathcal{L}[g] + \mathcal{L}[k] \cdot \mathcal{L}[f] \\ \mathcal{L}[f] &= \frac{\mathcal{L}[g]}{1 - \mathcal{L}[k]}\end{aligned}$$

Example 4.2

Consider $f(t) = t^3 + \int_0^t \sin(t - \tau)f(\tau)d\tau$.

$$F(s) = \frac{3!}{s^4} + \mathcal{L}[\sin t] F(s)$$

$$\vdots$$


$$F(s) = 3!(s^{-4} + s^{-6})$$

$$f(t) = t^3 + \frac{t^5}{20}$$

§5 | Lec 5: Oct 6, 2021

§5.1 Dirac Delta “Function”

Visually:



The limit of a function concentrated at zero, with integral

$$\int_{-\infty}^{\infty} \delta(t) dt = 1$$

Formally:

$$\delta : f(t) = \int_{-\infty}^{\infty} f(\tau) \delta(t - \tau) d\tau \implies f = f * \delta$$

δ “picks out” a pointwise value of any function we integrate against/convolve with. For finite dimension, let $\mathbf{f} \in \mathbb{R}^n$ and $\mathbf{e}_i = [0, \dots, 0, 1, 0, \dots]$. So

$$f_i = \mathbf{f} \cdot \mathbf{e}_i$$

For infinite dimension, $f(t) : \mathbb{R} \rightarrow \mathbb{R}$ for $t \in \mathbb{R}$,

$$f(t) = \int_{\mathbb{R}} f(\tau) \delta(t - \tau) d\tau$$

where $\delta(\tau - t) = \delta(t - \tau) = \delta_t(\tau)$. These two notions are analogous, in a sense. Solving a linear finite dimensional system

$$\mathbf{h} \in \mathbb{R}^n, \quad L \in \mathbb{R}^{n \times n}$$

Solve $L\mathbf{f} = \mathbf{h}$. If we know $L\mathbf{f}_i = \mathbf{e}_i$ where

\mathbf{e}_i : unit vector

\mathbf{f}_i : unit response vector

1. $\mathbf{h} = \sum h_i \mathbf{e}_i$

2. Linear superposition means

$$\mathbf{f} = \sum h_i \mathbf{f}_i$$

and

$$\begin{aligned} L\mathbf{f} &= L\left(\sum_i h_i \mathbf{f}_i\right) \\ &= \sum_i h_i L\mathbf{f}_i \\ &= \sum_i h_i \mathbf{e}_i \\ &= \mathbf{h} \end{aligned}$$

Solving ∞ -dim ODE

$$f'' + af' + bf = h(t) \quad (L[f] = h)$$

Let's say we know

$$g_t'' + ag_t' + bg = \delta_t$$

1. $h = h * \delta$
2. Then,

$$\begin{aligned} f &= h * g \\ &= \int_0^t g_t(\tau) h(\tau) d\tau \\ &= \int_0^t g(t - \tau) h(\tau) d\tau \end{aligned}$$

where g is known as the Green function.

$$\begin{aligned} e_i &\approx \delta_t \\ \mathbf{f}_i &\approx g_t \mathbf{f} = \sum h_i \mathbf{f}_i \approx f = h * g \end{aligned}$$


§6 | Lec 6: Oct 08, 2021

§6.1 Existence & Uniqueness of ODE Solutions

Intuitively, $f(t, x)$ is continuous seems like it guarantees a solution – **this is not true!**

1. Failure of existence over \mathbb{R} .


$$\frac{dx}{dt} = x^2, \quad x(0) = 1$$



We can easily solve this and obtain $x(t) = \frac{1}{1-t}$ which blows up in finite time.

2. What about uniqueness?

$$\frac{dx}{dt} = 3x^{\frac{2}{3}}, \quad x(0) = 0$$



This has infinite number of solution through $(0, 0)$ – non-unique. Notice that $x' = 3x^{\frac{2}{3}}$ is an autonomous ODE where the solution is $x(t) = t^3$. However, $x(t) = 0$ is also a solution which shows that solutions are not unique.

Question 6.1. What can prove existence and uniqueness?

1. Converting to “nicer” problem, $DE \iff$ integral equation
2. Devise an iterative algorithm to approximate solutions (Picard iteration)
3. Prove the algorithm converges to a unique solution

§7 | Lec 7: Oct 11, 2021

§7.1 Picard Iteration

Goal: Find sufficient conditions to prove existence and uniqueness of solution to ODE

$$\dot{x} = f(t, x(t)), \quad x(t_0) = x_0$$

Idea:

1. Smoother is better (integration is preferred over differentiation). Make things smoother by integrating

$$\dot{x}(t) = f(t, x(t)), \quad x(t_0) = x_0$$

Then, we can transform it into an integral equation

$$x(t) = x_0 + \int_{t_0}^t f(t', x(t')) dt'$$

Notice that f is continuous and x is continuous imply x is differentiable.

2. Iteration: If we can't solve it at first, try again.

Example 7.1

Newton's root-finding algorithm

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Picard Iteration: Iterative approximation to solutions of the integral equation

$$x(t) = x_0 + \int_{t_0}^t f(t', x(t')) dt'$$

Start with a guess for the function $x_0(t) = x_0$ (can be a constant)

$$x_{n+1}(t) = x_0 + \int_{t_0}^t f(t', x_n(t')) dt'$$

In general,

$$x_0(t) \xrightarrow{\text{Picard}} x_1(t) \xrightarrow{\text{Picard}} x_2(t) \xrightarrow{\text{Picard}} x_3(t) \xrightarrow{\infty} \dots$$

If $x_{n+1}(t) = x_n(t) = \bar{x}(t)$, then $\bar{x}(t)$ has to solve the IE. We want $\lim_{n \rightarrow \infty} x_n(t) \rightarrow x(t)$ solves IE.

Example 7.2

Consider $\dot{x}(t) = x(t)$, $x(0) = 1$. This is equivalent to the following integral equation

$$x(t) = 1 + \int_0^t x(t') dt'$$

Picard:

$$x_0(t) = 1$$

$$\begin{aligned} x_1(t) &= 1 + \int_0^t x_0(t') dt' = 1 + \int_0^t 1 dt' \\ &= 1 + t \end{aligned}$$

$$\begin{aligned} x_2(t) &= 1 + \int_0^t 1 + t dt \\ &= 1 + t + \frac{t^2}{2!} \end{aligned}$$

\vdots

$$x_n(t) = \sum_{k=0}^n \frac{t^k}{k!}$$

Thus,

$$\lim_{n \rightarrow \infty} x_n(t) \rightarrow e^t$$

§8 | Lec 8: Oct 13, 2021

§8.1 Continuity

Limit of continuous function is not necessarily continuous.

Example 8.1

Consider $x_n(t) = t^n$ on $[0, 1]$

$$x_0 = 1$$

$$x_1 = t$$

$$x_2 = t^2$$

$$\vdots$$

$$\bar{x} = \lim_{n \rightarrow \infty} x_n = \begin{cases} 0, & t < 1 \\ 1, & t = 1 \end{cases}$$

which is discontinuous.

Idea: We need “more” continuity. Given x , and given any $\varepsilon > 0$, if $|x - x'| < \delta(x, \varepsilon)$ then $|f(x) - f(x')| < \varepsilon$.

Example 8.2

Consider $f(x) = x$ on \mathbb{R} . We can see that

$$|x - x'| < \varepsilon \quad \forall |x - x'| < \varepsilon$$

in which we pick $\delta(x, \varepsilon) = \varepsilon$.

How about $f(x) = x^2$ on \mathbb{R} ?

$$|x^2 - y^2| < \varepsilon$$

If we pick $\delta(x, \varepsilon) = \varepsilon$, then $|x - y| < \delta = \varepsilon$ which does not necessarily imply $|x^2 - y^2| < \varepsilon$ because

$$\begin{aligned} |x^2 - y^2| &= |(x + y)(x - y)| \\ &= |x + y| |x - y| \\ &\leq \varepsilon |x + y| \end{aligned}$$

$|f(x) - f(y)| > \varepsilon$. So we need to pick smaller δ as x and y get larger. It would work for $\delta = \frac{\varepsilon}{2 \max(|x|, |y|)}$.

Question 8.1. Is $\frac{1}{x}$ continuous?

Ans: It depends on the domain. If we're talking about \mathbb{R} , it doesn't work at 0; on $(0, \infty)$, yes it's continuous.

Definition 8.3 (Uniform Continuity) — $\forall \varepsilon > 0, \exists \delta(\varepsilon) > 0$ s.t. $|x - y| < \delta \implies |f(x) - f(y)| < \varepsilon$.

Remark 8.4. Notice that the definition is similar to continuity except that δ doesn't depend on x .

Example 8.5

x^2 on \mathbb{R} is not uniformly continuous but x^2 on $(a, b) \subseteq \mathbb{R}$ is continuous since

$$\delta = \frac{\varepsilon}{\max(|x|, |y|)} = \frac{\varepsilon}{\max(|a|, |b|)}$$

Remark 8.6. Uniform continuity also depends on the domain as continuity does.

Exercise 8.1. Is $x^{\frac{1}{2}}$ uniformly continuous on $[0, 1]$?

Lipschitz Continuity: “gradient is bounded”

$$\frac{|f(x) - f(y)|}{|x - y|} < L < \infty$$

We can pick $\delta = \frac{\varepsilon}{L}$ everywhere.

Example 8.7 • x^2 on \mathbb{R} is not Lipschitz but it is on a finite interval.

• $x^{\frac{1}{2}}$ is not Lipschitz continuous on $[0, 1]$. However, it's uniformly continuous.

§9 | Lec 9: Oct 15, 2021

§9.1 Picard's Theorem


Let's prove local existence of the theorem.

Theorem 9.1 (Picard)

If $f(t, x)$ and $\partial_x f(t, x)$ are continuous function on a bounded rectangle $R = [t_1, t_2] \times [x_1, x_2]$ and (t_0, x_0) is in interior of R ($t_1 < t_0 < t_2$, $x_1 < x_0 < x_2$). Then \exists a smaller rectangle $R' = [t_0 - a, t_0 + a] \times [x_0 - b, x_0 + b]$ s.t. ODE

$$\dot{x}(t) = f(t, x(t)), \quad x(t_0) = x_0$$

has a solution in R' .



Note: Since R closed and bounded, then f , $\partial_x f$ are bounded, i.e.,

$$\begin{aligned} \max_R f(t, x) &= M \\ \max_R \partial_x f(t, x) &= L \end{aligned}$$

Thus, f is Lipschitz.

Proof Outline:

1. Solving ODE \iff Soling IE
2. Approximate solutions using Picard iteration

$$x_0(t) = x_0, \quad x_n(t) = x_0 + \int_{t_0}^t f(t', x_{n-1}(t')) dt'$$

3. Prove Picard iterates converges

$$\lim_{n \rightarrow \infty} x_n(t) \rightarrow \bar{x}(t)$$

4. Prove limit $\bar{x}(t)$ solves IE.
5. Prove limit $\bar{x}(t)$ is continuous.

6. Prove limit $\bar{x}(t)$ is unique.
7. How big is $R' = [t_0 - a, t_0 + a] \times [x_0 - b, x_0 + b]$?

$$\text{Pick } a \ni aL < 1 \text{ \& } b = Ma \leq |x_0 - x_1| |x_0 - x_2|$$

Proof. 2. Prove Picard iterates converge

a) We have

$$\lim_{n \rightarrow \infty} x_n(t) \iff \lim_{n \rightarrow \infty} x_0(t) + \sum_{k=1}^n x_k(t) - x_{k-1}(t)$$

telescoping sum!

- b) Series $x_0(t) + \sum_{k=1}^n x_k(t) - x_{k-1}(t)$ converges by Weierstrass M-test - If $|f_n(x)| < M_n$
 $\forall n \in \mathbb{N}, x \in D$ and $\sum_{n=0}^{\infty} M_n$ converges, then

$$\sum_{n=0}^{\infty} f_n(x)$$

converges absolutely and uniformly.

- i) Show $x_i(t)$ are all in $R' \subseteq R$ so we can use bounds L, M .

$$\begin{aligned} |x_0(t) - x_0| &= 0 \\ |x_1(t) - x_0| &= \left| \int_{t_0}^t f(t', x_0(t')) dt' \right| \\ &\leq \int_{t_0}^t |f(t', x_0(t'))| dt \\ &\leq \int_{t_0}^t M dt \\ &\leq Ma = b \end{aligned}$$

Thus, $x_1(t)$ is in the rectangle. By induction, every $x_n(t)$ in $R' \subseteq R$.

- ii) Show $\sum_{i=1}^{\infty} |x_i(t) - x_{i-1}(t)|$ is bounded.

Define $\Delta = \max_{R'} |x_1(t) - x_0|$. Then

$$\begin{aligned} |x_2(t) - x_1(t)| &= \left| \int_{t_0}^t f(t', x_1(t')) - f(t', x_0(t')) dt' \right| \\ &\leq \int_{t_0}^t |f(t', x_1(t')) - f(t', x_0(t'))| dt' \\ &\leq \int_{t_0}^t L |x_1(t') - x_0(t')| dt' \\ &\leq \Delta aL \end{aligned}$$

and

$$\begin{aligned} |x_3(t) - x_2(t)| &= \left| \int_{t_0}^t f(t, x_2(t)) - f(t, x_1(t)) dt \right| \\ &\leq \int_{t_0}^t |f(t, x_2(t)) - f(t, x_1(t))| dt \\ &\leq \int_{t_0}^t L |x_2(t') - x_1(t')| dt' \\ &\leq L(\Delta aL)(t - t_0) \\ &\leq \Delta(aL)^2 \end{aligned}$$

Every $|x_n(t) - x_{n-1}(t)|$ depends on $|x_{n-1}(t) - x_{n-2}(t)|$ recursively. The general pattern is

$$\begin{aligned} |x_n(t) - x_{n-1}(t)| &\leq \Delta(aL)^{n-1} \\ \sum_{n=1}^{\infty} |x_n - x_{n-1}| &\leq \sum_{n=0}^{\infty} \Delta(aL)^n \\ &= \frac{\Delta}{1 - aL} \\ &< \infty \end{aligned}$$

Thus, $\sum x_n - x_{n-1}$ converges absolutely and uniformly by the Weierstrass M-test. Therefore,

$$\lim_{n \rightarrow \infty} x_n(t) = \bar{x}(t) \text{ exists!}$$

3. \bar{x} solves I.E.

Idea: We know $|\bar{x} - x_n|$ gets small so break $\left| \bar{x} - x_0 - \int_{t_0}^t f(t', \bar{x}(t')) dt' \right|$ into pieces like $|\bar{x} - x_n(t)|$.

$$\text{subtract } x_n(t) - x_0 - \int_{t_0}^t f(t', x_{n-1}(t')) dt' = 0$$

$$\text{Let } \kappa = \left| \bar{x} - x_0 - \int_{t_0}^t f(t', \bar{x}(t')) dt' \right|.$$

$$\begin{aligned} \kappa &= \left| -(x_n - x_0 - \int_{t_0}^t f(t', x_{n-1}(t')) dt') \right| \\ &\leq |\bar{x} - x_n| + \left| \int_{t_0}^t f(t, \bar{x}) - f(t, x_{n-1}) dt \right| \\ &\leq |\bar{x} - x_n| + \int_{t_0}^t |f(t, \bar{x}) - f(t, x_{n-1})| dt \\ &\leq |\bar{x} - x_n| + aL |\bar{x} - x_{n-1}| \end{aligned}$$

which approaches 0 as $n \rightarrow \infty$ because $\lim_{n \rightarrow \infty} x_n = \bar{x}$.

4. $\bar{x} = \lim_{n \rightarrow \infty} x_n$ is continuous, i.e., given $\varepsilon > 0$, $\exists \delta > 0$ s.t.

$$|t - t'| < \delta \implies |\bar{x}(t) - \bar{x}(t')| < \varepsilon$$

Idea: Split into known things

$$\begin{aligned} |\bar{x}(t) - \bar{x}(t')| &= |\bar{x}(t) - x_n(t) + x_n(t) - x_n(t') + x_n(t') - \bar{x}(t)| \\ &\leq |\bar{x}(t) - x_n(t)| + |x_n(t) - x_n(t')| + |x_n(t') - \bar{x}(t)| \end{aligned}$$

We pick n s.t. $|\bar{x}(t) - x_n(t)| < \frac{\varepsilon}{3} \forall t$ which is possible because Weierstrass implies uniform convergence. Then pick δ s.t.

$$|x_n(t) - x_n(t')| < \frac{\varepsilon}{3} \quad \forall |t - t'| < \delta$$

which is possible because x_n is continuous.

5. \bar{x} is unique.

Idea: Prove $|\bar{x} - \tilde{x}| \leq |\bar{x} - \tilde{x}|$.

- If \tilde{u} is other solution, it also exists in R' .

Proof. (by contradiction) If not, then

$$|\tilde{x}(t_*) - x_0| = b = Ma$$

for some $|t_* - t| < a$. But

$$\begin{aligned} |\tilde{x}(t_*) - x_0| &= \left| \int_{t_0}^{t_*} f(t', \tilde{x}(t')) dt' \right| \\ &\leq \int_{t_0}^{t_*} |f(t', \tilde{x}(t'))| dt' \\ &\leq M(t_* - t_0) \\ &< Ma = b \end{aligned}$$

Contradiction! □

- Have

$$\begin{aligned} |\bar{x}(t) - \tilde{x}(t)| &= \left| \int_{t_0}^t f(t', \bar{x}(t')) - f(t', \tilde{x}(t')) dt' \right| \\ &\leq \int_{t_0}^t |f(t', \bar{x}(t')) - f(t', \tilde{x}(t'))| dt' \\ &\leq \int_{t_0}^t L \max |\bar{x}(t') - \tilde{x}(t')| dt' \\ &\leq La \max |\bar{x}(t') - \tilde{x}(t')| \\ \max |\bar{x}(t) - \tilde{x}(t)| &\leq \max |\bar{x}(t) - \tilde{x}(t)| \end{aligned}$$

which is only possible if $\bar{x}(t) - \tilde{x}(t) = 0$, i.e., solution is unique. □


§10 | Lec 10: Oct 18, 2021

§10.1 Fourier Series

Goal: Solve linear PDE: 3 canonical examples


1. Heat/Diffusion equation

$$\partial_t u(t, x) - \partial_x^2 u(t, x) = 0$$



2. Wave equation

$$\partial_t^2 u = \partial_x^2 u$$



3. Laplace equation:

$$\partial_x^2 u + \partial_y^2 u = 0$$

Question 10.1. How do we solve linear PDEs?

Use linearity to split big problems into small ones that you can solve (find the eigenvectors). Then we split 1 PDE $\rightarrow \infty$ ODEs. First, let's define Fourier series.

Definition 10.1 (Fourier Series) — Fourier Series is a function written as a sum of sines and cosines

$$\begin{aligned} f(x) &= \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \sin(nx) + b_n \cos(nx) \\ &= \sum_{n=-\infty}^{\infty} c_n e^{inx} \end{aligned}$$

where $c_n = c_r + ic_{in}$.

They have amazing properties:

1. They can approximate almost anything

- analytic function
- smooth function
- periodic function
- differentiable function
- continuous/discontinuous function

2. They simplify differentiation!

$$\begin{aligned}\frac{d}{dx}e^{ikx} &= ik e^{ikx} \\ \frac{d^2}{dx^2} \sin kx &= -k^2 \sin kx \\ \frac{d^2}{dx^2} \cos kx &= -k^2 \cos kx\end{aligned}$$

Just like Laplace transform, Fourier series transform differentiation into multiplication problem (easier to deal with).

3. Fourier series are orthogonal

$$\int_{-\pi}^{\pi} \sin mx \cos nx \, dx = 0$$

or

$$\int_{-\pi}^{\pi} \sin mx \sin nx \, dx = 0 \quad \text{if } m \neq n$$

or

$$\int_{-\pi}^{\pi} \cos mx \cos nx \, dx = 0 \quad \text{if } m \neq n$$

This gives easy formulas

From these facts follow from linear algebra, because Fourier series are eigenfunctions of differentiation. They are the correct basis to solve linear PDEs.

§11 | Lec 11: Oct 20, 2021

§11.1 Coefficients of Fourier Series

Question 11.1. How do we calculate Fourier Series $a_n, b_n = ?$

Consider the domain: $[-\pi, \pi]$, finite dimensions N , vector

$$\mathbf{u} = \sum u_i \mathbf{e}_i$$

How do we calculate u_i ?

$$\begin{aligned} \mathbf{u} \cdot \mathbf{e}_j &= \left(\sum_{i=1}^N u_i \mathbf{e}_i \right) \cdot \mathbf{e}_j \\ &= \sum_{i=1}^N u_i (\mathbf{e}_i \cdot \mathbf{e}_j) \\ &= \sum_{i=1}^N \delta_{ij} \end{aligned}$$

where

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

We want to do this in ∞ dimensions – inner product

$$\begin{aligned} N : \langle u, v \rangle &= u \cdot v = \sum_{i=1}^N u_i v_i \\ \infty : \langle u, v \rangle &\propto \int_a^b u(x) v(x) dx \end{aligned}$$

Inner Product: $\langle u, v \rangle \rightarrow \mathbb{R}$ takes in two function & spits out a number. It has to satisfy the following properties

1. Bilinear

$$\langle au + bv, w \rangle = a \langle u, w \rangle + b \langle v, w \rangle$$

2. Symmetric $\langle u, v \rangle = \langle v, u \rangle$.

3. Positivity: $\langle u, u \rangle > 0$ unless $u = 0$.

Inner products are important

- They imply a norm $\|u\| = \sqrt{\langle u, u \rangle}$
- Cauchy-Schwarz Inequality

$$\langle u, v \rangle^2 \leq \langle u, u \rangle \langle v, v \rangle$$

- Triangle inequality

$$\|u + v\| \leq \|u\| + \|v\|$$

Exercise 11.1. Prove these properties.

Now, we will use inner products to calculate Fourier. Define

$$\langle u, v \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} u(x)v(x) dx$$

Under this inner product, $\sin kl$, $\cos kl$ are orthogonal functions, i.e.,

$$\begin{aligned}\langle \sin kx, \cos lx \rangle &= 0 \quad \forall k, l \\ \langle \sin kx, \sin lx \rangle &= 0 \quad \text{if } k \neq l \\ \langle \cos kx, \cos lx \rangle &= 0 \quad \text{if } k \neq l\end{aligned}$$

Note: $1 = \cos 0x$

Proof. Left as exercise, but use

$$\begin{aligned}\cos((k+l)x) &= \cos kx \cos lx - \sin kx \sin lx \\ \sin((k+l)x) &= \sin kx \cos lx + \sin lx \cos kx\end{aligned}$$

Also,

$$\begin{aligned}\langle \sin kx, \sin kx \rangle &= 1 \\ \langle \cos kx, \cos kx \rangle &= 1 \quad k \neq 0 \\ \langle 1, 1 \rangle &= 2\end{aligned}$$

□

We have

$$\begin{aligned}f(x) &= \frac{a_0}{2} + \sum a_k \cos kx + b_k \sin kx \\ \langle f, \cos lx \rangle &= \left\langle \frac{a_0}{2} + \sum a_k \cos kx + b_k \sin kx, \cos lx \right\rangle \\ &= \frac{a_0}{2} \langle 1, \cos lx \rangle + \sum_{k=1}^{\infty} a_k \langle \cos kx, \cos lx \rangle + \sum_{k=1}^{\infty} b_k \langle \sin kx, \cos lx \rangle \\ \langle f, \cos lx \rangle &= a_l \\ \langle f, \sin lx \rangle &= b_l\end{aligned}$$

So we can write any function $f(x)$

$$f(x) = \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx$$

where


$$\begin{aligned}a_k &= \langle f, \cos kx \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos kx dx \\ b_k &= \langle f, \sin kx \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin kx dx\end{aligned}$$

Question 11.2. Are these orthogonal functions under $\langle u, v \rangle$?

Question 11.3. Are there any other kind of L^2 inner product?

Example 11.1

Consider $f(x) = x$



We have

$$\begin{aligned}
 x &= \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx \\
 a_k &= \langle x, \cos kx \rangle \\
 &= \frac{1}{\pi} \int_{-\pi}^{\pi} x \cos kx \, dx \\
 &= 0 - 0 - 0 = 0 \quad (\text{integration by parts}) \\
 b_k &= \langle x, \sin kx \rangle \\
 &= \frac{1}{\pi} \int_{-\pi}^{\pi} x \sin kx \, dx \\
 &= \frac{1}{\pi} \left[-\pi \frac{\cos k\pi}{k} - (-(-\pi)) \frac{\cos(-k\pi)}{k} \right] \quad (\text{integration by parts}) \\
 &= \frac{2(-1)^{k+1}}{k}
 \end{aligned}$$

Thus,

$$x \sim \sum_{k=1}^{\infty} \frac{2(-1)^{k+1}}{k} \sin kx$$

To show that infinite series converges

$$\sum_{k=1}^{\infty} \left| \frac{2(-1)^{k+1}}{k} \right| < 2 \sum_{k=1}^{\infty} \frac{1}{k}$$

which is conclusive (by Weierstrass-M test).

§12 | Lec 12: Oct 22, 2021

§12.1 Convergence of Fourier Series

Consider the last example from last lecture

$$f(x) = x \sim \sum_{k=1}^{\infty} \frac{2(-1)^{k+1}}{k} \sin kx$$


Question 12.1. In what sense does it converge? (What's happening at $\pm\pi$)

Fourier series must be 2π periodic (because $\cos kx$, $\sin kx$ are 2π -periodic) so the y must converge to a 2π -periodic extension of the function.

$$\tilde{f}(x + 2\pi) = \tilde{f}(x)$$

Note: x is C' (derivative continuous) but \tilde{x} is not C' . It is piecewise C' (C' : f continuous and $\frac{df}{dx}$ is continuous).

Piecewise C' on $[a, b]$



f is C' except at finitely many points. At any bad point we have

$$\begin{cases} f(x^-) = \lim_{h \rightarrow 0} f(x-h) & \text{if } f(x^+) \neq f(x^-) \text{ jump} \\ f(x^+) = \lim_{h \rightarrow 0} f(x+h) \\ f'(x^-) = \lim_{h \rightarrow 0} f'(x-h) & \text{if } f(x^+) = f(x^-) \\ f'(x^+) = \lim_{h \rightarrow 0} f'(x+h) & \text{but } f'(x^+) \neq f'(x^-) \text{ corner} \end{cases}$$

Theorem 12.1 (Fourier Convergence)

If $\tilde{f}(x)$ is 2π -periodic, piecewise C' function, then its Fourier series converges to \tilde{f} everywhere except jump points x where the series converges to $\frac{f(x^+) + f(x^-)}{2}$

Question 12.2. Recall the example at the beginning, why is there no cosines for x ?

Odd/even symmetries!

Fact 12.1. We have

$$\begin{aligned} \text{odd} + \text{odd} &= \text{odd} \\ \text{even} + \text{even} &= \text{even} \end{aligned}$$

and

$$\begin{aligned}\text{odd} \times \text{odd} &= \text{even} \\ \text{even} \times \text{even} &= \text{even} \\ \text{odd} \times \text{even} &= \text{odd}\end{aligned}$$

and

$$\begin{aligned}\int_{-a}^a \text{odd } dx &= 0 \\ \int_{-a}^a \text{even } dx &= 2 \int_0^a \text{even } dx\end{aligned}$$

This implies odd functions f have sine series and even functions have cosine series.

§13 | Lec 13: Oct 27, 2021

Recap:

$$f(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx$$

where the coefficients are calculated as follows

$$\begin{aligned} a_k &= \langle f, \cos kx \rangle \\ b_k &= \langle f, \sin kx \rangle \\ \langle u, v \rangle &= \frac{1}{\pi} \int_{-\pi}^{\pi} u(x)v(x) dx \end{aligned}$$

Symmetry simplifies a_k, b_k . Fourier series converges for periodic and piecewise C^1 functions.

§13.1 Complex Fourier Series

Recall the Euler's formula

$$e^{ikx} = \cos kx + i \sin kx$$

Also,

$$\begin{aligned} \cos kx &= \frac{e^{ikx} + e^{-ikx}}{2} \\ \sin kx &= \frac{e^{ikx} - e^{-ikx}}{2i} \end{aligned}$$

So,

$$f \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx \quad \leftrightarrow \quad \sum_{k=-\infty}^{\infty} c_k e^{ikx}$$

We want $c_k = \langle f, e^{ikx} \rangle$

$$\langle e^{ikx}, e^{ikx} \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} e^{2ikx} dx$$

which is not necessarily positive and we want it to be strictly positive, i.e., norm.

$$\begin{aligned} \int_{-\pi}^{\pi} e^{2ikx} dx &= \left[\frac{e^{2ikx}}{2ik} \right]_{-\pi}^{\pi} \\ &= \frac{e^{2\pi ki} - e^{-2\pi ki}}{2ik} \\ &= \frac{\sin 2\pi k}{k} \\ &= 0 \end{aligned}$$

To fix this, let's define Hermitian inner product

$$\langle f, g \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \overline{g(x)} dx$$

where $x \in (-\pi, \pi]$ and $f, g : (-\pi, \pi] \rightarrow \mathbb{C}$. So

$$\begin{aligned} c_k &= \langle f, e^{ikx} \rangle \\ c_k &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx \end{aligned}$$

Question 13.1. How do Fourier series work with integration?

Integration makes things smoother. We have

$$f(x) \sim \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx$$

$$\int f(x) dx \sim \int \frac{a_0}{2} dx + \sum_{k=1}^{\infty} a_k \int \cos kx dx + b_k \int \sin kx dx$$

Question 13.2. Is this okay?

Notice that

$$\int \cos kx dx = \frac{\sin kx}{k} \quad \int \sin kx dx = \frac{-\cos kx}{k}$$

Problem: If $f(x) = 1$, then

$$f \sim 1$$

$$\int_0^x f dx \sim 2 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \sin kx$$

Constants terms in Fourier series are bad under integration.

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx$$

Integration is fine if the function has mean 0

$$\int_{-\pi}^{\pi} f(x) dx = 0$$

Compare $f(x) = 1$ and $g(x) = x$.

Remark 13.1. Fourier series need piecewise C^1 . To have Fourier of f' , it must be C^1 so f must be continuous (can have corners but not jumps).

$$f = a_0 + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx$$

$$f' = -ka_k \sin kx + kb_k \cos kx$$

if f is continuous.

Summary:

- Integrate: divide by k
- Differentiation: multiply by k

§14 | Lec 14: Oct 29, 2021

§14.1 Rescaling Intervals of Fourier Series

We know Fourier series on $[-\pi, \pi]$. What about $[-l, l]$? We use coordinate transformation

$$\begin{aligned} y &= \frac{\pi}{l} x \\ F(y) &= f(x(y)) \\ F(y(x)) &= f(x) \end{aligned}$$

We have

$$F(y) = f(x(y)) = f\left(\frac{l}{\pi}y\right)$$

So $F(y) = \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos ky + b_k \sin ky$

$$\begin{aligned} a_k &= \frac{1}{\pi} \int_{-\pi}^{\pi} F(y) \cos ky \, dy \\ &= \frac{1}{\pi} \int_{-l}^l F(y(x)) \cos ky(x) \frac{\pi}{l} \, dx \\ &= \frac{1}{l} \int_{-l}^l f(x) \cos \left(\frac{k\pi}{l}x\right) \, dx \end{aligned}$$

So

$$f(x) = F(y(x)) = \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos \frac{k\pi}{l}x + b_k \sin \frac{k\pi}{l}x$$

We can find b_k similarly.

§15 | Lec 15: Nov 1, 2021

§15.1 The Relationship between Smoothness and Fourier Coefficients

Smother functions (more differentiable) have faster decaying Fourier coefficients. (infinitely differentiable leads to exponential decay).

Example 15.1 • Discontinuous function $\rightarrow c_k \propto \frac{1}{k}$

- $C^0 \rightarrow c_k \propto \frac{1}{k^2}$
 - $C^1 \rightarrow c_k \propto \frac{1}{k^3}$
 - $C^2 \rightarrow c_k \propto \frac{1}{k^4}$
- Why?

Recall these definitions

Definition 15.2 — $\forall \varepsilon, x \exists N$ s.t.

$$|f_n(x) - f(x)| < \varepsilon \quad \forall n > N(x)$$

Then, $f_n(x) \rightarrow f(x)$ (pointwise convergence).

Definition 15.3 — $\forall \varepsilon, \exists N$ s.t.

$$|f_n(x) - f(x)| < \varepsilon \quad \forall n > N, \quad \forall x$$

Then, $f_n(x) \rightarrow f(x)$ (uniform convergence).

Series converges $\sum_{k=1}^{\infty} f_k(x) \rightarrow g(x)$ if

$$s_n(x) = \sum_{k=1}^n f_k(x) \rightarrow g(x) \text{ as } n \rightarrow \infty$$

Weierstrass M-test: If $|f_n(x)| < M_n$ and $\sum_{n=1}^{\infty} M_n < \infty$, then $\sum_{n=1}^{\infty} f_n(x)$ converges (absolutely/uniformly). So the limit is continuous if f_n are continuous.

Consider a complex Fourier series

$$f \sim \sum_{k=-\infty}^{\infty} c_k e^{ikx}$$

Theorem 15.4

If $\sum_{k=-\infty}^{\infty} |c_k| < \infty$, then the Fourier series is “good”, i.e., the limit of the Fourier series is continuous.

Proof. Weierstrass!

$$|c_k e^{ikx}| \leq |c_k| |e^{ikx}| = |c_k|$$

□

Corollary 15.5

If $|c_k| < \frac{M}{|k|^\alpha}$ where $\alpha > 1$. Then Fourier series is continuous.

Proof. $\sum_{k=1}^{\infty} \frac{M}{k^\alpha} < \infty$ for $\alpha > 1$ by comparison test. □

Note:

$$f \sim \sum_{k=-\infty}^{\infty} c_k e^{ikx}$$

$$f' \sim \sum_{k=-\infty}^{\infty} ikc_k e^{ikx}$$

Differentiation: $c_k \rightarrow ikc_k$ or $|c_k| \rightarrow k|c_k|$

Theorem 15.6

If $\sum_{k=1}^{\infty} |k|^n |c_k| < \infty$ where $f \sim \sum_{k=-\infty}^{\infty} c_k e^{ikx}$, then $f^{(n)}$ is continuous (f is C^n)

Proof. Have

$$f^{(n)} \sim \sum_{k=-\infty}^{\infty} (ik)^n c_k e^{ikx}$$

then Weierstrass $|(ik)^n c_k e^{ikx}| \leq |k|^n |c_k|$ □

Corollary 15.7

If $|c_k| < \frac{M}{|k|^\alpha}$ where $\alpha > n + 1$ then f is n times differentiable.

Proof. Comparison test: $c_k = \frac{1}{k^2}$, then

$$|c_k| < \frac{1}{k^{1.5}} \propto \frac{1}{k}$$

So,

$$\begin{aligned} \frac{1}{k^2} &\rightarrow C^0 \\ \frac{1}{k^3} &\rightarrow C^1 \\ \frac{1}{k^4} &\rightarrow C^2 \\ &\vdots \end{aligned}$$

□

§16 | Lec 16: Nov 3, 2021

§16.1 Hilbert Spaces & Convergence in Norm

Goal: Prove Fourier series converge “in norm”. First, we need some definitions.

Definition 16.1 (L^2 integrable) — f is L^2 integrable if $\|f\|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |f(x)|^2 dx < \infty$.

Definition 16.2 (Hilbert Space) — Hilbert space is vector space of L^2 integrable function

Proof. Have

- Has a 0 (0 function)
- Closed under addition

$$\|f + g\| \leq \|f\| + \|g\| < \infty$$

- Closed under scalar multiplication

$$\|cf\| = |c|\|f\| < \infty$$

□

- test other axioms ...

Note: L^2 function have Fourier series.

Proof. $|c_k| = |\langle f, e^{ikx} \rangle| \leq \|f\| \|e^{ikx}\| < \infty$ (Cauchy-Schwarz).

□

Note: L^2 functions are “abnormal” TBA

Fact 16.1. Hilbert spaces are complete (every “convergent” sequence has a limit that is L^2)

Definition 16.3 — “Convergent” means Cauchy sequence for sequence $a_n \rightarrow a$. We need

$$\text{Cauchy} : \forall \varepsilon, \exists N \ni |a_m - a_n| < \varepsilon \quad \forall m, n > N$$

Aside: Completeness is the difference between rationals \mathbb{Q} , and reals \mathbb{R} (\mathbb{Q} isn’t complete because π is limit of sequence in \mathbb{Q} but $\pi \notin \mathbb{Q}$). Completeness matters for taking limits.

Definition 16.4 (Convergence in Norm) — $f_n(x) \rightarrow f(x)$ if $\|f_n(x) - f(x)\| \rightarrow 0$ as $n \rightarrow \infty$.

We’ll prove Fourier series converge to their function in norm in a general way for a general ∞ -dim vector space V with an inner product.

Definition 16.5 (Orthonormal System) — Orthonormal system $\phi_1, \phi_2, \dots \in V$

$$\langle \phi_i, \phi_j \rangle = \delta_{ij} = \begin{cases} 0, & i \neq j \\ 1, & i = j \end{cases}$$

Definition 16.6 (General Fourier Series) — $f \sim \sum_{k=1}^{\infty} c_k \phi_k$ where $c_l = \langle f, \phi_l \rangle$ and c_k comes from $\langle f, \phi_l \rangle$ on both sides.

Theorem 16.7

The truncated Fourier series

$$s_n = \sum_{k=1}^{\infty} c_k \phi_k$$

is the best approximation to f in least squares sense, that is, consider $V_n = \text{span}\{\phi_1, \dots, \phi_n\}$ and take any $p_n = \sum_{k=1}^n d_k \phi_k \in V_n$ then

$$\|s_n - f\| \leq \|p_n - f\| \quad \forall p_n \in V_n$$

Proof. We have

$$\begin{aligned} p_n &= \sum_{k=1}^n d_k \phi_k \\ s_n &= \sum_{k=1}^n c_k \phi_k \\ c_k &= \langle f, \phi_k \rangle \end{aligned}$$

Then,

$$\begin{aligned} \|p_n\|^2 &= \langle p_n, p_n \rangle \\ &= \left\langle \sum_{k=1}^n d_k \phi_k, \sum_{l=1}^n d_l \phi_l \right\rangle \\ &= \sum_{k=1}^n \sum_{l=1}^n d_k d_l \langle \phi_k, \phi_l \rangle \\ &= \sum_{k=1}^n \sum_{l=1}^n d_k d_l \delta_{kl} \\ &= \sum_{k=1}^n |d_k|^2 \end{aligned}$$

and

$$\begin{aligned} \|p_n - f\|^2 &= \langle p_n - f, p_n - f \rangle \\ &= \langle p_n, p_n \rangle - 2\langle p_n, f \rangle + \langle f, f \rangle \\ &= \sum_{k=1}^n |d_k|^2 - 2 \left(\sum_{k=1}^n d_k \langle \phi_k, f \rangle \right) + \|f\|^2 \\ &= \sum_{k=1}^n |d_k - c_k|^2 - \sum_{k=1}^n |c_k|^2 + \|f\|^2 \end{aligned}$$

Pick $d_k = c_k$ – norm minimized by s_n . □

§17 | Lec 17: Nov 5, 2021

§17.1 Pointwise Convergence of Fourier Series

We know Fourier series converge in norms for continuous, piecewise C^1 , periodic functions. But Fourier series seemed to work even for discontinuous functions too (Gibbs phenomenon!). Today we will prove it works pointwise for discontinuous function if $s_n = \sum_{k=-n}^n c_k e^{ikx}$. Prove

$$\lim_{n \rightarrow \infty} s_n(x) = \frac{1}{2} (f(x^+) + f(x^-))$$

1. Use the formulas for c_k

$$\begin{aligned} s_n &= \sum_{k=-n}^n \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) e^{-iky} dy \right) e^{ikx} \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) \underbrace{\left(\sum_{k=-n}^n e^{ik(x-y)} \right)}_{dy?} dy \end{aligned}$$

Notice that $\sum_{k=-n}^n e^{ikx}$ is a geometric series

$$\begin{aligned} \sum_{k=-n}^n e^{ikx} &= e^{-inx} \left(\frac{e^{i(2n+1)x} - 1}{e^{ix} - 1} \right) \\ &= \vdots \\ &= \frac{\sin\left((n + \frac{1}{2})x\right)}{\sin \frac{1}{2}x} \end{aligned}$$

So

$$\begin{aligned} s_n(x) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) \frac{\sin\left((n + \frac{1}{2})(x - y)\right)}{\sin \frac{1}{2}(x - y)} dy \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x + y) \frac{\sin\left((n + \frac{1}{2})y\right)}{\sin \frac{1}{2}y} dy \\ &= \frac{1}{2\pi} \int_0^{\pi} f(x + y) \frac{\sin\left((n + \frac{1}{2})y\right)}{\sin \frac{1}{2}y} dy + \frac{1}{2\pi} \int_{-\pi}^0 f(x + y) \frac{\sin\left((n + \frac{1}{2})y\right)}{\sin \frac{1}{2}y} dy \end{aligned}$$

WTS:

$$\begin{aligned} \lim_{n \rightarrow \infty} \text{top} &= f(x^+) \\ \lim_{n \rightarrow -\infty} \text{bottom} &= f(x^-) \end{aligned}$$

Note

$$\begin{aligned} \frac{1}{\pi} \int_0^{\pi} \frac{\sin\left((n + \frac{1}{2})y\right)}{\sin y} dy &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{k=-n}^n e^{iky} dy \\ &= 1 \end{aligned}$$

which is valid if only e^{i0} counts.

2. Prove a difference integral $\rightarrow 0$ by showing that it's a Fourier coefficient. Prove

$$\frac{1}{2\pi} \int_0^{\pi} (f(x + y) - f(x^+)) \frac{\sin\left((n + \frac{1}{2})y\right)}{\sin \frac{1}{2}y} dy = 0$$

Notice that

$$g(y) \equiv \frac{f(x+y) + f(x^-)}{\sin \frac{1}{2}y}$$

is piecewise continuous $\forall y \in [0, \pi]$. We need $\int_0^\pi g(y) \sin \left(n + \frac{1}{2}\right) y dy = 0$. Note that

$$\sin \left(n + \frac{1}{2}\right) y = \sin \frac{1}{2}y \cos ny + \cos \frac{1}{2}y \sin ny$$

Then,

$$\int_0^\pi \left(g(y) \sin \frac{y}{2}\right) (0)ny + \left(g(y) \cos \frac{y}{2} \sin ny\right)$$

But we know Fourier coefficients decay for all L^2 integrable functions. So these terms $\rightarrow 0$ and we prove pointwise convergence.

§18 | Lec 18: Nov 8, 2021

§18.1 Heat Equation

We've been learning about Fourier series

$$\begin{aligned} f(x) &= \frac{a_0}{2} + \sum_{k=1}^{\infty} a_k \cos kx + b_k \sin kx \\ &= \sum_{k=-\infty}^{\infty} c_k e^{ikx} \end{aligned}$$

Question 18.1. Why is Fourier series useful?

It's used to solve PDEs; specifically, we want to investigate the heat equation in this lecture. For us,

$$\partial_t u(t, x) - \kappa \partial_x^2 u(t, x) = 0$$

where κ is constant (diffusing constant). Note that

- 1x time derivative \implies 1x initial condition
- 2x space derivative \implies 2x boundary condition

Types of boundary condition

$$\begin{aligned} u(t, 0) &= \alpha(t) && \text{(Dirichlet boundaries)} \\ \partial_x u(t, 0) &= \mu(t) && \text{(Neumann boundaries)} \\ \partial_x u + \beta(t)u &= \tau(t) && \text{(Robin/Mixed boundaries)} \end{aligned}$$

Homogeneous \implies RHS = 0, i.e., $u = 0$, $\partial_x u = 0$, or $\partial_x u + \beta u = 0$.

Question 18.2. How do we solve this? (infinitely harder than an ODE!)

Assume $u(t, x) = T(t)X(x)$. Substitute into $\partial_t u - \kappa \partial_x^2 u = 0$

$$\begin{aligned} T'(t)X(x) - \kappa T(t)X''(x) &= 0 \\ \frac{T'(t)}{\kappa T(t)} &= \frac{X''(x)}{X(x)} \\ \frac{T'(t)}{\kappa T(t)} &= \frac{X''(x)}{X(x)} \end{aligned}$$

This can only be true if neither depends on t or x , i.e., constant. So

$$\frac{T'}{\kappa T} = \frac{X''}{X} = \lambda$$

What sign is λ ?

- If $\lambda > 0$, we get exponential growth which isn't physical.

$$\begin{aligned} X'' &= \lambda X \\ X &= Ae^{\sqrt{\lambda}x} + Be^{-\sqrt{\lambda}x} \\ T' &= \kappa \lambda T \\ T &= Te^{\kappa \lambda t} \end{aligned}$$

- $\lambda = 0$,

$$\begin{aligned} X'' &= 0 \\ X &= Ax + B \\ T' &= 0 \\ T &= T_0 \end{aligned}$$

- If $\lambda < 0$, redefine $\lambda \rightarrow -\lambda$


$$\begin{aligned} X'' &= -\lambda X \\ X &= A \sin \sqrt{\lambda} x + B \cos \sqrt{\lambda} x \\ T' &= -\kappa \lambda T \\ T &= T_0 e^{-\kappa \lambda t} \end{aligned}$$

So either $\lambda = 0$, or $\lambda < 0$.

Example 18.1

Consider

$$\begin{aligned} \partial_t u - \partial_x^2 u &= 0 \\ u(0, x) &= u_0(x) \\ u(t, 0) &= 0 \\ u(t, l) &= 0 \end{aligned}$$



If we assume $u(t, x) = T(t)X(x)$ we find

$$\frac{T'}{\kappa T} = \frac{X'}{X} = \lambda$$

If $\lambda = 0$, $X = Ax + B$, but $u(t, 0) = u(t, l) = 0 \implies A = B = 0$. So $\lambda < 0$. Let's write $\lambda = -\omega^2$,

$$\begin{aligned} X'' &= -\omega^2 X \\ X &= A \cos \omega x + B \sin \omega x \end{aligned}$$

Use BC's to get A, B .

$$\begin{aligned} u(t, 0) = T(t)X(0) = 0 &\implies X(0) = A = 0 \\ X &= \sin \omega x \end{aligned}$$

Example 18.2 (Cont'd)

And we have

$$\begin{aligned} u(t, l) = T(t)X(l) &= B \sin \omega l = 0 \\ \implies \omega l &= k\pi, \quad k = 1, 2, \dots \end{aligned}$$

Thus, $\omega = \frac{k\pi}{l}$, and it's an eigenvalue and $\sin \frac{n\pi}{l}x$ is an eigenfunction.

The final solution is

$$u(t, x) = \sum_{n=1}^{\infty} \left(\hat{u}_{0,k} e^{-\frac{n^2 \pi^2}{l^2} \kappa t} \sin \left(\frac{n\pi}{l} x \right) \right)$$

We get $\hat{u}_{0,k}$ from the Fourier series of the function $u_0(x)$.

Note: The Fourier coefficients decay more quickly as k gets larger so diffusion smooth things out. If we have source term

$$\partial_t u - \partial_x^2 u = f(t, x)$$

We can express f as a Fourier series and solve an ODE for each Fourier coefficient.

Summary:

1. We assume separable solution: $u(t, x) = T(t)X(x)$
2. Substituting gives an eigenvalue problem

$$X'' = \lambda X$$

3. The boundary conditions imply $\lambda = -\omega^2$ where $\omega = \frac{n\pi}{l}$, $x \in [0, l]$.
4. Linearity mean we sum up all the eigenfunctions

$$u(t, x) = \sum_{n=1}^{\infty} \left(\hat{u}_{0,k} e^{-\kappa \frac{(k-\pi)^2}{2^2} t} \right) \sin k \frac{\pi}{l} x$$

5. We use the initial condition to determine the Fourier series coeff, $\hat{u}_{0,k}$

§19 | Lec 19: Nov 10, 2021

§19.1 Wave Equation

Goal: Solve the wave equation

1. Look for separable solutions

$$u(t, x) = T(t)X(x)$$

to

$$\partial_t^2 u(t, x) = c^2 \partial_x^2 u(t, x)$$

c : wave speed ($\frac{\text{space}}{\text{time}}$).

$$\partial_t^2(TX) = c^2 \partial_x^2(TX)$$

$$T''X = c^2 TX''$$

$$\frac{T''(t)}{T(t)} = \frac{c^2 X''(x)}{X(x)} = \lambda$$

a) $\lambda = \omega^2 > 0$, $T'' = \omega^2 T \implies T = e^{\omega t}$ or $T = e^{-\omega t}$ and $X'' = \frac{\omega^2}{c^2} X$, $X = e^{\frac{\omega x}{c}}$ or $X = e^{-\frac{\omega x}{c}}$

b) $\lambda = 0$:

$$T'' = 0 \implies T = A + Bt$$

$$X'' = 0 \implies X = C + Dx$$

$$TX = a + bt + cx + d + x$$

c) $\lambda = -\omega^2 < 0$

$$T'' = -\omega^2 T \implies T = \sin \omega t \text{ or } T = \cos \omega t$$

$$X'' = \frac{-\omega^2}{c^2} X \implies X = \frac{\sin \omega x}{c} \text{ or } X = \frac{\cos \omega x}{c}$$

Next, let's decide on the sign of λ using the boundary conditions, e.g., homogeneous, Dirichlet, boundary conditions

$$u(t, 0) = u(t, l) = 0$$

If $\lambda > 0$

$$u = Ae^{\omega t} e^{\frac{\omega x}{c}} + Be^{-\omega t} e^{\frac{\omega x}{c}} + Ce^{\omega t} e^{-\frac{\omega x}{c}} + De^{-\omega t} e^{-\frac{\omega x}{c}}$$

but the boundary condition implies that $A = B = C = D = 0$.

If $\lambda = 0$, similarly $u = 0$ is only possibility.

If $\lambda < 0$, $u = T(t)X(x)$, $T = \sin \omega t, \cos \omega t$, $X = \sin \frac{\omega x}{c}, \cos \frac{\omega x}{c}$

$$u(t, 0) = T(t)X(0) = X(0) = 0$$

$$\implies A \sin \frac{\omega 0}{c} + B \cos \frac{\omega 0}{c} = B = 0$$

So $X = A \sin \frac{\omega x}{c}$.

$$u(t, l) = T(t)X(l) = A \sin \frac{\omega l}{c}$$

$$\implies \frac{\omega l}{c} = n\pi, \quad n = 1, \dots$$

$$\omega = \frac{n\pi c}{l}$$

In general, the solution is

$$u(t, x) = \sum_{n=1}^{\infty} \left(A_n \cos \frac{n\pi c}{l} t + B_n \sin \frac{n\pi c}{l} t \right) \sin \frac{n\pi x}{l}$$

TO find A_n, B_n , we use the initial conditions

$$u(0, x) = f(x), \quad \partial_t u(0, x) = g(x)$$

Example 19.1

Consider $\partial_t^2 u = \partial_x^2 u$, $x \in [0, 1]$

$$\begin{aligned} u(t, 0) &= u(t, 1) = 0 \\ u(0, x) &= \begin{cases} x, & 0 \leq x \leq \frac{1}{2} \\ 1 - x, & \frac{1}{2} \leq x \leq 1 \end{cases} \\ \partial_t u(0, x) &= 0 \\ c = 1, \quad l = 1 &\implies \omega_n = n\pi \end{aligned}$$

Using Dirichlet, the general solution is

$$u(t, x) = \sum_{n=1}^{\infty} (a_n \cos n\pi t + b_n \sin n\pi t) \sin n\pi x$$

We want to get a_n, b_n with ICs. For $t = 0$,

$$\begin{aligned} u(0, x) &= \sum_{n=1}^{\infty} a_n \sin n\pi x \\ &= \begin{cases} x, & 0 \leq x \leq \frac{1}{2} \\ 1 - x, & \frac{1}{2} \leq x \leq 1 \end{cases} \\ a_n &= \frac{1}{2} \int_0^1 f(x) \sin n\pi x dx \end{aligned}$$

In general,

$$\begin{aligned} \sum_{n=1}^{\infty} a_n \int_0^1 \sin n\pi x \sin m\pi x dx &= \int_0^{\frac{1}{2}} x \sin n\pi x dx + \int_{\frac{1}{2}}^1 (1 - x) \sin n\pi x dx \\ &= \begin{cases} 0, & n \text{ even} \\ 2 \left(-\left[\frac{1}{2} \frac{\cos n\pi/2}{n\pi} \right] + \frac{1}{(n\pi)^2} [\sin n\pi]_0^{\frac{1}{2}} \right) & n \text{ odd} \end{cases} = \dots = \frac{4}{\pi^2} \frac{\sum (-1)^k \cos(2k+1)\pi/2 \sin(2k+1)\pi/2}{2k+1} \end{aligned}$$

§20 | Lec 20: Nov 12, 2021

§20.1 Midterm 2


Need to know:

- How to calculate Fourier series
- How to test convergence?
- How does convergence relate to continuity?
- How does symmetry affect Fourier series?
- How does rescaling affect Fourier series?
- How does differentiation/integration affect Fourier series?
- How does smoothness (C^n) relate to Fourier series?
- How to analyze generalized Fourier series in Hilbert spaces?
- How to use Fourier series to solve PDEs?

§20.2 Laplace Equations

Consider

$$\partial_x^2 u + \partial_y^2 u = 0$$



The Laplace equation satisfies the Maximum principle, i.e., the maximum (and minimum) value of a solution to the Laplace equation must occur on the boundary,

Proof. If a function u has a maximum at (x, y) then

1. $\partial_x^2 u < 0$
2. $\partial_y^2 u < 0$

But Laplace says $\partial_x^2 u + \partial_y^2 u = 0$. Thus, we can't have local maximum (or minimum) in the domain. \square

Theorem 20.1

Solutions to Laplace equation, with given boundary conditions, are unique.

Proof. Suppose there exist u_1 and u_2 where

$$\begin{aligned}\partial_x^2 u_1 + \partial_y^2 u_1 &= 0, & u_1 &= f \text{ on the boundary} \\ \partial_x^2 u_2 + \partial_y^2 u_2 &= 0, & u_2 &= f \text{ on the boundary}\end{aligned}$$

Consider $u_1 - u_2 = \Delta u$. Since Laplace equation is linear, Δu solves Laplace. And we know that $\Delta u = 0$ on the boundary. Therefore, $\Delta u = 0$ everywhere (by maximum principle). \square

Example 20.2

Consider

$$\begin{array}{ccccc} & 1 & u(x, 1) = 0 & 1 & \\ & \downarrow & & \downarrow & \\ u(0, y) = 0 & \square & \partial_x^2 u + \partial_y^2 u = 0 & \square & u(1, y) = 0 \\ & \uparrow & & \uparrow & \\ & 0 & & 1 & \end{array}$$

and

$$u(0, x) = \begin{cases} x \\ 1 - x \end{cases}$$

Have $u = X(x)Y(y)$

$$X''Y + XY'' = 0$$

$$\frac{X''}{X} = \frac{-Y''}{Y} = \lambda$$

- $\lambda = -\omega^2 < 0$

$$\begin{aligned}X'' &= -\omega^2 X \implies X = \cos \omega x, \sin \omega x \\ Y'' &= \omega^2 Y \implies Y = e^{\omega y}, e^{-\omega y}, \cosh y, \sinh y\end{aligned}$$

- $\lambda = 0,$

$$\begin{aligned}X &= 1, x \\ Y &= 1, y\end{aligned}$$

- $\lambda = \omega^2 > 0$

$$\begin{aligned}X &= e^{\omega x}, e^{-\omega x} \\ y &= \cos \omega y, \sin \omega y\end{aligned}$$

Example 20.3 (Cont'd)

Only $\lambda = -\omega^2 < 0$ works. For $X = \sin n\pi x$,

$$X(0) = X(1) = 0 \implies \omega = n\pi$$

Therefore,

$$\begin{aligned} Y &= Ae^{n\pi}Be^{-n\pi y} \\ Y(1) = 0 &\implies Ae^{n\pi} + Be^{-n\pi} = 0 \\ \iff Y(y) &\propto \sinh n\pi(y - 1) \end{aligned}$$

In general,

$$u = \sum_{n=1}^{\infty} a_n \sinh n\pi(1 - y) \sin n\pi x$$

and

$$\begin{aligned} u(x, 0) &= \sum_{n=1}^{\infty} a_n \sinh n\pi \sin n\pi x = \begin{cases} x, & x < \frac{1}{2} \\ 1 - x, & x > \frac{1}{2} \end{cases} \\ &= \sum_{k=1}^{\infty} (-1)^k \frac{4}{\pi^2(2k+1)^2} \sin n\pi x \end{aligned}$$

where

$$a_{2k+1} = \frac{(-1)^k 4}{\pi^2(2k+1)^2}$$