Recoil transport and environmental exposure of black-hole merger remnants

Venkata Hemanth Athota[1,*]

[1]*Independent researcher*

Black-hole mergers in stellar clusters leave remnants whose recoil carries them through environments of different density and composition. Their availability for subsequent interactions depends on residence in those environments, as well as on gravitational retention. We study when continuous drift and diffusion preserve finite-time residence under discrete gravitational encounters. Published multimass distributions for six Galactic clusters supply consistent potentials, species distributions and local scattering backgrounds. Collisionless controls connect these inputs to orbital occupation, while a conditional Omega Centauri calculation compares finite encounters with a diffusion process having the same local velocity-change moments. A backward occupation equation relates their mean-residence difference to a nonlocal generator residual. We test a first-response approximation through complete-transition replacements, with predictions recorded before independent mixture trajectories are generated. The numerical endpoint distributions differ, but the response test remains limited by sampling precision, refinement uncertainty and numerical-boundary continuation. It therefore does not establish or reject the proposed prediction accuracy. A fixed precision comparison finds that a symmetric covariance root substantially reduces response-estimator variance at similar measured cost, while proportional time stratification adds no resolved benefit. This improves the feasibility of a further test without resolving prediction accuracy. The calculations distinguish exact response identities, measurable encounter effects and the numerical precision required for an observable-specific transport criterion. They remain conditional on the adopted hosts and encounter prescription, without a calibrated cluster residence or merger-rate prediction.

I. INTRODUCTION

A black hole formed by a merger can receive a recoil from anisotropic gravitational-wave emission. Within a stellar cluster, its subsequent motion reflects both that recoil and the state of the system in which the merger occurs. The progenitor's center-of-mass motion, merger position, host potential, and gravitational scattering determine the remnant's evolving position and velocity. A gravitationally bound remnant need not remain near the center, and entry into a dense region does not imply that the remnant comes to rest there.

The distinction matters whenever the surrounding population influences subsequent evolution. Residence measures the time spent in a specified region, whereas density exposure weights the trajectory by the material encountered along it. Neither is fixed by a retention flag alone. Their calculation requires a dynamical description with enough information to distinguish repeated excursions, continued wandering, and changes in the environment during the remnant's motion.

The kinetic description follows from the velocity-change moments of inverse-square interactions [1]. Anisotropic relaxation calculations connect such moments to direct stellar dynamics [2], while kicked-black-hole simulations demonstrate that gravitational encounters can increase or decrease the remnant's orbital energy [3]. Full encounter distributions have also been studied for stellar energy relaxation around massive black holes [4]. That nuclear setting differs from the cluster-remnant problem considered here; the relevant methodological connection is the distinction between an encounter transition law and its diffusion approximation.

Our primary question is when a continuous drift-and-diffusion description preserves the finite-time residence and exposure of a kicked remnant. We hypothesize that the relevant accuracy depends on the velocity increments sampled by the remnant and on their effect on its subsequent occupation of dense regions. The mean population size or a central diffusion coefficient alone does not specify that effect. We express this question through a matched encounter comparison and a backward equation for expected remaining occupation in Sec. XI. This establishes an observable-specific theoretical target for the calculations.

Step 1 defines merger-conditioned transport in position and velocity, with a consistent matter inventory, reference frame and encounter prescription. Step 2 defines entry, residence and species-resolved exposure on the resulting histories. Their common output is a timed description of occupation; it retains continued motion after entry and the status of incomplete histories.

We use published multimass-model posteriors for six Galactic clusters as the dynamical backgrounds [5–7]. Each adopted realization supplies a potential and mass-resolved phase-space distributions. Reference reconstructions and a joint-parameter sensitivity pilot establish different reservoir sizes and radial scattering profiles. A detailed NGC 3201 calculation develops the host-to-collision connection, and the same calculations extend across the six hosts. Galactic reference orbits supply their external-force diagnostics. Finite-angle calculations then provide event rates, matched moments and samples from one common encounter law. Collisionless reference paths then measure finite-time residence and path-integrated encounter budgets in each host. A conditional ω Cen pilot then compares finite-encounter and diffusion residence under a common numerical collision law. Its grid and time-step comparisons retain sampling uncertainty separately from physical calibration. A separate complete-transition response test records predictions before independent mixture validation, including a previously unused oblique launch.

* bharadwaj.hemanth.002@gmail.com

II. OBSERVATIONAL INPUTS AND COMPARISON METHOD

A. Sample and source distinctions

We examine NGC 0104 (47 Tuc), NGC 1851, NGC 3201, NGC 5139 (ω Centauri), NGC 6121 (M4), and NGC 6397. This deliberately selected sample provides varied mass and size scales and public kinematic and model products. It is not a complete cluster census or a selection-corrected population of black-hole-merger hosts.

For five clusters we use the original binned Hubble Space Telescope (HST) proper-motion products of Libralato et al. [8]. For ω Centauri we use the logarithmic radial bins released with Häberle et al. [9]. The former dispersion profiles use bright subgiant and red-giant tracers, whereas the latter use the full magnitude range. Stellar mass and selection can affect dispersion in a partially relaxed system; converting the units cannot remove this distinction.

We inspect the original Gaia EDR3 field catalogs and fitted radial curves of Vasiliev and Baumgardt [10], separately from compiled kinematic profiles. Membership probabilities, quality flags, measurement errors, and source identifiers describe the field catalogs. They do not make every listed star a cluster member, and raw proper-motion spread is not an intrinsic velocity dispersion. The released smooth radial curves are correlated fit products; their grid points are not independent observational bins.

Structural comparison values are taken from one captured path of the Baumgardt cluster database associated with the modeling and distance studies in Refs. [11, 12].[1] These values are model-derived estimates. In particular, a three-dimensional half-mass radius is distinct from a projected half-light or core radius. The compiled surface profiles mix normalized stellar counts and, for some sources, surface brightness; they are not directly a three-dimensional mass density.

For a separate view of host uncertainty, we inspect the released standard-fit multimass posterior samples discussed by Dickson et al. [6, 7]. Original observations, compilations, and these posterior products share underlying evidence. We keep the views separate rather than multiply them together as independent likelihood factors. The complete data collection comprises 51 captured files, including supporting source definitions and historical candidate observation bundles.

[1] The selected structural table is the database's /newdata/ parameter table. The root-path table is retained as a separate comparison; values from the two paths are not merged.

B. Native measurements, units, and shared uncertainty

For angular radius α, distance D, and proper-motion dispersion σ_μ, the small-angle conversions are

$$\frac{R}{\mathrm{pc}} = \frac{1000}{206264.806}\,\frac{D}{\mathrm{kpc}}\,\frac{\alpha}{\mathrm{arcsec}}, \tag{1}$$

$$\frac{\sigma_\perp}{\mathrm{km\,s^{-1}}} = 4.74047046\,\frac{D}{\mathrm{kpc}}\,\frac{\sigma_\mu}{\mathrm{mas\,yr^{-1}}}. \tag{2}$$

The radial and tangential proper-motion components are projected components, not radial and tangential velocities in three-dimensional space. Line-of-sight dispersions already expressed in physical velocity units require no proper-motion conversion. Native measurements and asymmetric errors are retained alongside the display conversions.

For a vector of converted quantities $\mathbf{y}(D)$, a shared distance error contributes, to first order,

$$C_{ab}^{(D)} = \frac{\partial y_a}{\partial D}\frac{\partial y_b}{\partial D}\sigma_D^2. \tag{3}$$

This is one covariance contribution, not the complete uncertainty model. Independence of distance and the measured quantities is not guaranteed, particularly for a kinematic distance. Joint propagation must retain any cross-covariances. Our plots condition on a catalog distance and show converted measurement errors only; distance derivatives and uncertainty are recorded separately. They are not added as independent errors in each radial bin.

Using total mass M and three-dimensional half-mass radius r_h from the same catalog view, define comparison scales

$$r_0 = r_h, \qquad v_0 = \sqrt{\frac{GM}{r_h}}, \qquad t_0 = \frac{r_h}{v_0}. \tag{4}$$

We use $G = 4.30091727 \times 10^{-3}\,\mathrm{pc\,(km\,s^{-1})^2}\,M_\odot^{-1}$ and convert $1\,\mathrm{pc}/(\mathrm{km\,s^{-1}}) = 0.97779222\,\mathrm{Myr}$. The quantities R/r_h and $\sigma_\perp/v_0$ provide common comparison coordinates. This does not impose a shared density profile or make the clusters homologous. Neither v_0 nor t_0 is a measured dispersion, escape speed, or relaxation time. Uncertainty in the joint mass, radius, and distance scales is not propagated into these display plots.

C. Joint host parameters

The inspected posterior products share the coordinate pattern

$$\boldsymbol{\theta}_c = (W_0, M, r_h, \ell_a, g_\mathrm{t}, \delta, s^2, F, \alpha_1, \alpha_2, \alpha_3, f_\mathrm{ret}, D)_c, \tag{5}$$

for cluster c. Here ℓ_a is the base-10 logarithm of the dimensionless anisotropy radius, g_t controls truncation, and δ governs the mass dependence of the velocity scale. These quantities belong to the specified lowered-isothermal and stellar-evolution model [5, 6]. The mass-function slopes α_i and black-hole retention parameter f_ret retain that model's conventions.

The stored retention parameter is a percentage, not the present-day black-hole mass fraction. The coordinates s^2 and F are likelihood nuisance parameters rather than forces or velocity factors. The symbol $g_{\rm t}$ is separate from the geometric region function and Rosenbluth potentials below.

We normalize each stored log weight ℓ_k^w as

$$w_k = \frac{\exp(\ell_k^w - \ell_{\max}^w)}{\sum_l \exp(\ell_l^w - \ell_{\max}^w)}. \tag{6}$$

Moments and marginal quantiles are computed directly from complete weighted rows, without resampling or evaluating a host model. For example,

$$\bar{\boldsymbol{\theta}} = \sum_k w_k \boldsymbol{\theta}_k, \tag{7}$$

$$\mathsf{C}_\theta = \sum_k w_k (\boldsymbol{\theta}_k - \bar{\boldsymbol{\theta}})(\boldsymbol{\theta}_k - \bar{\boldsymbol{\theta}})^{\mathsf{T}}. \tag{8}$$

This is the covariance of the weighted empirical posterior; it is not a finite-sample unbiased estimator. Marginal quantiles use interpolation in the sorted weighted cumulative distribution. Nuisance coordinates remain in the joint records, and marginal summaries do not define independent parameter draws.

Exact observation-file and software-revision identifiers were not found in the inspected run metadata. Historical candidate inputs are therefore documented without claiming a reproduced fit. Current resources cannot automatically replace the inputs to an older posterior. The distinct NGC 6397 no-black-hole alternative discussed in Ref. [7] has a released summary and figures but no corresponding chain in the inspected release.

III. OBSERVATIONAL FINDINGS

A. Coverage and source reconciliation

Table I summarizes the inspected products. The six original Gaia field catalogs contain 621,878 rows, with no repeated source identifier within a cluster. The HST products provide 107 radial bins and 321 component records. Combined, projected radial, and projected tangential dispersions in a bin share stars; the combined value is not a third independent measurement. The released Gaia curves contain 101 grid points per cluster, or 606 correlated samples in total.

The separate compilation contains 418 kinematic rows, of which 20 require reconciliation: four zero-radius entries for NGC 1851, one all-zero entry for ω Centauri, and 15 NGC 6397 entries whose dispersion equals both quoted errors. Four NGC 1851 records also share one radius, source, and component key. These observations motivate flags, not guessed corrections or a claim that every flagged value is physically impossible. The compilation is retained for comparison and is not treated as an approved likelihood input. The 422 surface-profile records likewise retain their unresolved absolute mass normalization.

TABLE I. Source coverage. Gaia field rows include nonmembers. HST bins refer to the original proper-motion profiles; the combined, radial and tangential quantities share stars. Flagged rows belong to the separate 418-row compilation and are not corrected or used as likelihood inputs.

Cluster	Gaia field rows	HST bins	Flagged rows
47 Tuc	144,761	28	0
NGC 1851	12,688	20	4
NGC 3201	63,171	6	0
ω Cen	228,055	40	1
M4	69,417	6	0
NGC 6397	103,786	7	15
Total	621,878	107	20




FIG. 1. Spans of original HST median-bin radii in common comparison coordinates. Numbers at the right give radial-bin counts. Dashed lines mark the intersection of the six spans at fixed catalog scales. They do not represent bin edges or a common three-dimensional domain.

Figure 1 shows the variation in radial sampling. At the catalog point scales, the intersection of the six HST median-radius spans is approximately 0.081–0.251 in projected R/r_h. This is not exact shared annular coverage: most HST bin edges are unavailable, and a span does not establish sampling density between its endpoints. No common radial interpolation or likelihood is constructed.

B. Structural scales and projected kinematics

Table II reports the selected structural estimates and derived scales. Catalog mass ranges from $8.88 \times 10^4\,M_\odot$ for M4 to $3.32 \times 10^6\,M_\odot$ for ω Centauri; the listed central escape speeds range from 18.6 to 61.9 km s^{-1}. These are conditional model estimates, not a measured retention law for merger remnants. Retention would additionally depend on position, inherited progenitor motion, recoil direction, external gravity, and the criterion used to classify escape.

The derived time scales span approximately 0.155–

TABLE II. Structural-catalog estimates and derived display scales. The selected catalog path supplies the quoted mass and distance errors, but no errors for r_h or $v_{\rm esc,0}$; missing uncertainty is not zero. Model systematics and joint scale uncertainty are not included. The gravitational scales v_0 and t_0 are defined in Eq. (4); t_0 is not a friction time.

Cluster	D (kpc)	$M/(10^5 M_\odot)$	r_h (pc)	$v_{\rm esc,0}$ (km s^{-1})	v_0 (km s^{-1})	t_0 (Myr)
47 Tuc	4.52 ± 0.03	8.900 ± 0.050	6.24	51.0	24.77	0.246
NGC 1851	11.95 ± 0.13	2.780 ± 0.040	3.11	42.7	19.61	0.155
NGC 3201	4.74 ± 0.04	1.800 ± 0.030	8.37	18.6	9.62	0.851
ω Cen	5.43 ± 0.05	33.200 ± 0.100	9.85	61.9	38.07	0.253
M4	1.85 ± 0.02	0.888 ± 0.011	3.81	19.5	10.01	0.372
NGC 6397	2.48 ± 0.02	0.901 ± 0.009	3.17	23.4	11.06	0.280

0.851 Myr. They establish dimensional comparison units, not return or damping times. Radius and escape-speed errors are not supplied with these catalog entries; the quoted mass and distance errors also do not capture every model systematic. Mixing a catalog mass with a radius or distance from a different fit would create an unsupported joint parameter combination.

Figure 2 compares original HST radial and tangential dispersions at the catalog scales. It displays the available projected constraints without fitting a common profile. Tracer selection, scale uncertainty, calibration, and radial sampling differ across clusters. Consequently the figure neither demonstrates universal kinematics nor statistically rejects them. HST transformations can remove sky-plane streaming signals [8]; absence of such a signal in these dispersion products is not evidence of zero cluster rotation.

C. Posterior dependence and model sensitivity

Table III reports selected marginal summaries of the published standard-fit samples. The effective sample sizes, calculated as $1/\sum_k w_k^2$, reproduce all six printed source values and range from approximately 5,171 to 6,889. Twelve parameter rows differ beyond three-decimal rounding in a mean or standard deviation. The original summary procedure averages statistics from resampled runs, whereas Eqs. (6)–(8) describe the original weighted sample. This estimator distinction permits close agreement without rounding-identical moments; it does not establish exact source-fit reproduction.

Mass and radius have weighted correlations of 0.900 in NGC 3201 and 0.944 in ω Centauri. Independent sampling of their marginal distributions would discard these dependencies. Because the scales in Eq. (4) involve ratios and products of these parameters, uncertainty should be propagated from complete host realizations. This requirement also applies to quantities derived from a realization, including its potential and species distributions.

Figure 3 compares the published multimass marginal estimates with the structural catalog. For example, NGC 6397 has catalog $r_h = 3.17$ pc, compared with a standard-fit posterior median 4.887 pc and marginal 16th–84th interval $[4.747, 5.044]$ pc. The latter is not the unavailable alternative no-black-hole chain. This difference identifies sensitivity to model and data choices; it is neither temporal expansion measured between releases nor a statistical rejection using independent datasets.

D. What the comparison establishes

The analysis supports a common field and unit convention with a distinct joint host distribution for each cluster. It does not establish six interchangeable scattering backgrounds. NGC 3201 and 47 Tuc provide initial candidates for reproducing a published host and checking its projected observables, with rotation explicitly assessed for 47 Tuc. NGC 1851 and M4 extend the structural comparison. NGC 6397 tests sensitivity in a core-collapsed system, while ω Centauri requires particular attention to rotation, flattening, and mass-dependent tracer selection [7, 9].

Published density and cumulative-mass bands do not by themselves provide mutually consistent local host realizations. Independently assembling median profiles could violate the consistency between mass density, gravity, and velocity structure. These observational findings constrain how the following transport problem should be supplied with inputs. Section V evaluates a specific reconstructed realization, while preserving the distinction between numerical consistency and physical calibration.

IV. STEP 1: RECOIL TRANSPORT

A. Matter distributions and reference frame

Use physical units and cluster-relative coordinates in translating, nonrotating axes. Let $\mathbf{R}(t)$ be the reference position in an inertial frame, and let $(\mathbf{r}, \mathbf{v})$ denote the remnant's relative position and velocity. The host description Θ contains its matter distributions, gravitational boundary conditions, and any supplied temporal evolution.

For species j with individual mass m_j, define a number phase-space distribution $F_j(\mathbf{r}, \mathbf{u}, t \mid \Theta)$. Thus $F_j\, d^3r\, d^3u$ is an expected object count. Its dimensions are inverse volume times inverse velocity cubed. The number and mass densities




FIG. 2. Original projected radial and tangential HST proper-motion dispersions in common dimensionless coordinates. Circles and squares distinguish components. Five clusters use the bright-tracer profiles of Libralato et al. [8]; ω Centauri uses the full-magnitude-range profiles of Häberle et al. [9]. Error bars retain the source measurement-error convention and condition on catalog point scales. Shared distance and joint structural uncertainty are not fully propagated. Components and bins need not be statistically independent.

TABLE III. Direct weighted marginal medians and 16th–84th percentiles from the published standard-fit joint samples of Dickson et al. [6, 7]. These are conditional model estimates, not new fits. The NGC 6397 row is the standard fit; its distinct no-black-hole alternative lacks a released chain in the inspected product. Correlations refer to the complete weighted samples.

Cluster	$M/(10^5 M_\odot)$	r_h (pc)	D (kpc)	Corr(M, r_h)
47 Tuc	9.045 [8.996, 9.094]	6.714 [6.679, 6.749]	4.420 [4.408, 4.431]	0.327
NGC 1851	3.250 [3.211, 3.288]	3.712 [3.670, 3.757]	12.183 [12.106, 12.268]	-0.235
NGC 3201	1.757 [1.699, 1.814]	9.598 [9.070, 10.152]	4.636 [4.603, 4.671]	0.900
ω Cen	30.824 [30.693, 31.489]	9.419 [9.402, 9.529]	5.300 [5.287, 5.327]	0.944
M4	0.911 [0.894, 0.929]	3.920 [3.859, 3.982]	1.852 [1.838, 1.866]	0.316
NGC 6397	1.084 [1.061, 1.109]	4.887 [4.747, 5.044]	2.436 [2.422, 2.452]	0.782

are

$$n_j(\mathbf{r}, t) = \int F_j(\mathbf{r}, \mathbf{u}, t)\, d^3u, \tag{9}$$

$$\rho_j(\mathbf{r}, t) = m_j n_j(\mathbf{r}, t), \qquad \rho_c = \sum_j \rho_j. \tag{10}$$

The cluster potential satisfies

$$\nabla^2 \Phi_c = 4\pi G \rho_c, \tag{11}$$

together with specified boundary conditions. The tracked remnant's self-field is excluded. An unresolved binary can be represented by its center of mass only on encounter scales supporting that approximation; its mass must not also enter through separately counted components. A continuous mass spectrum requires a mass integral or a declared binning convention.

The same distributions determine streaming and velocity




FIG. 3. Structural-catalog estimates (open squares) and published standard-fit posterior marginals (circles with 16th–84th intervals). The mass axis is logarithmic. Catalog mass and distance errors are shown where provided; catalog radius errors are unavailable. Source observations overlap, and model assumptions differ. The NGC 6397 standard fit is kept distinct from its no-black-hole alternative. The comparison measures dependence on the adopted inference, not cluster evolution.

dispersion:

$$\overline{\mathbf{u}}_j = \frac{1}{n_j}\int \mathbf{u} F_j \, d^3u, \tag{12}$$

$$S_{j,ik} = \frac{1}{n_j}\int (u_i - \overline{u}_{j,i})(u_k - \overline{u}_{j,k}) F_j \, d^3u. \tag{13}$$

These moments are defined where $n_j > 0$. Distinct species may have different concentrations, streaming motions, and anisotropies. Density and a scalar dispersion do not generally determine the full velocity distribution needed for a collision calculation.

For an external gravitational potential $\Phi_{\rm ext}$, the relative external acceleration is

$$\mathbf{a}_{\rm ext,rel} = -\nabla\Phi_{\rm ext}(\mathbf{R}+\mathbf{r},t) - \ddot{\mathbf{R}}(t). \tag{14}$$

If the reference orbit follows $\ddot{\mathbf{R}} = -\nabla\Phi_{\rm ext}(\mathbf{R},t)$, Eq. (14) is the differential tidal acceleration. There are no Coriolis terms in these nonrotating axes. Internal stellar rotation remains possible through F_j.

This is a test-remnant description in a supplied background. Poisson's equation does not evolve the host. A frozen host needs a validity interval; an evolving host needs mutually consistent distributions and potential at each epoch. Backreaction, collective response, and energy feedback require additional equations and are not obtained by introducing a time argument alone.

B. Merger-conditioned initial conditions

Set elapsed time zero at the remnant-producing merger. The launch conditions are

$$\mathbf{r}(0) = \mathbf{r}_{\rm m}, \qquad \mathbf{v}(0) = \mathbf{v}_{\rm COM} + \mathbf{v}_{\rm k}, \tag{15}$$

$$m_{\rm r}(0) = m_{\rm rem}. \tag{16}$$

Here $\mathbf{v}_{\rm COM}$ is the progenitor binary's center-of-mass velocity, not its internal orbital velocity. The recoil vector $\mathbf{v}_{\rm k}$ must be transformed into the cluster frame before addition. Equation (15) uses nonrelativistic cluster kinematics; a relativistic merger calculation supplies recoil and remnant mass.

Let $\mathcal{D}$ denote observational or simulation constraints and $\mathcal{M}$ the merger-selection rule and relevant assembly information. The initial law is joint:

$$z_0 = (\mathbf{r}_{\rm m}, \mathbf{v}_{\rm COM}, \mathbf{v}_{\rm k}, m_{\rm rem}), \tag{17}$$

$$z_0 \sim \pi_0(z_0 \mid \Theta, \mathcal{D}, \mathcal{M}). \tag{18}$$

No independence among these variables or from the host is imposed. In particular, the initial specific angular momentum $\mathbf{r}_{\rm m}\times(\mathbf{v}_{\rm COM}+\mathbf{v}_{\rm k})$ need not vanish. A noncentral trajectory need not pass through the exact center.

A retained-only population is meaningful only after specifying the escape criterion and the interval over which retention is assessed. Initial negative cluster-relative energy does not by itself establish permanent retention in an evolving tidal environment. The base transport process is defined before any selection on future survival. Statistics conditioned on that survival require conditioning or reweighting its full path law; changing the launch distribution alone while leaving subsequent stochastic evolution unchanged is insufficient.

C. Velocity drift and diffusion

Define the smooth acceleration $\mathbf{a}_g = -\nabla\Phi_c + \mathbf{a}_{\rm ext,rel}$. A local diffusion description takes the Itô form

$$d\mathbf{r} = \mathbf{v}\,dt, \tag{19}$$

$$d\mathbf{v} = (\mathbf{a}_g + \mathbf{A})\,dt + \mathbf{B}\,d\mathbf{W}, \tag{20}$$

$$\mathbf{D} = \mathbf{B}\mathbf{B}^{\mathsf{T}}. \tag{21}$$

The three Wiener components satisfy $\mathbb{E}[dW_i\,dW_k] = \delta_{ik}\,dt$. After subtracting smooth gravity, the first and second collisional increments define

$$A_i = \lim_{\Delta t\to 0} \frac{\mathbb{E}[\Delta v_i^{\rm coll} \mid \mathbf{r}, \mathbf{v}, t]}{\Delta t}, \tag{22}$$

$$D_{ik} = \lim_{\Delta t\to 0} \frac{\mathbb{E}[\Delta v_i^{\rm coll}\Delta v_k^{\rm coll} \mid \mathbf{r}, \mathbf{v}, t]}{\Delta t}. \tag{23}$$

These limits belong to a coarse-grained approximation; they do not resolve the duration of each gravitational encounter. The dimensions of $\mathbf{A}$ and $\mathbf{D}$ are acceleration and velocity squared per time, respectively. Once $\mathbf{A}$ is the complete Itô first moment, an additional noise-induced drift must not be appended.

Between discrete transitions, remnant mass is fixed on each trajectory. At fixed $Q = (m_{\rm r}, \Theta, \mathcal{D}, \mathcal{M})$, let $p(\mathbf{r}, \mathbf{v}, t \mid Q)$ denote its phase-space density. The interior evolution is

$$\begin{aligned}\partial_t p = &-\partial_{r_i}(v_i p) - \partial_{v_i}[(a_{g,i} + A_i)p] \\ &+ \frac{1}{2}\partial_{v_i}\partial_{v_k}(D_{ik}p).\end{aligned} \tag{24}$$

Repeated Cartesian indices are summed. The factor of one half follows from Eq. (21). Initial, boundary, and termination conditions are needed to define a solution. When launch masses vary, each mass-dependent operator must retain its own conditioning; replacing it by an operator evaluated at a mean mass is not generally equivalent.

D. A candidate weak-scattering closure

For local, uncorrelated weak deflections, a leading-logarithm Rosenbluth closure supplies both coefficients from the same matter distribution [1, 2]. At fixed position and time, define

$$h_j(\mathbf{v}) = \int \frac{F_j(\mathbf{r}, \mathbf{u}, t)}{|\mathbf{v} - \mathbf{u}|}\,d^3u, \tag{25}$$

$$g_j(\mathbf{v}) = \int F_j(\mathbf{r}, \mathbf{u}, t)|\mathbf{v} - \mathbf{u}|\,d^3u. \tag{26}$$

For the number normalization in Eq. (9),

$$A_i = 4\pi G^2 \sum_j m_j(m_{\rm r} + m_j)\ln\Lambda_j\,\partial_{v_i}h_j, \tag{27}$$

$$D_{ik} = 4\pi G^2 \sum_j m_j^2 \ln\Lambda_j\,\partial_{v_i}\partial_{v_k}g_j. \tag{28}$$

A mass-normalized distribution instead absorbs a factor of m_j into each Rosenbluth potential. Mixing those conventions would change the collision coefficients.

The Coulomb logarithm $\ln\Lambda_j = \ln(b_{{\rm max},j}/b_{{\rm min},j})$ represents a declared range of encounter impact parameters. It is held fixed during the velocity integrals and differentiations above. Arbitrary velocity-dependent cutoffs require a consistent derivation rather than an unqualified substitution. Neither cutoff values nor a prescription are selected here.

Writing $\mathbf{w} = \mathbf{v} - \mathbf{u}$, the derivative kernels are

$$\partial_{v_i}h_j = -\int F_j \frac{w_i}{w^3}\,d^3u, \tag{29}$$

$$\partial_{v_i}\partial_{v_k}g_j = \int F_j\left(\frac{\delta_{ik}}{w} - \frac{w_i w_k}{w^3}\right)d^3u. \tag{30}$$

For nonnegative distributions and positive Coulomb logarithms, $\mathbf{D}$ is symmetric and positive semidefinite. Streaming and anisotropy enter through F_j; a Maxwellian is not imposed. Mean collisional drift need not oppose the cluster-frame velocity in every anisotropic or streaming population.

The closure excludes strong encounters and collective or long-range correlations. Its applicability depends on the scales and populations contributing to relaxation. Kicked-black-hole calculations showing departures from simplified friction models [3] therefore motivate independent validation, not an assumption that adding these coefficients is sufficient.

E. Energy, termination, and probability accounting

For fixed remnant mass, define specific cluster-relative energy

$$\epsilon = \frac{1}{2}|\mathbf{v}|^2 + \Phi_c(\mathbf{r}, t). \tag{31}$$

Itô's rule gives

$$\begin{aligned}d\epsilon = &\left[\partial_t\Phi_c + \mathbf{v}\cdot\mathbf{a}_{\rm ext,rel} + \mathbf{v}\cdot\mathbf{A} + \frac{1}{2}\,\mathrm{tr}\,\mathbf{D}\right]dt \\ &+ \mathbf{v}^{\mathsf{T}}\mathbf{B}\,d\mathbf{W}.\end{aligned} \tag{32}$$

The diffusion contribution is required. Individual trajectories can gain energy, and this energy is not a conserved tidal escape criterion. No condition resets position or velocity upon core entry.

For a common physical horizon T, define the modeled endpoint

$$T_* = \min\{T, \tau_{\rm esc}, \tau_{\rm strong}, \tau_{\rm valid}\}. \tag{33}$$

The stopping times represent declared escape, a strong encounter requiring an unavailable transition model, and loss of validity of the supplied description. A strong-encounter stop needs a detector, resolved objects, or a justified event model; missing encounter information is not evidence of no encounters. Weak scattering and explicitly resolved interactions must partition deflections without double-counting.

Let $p_{\rm alive}(\mathbf{r},\mathbf{v},t\mid Q)$ include only paths with $t<T_*$. Then

$$\int p_{\rm alive}\,d^3r\,d^3v = S(t\mid Q) = \Pr(t<T_*\mid Q). \tag{34}$$

Survival here means continued inclusion in the transport description. It does not mean that the black hole ceases to exist when a path is stopped. The transport operator requires the corresponding absorbing boundary or justified killing mechanism. A history-dependent stop requires an augmented state or a justified history-dependent treatment; it does not automatically close Eq. (24) in position and velocity.

The density in Eq. (34) is conditional on fixed remnant mass. A joint mass density is obtained by multiplication with the initial mass marginal π_m:

$$\begin{aligned} p^{\rm joint}_{\rm alive}&(\mathbf{r},\mathbf{v},m,t\mid\Theta,\mathcal{D},\mathcal{M})\\ &= \pi_m(m\mid\Theta,\mathcal{D},\mathcal{M})\,p_{\rm alive}(\mathbf{r},\mathbf{v},t\mid Q_m), \end{aligned} \tag{35}$$

where $Q_m=(m,\Theta,\mathcal{D},\mathcal{M})$. Integrating a fixed-mass conditional density over mass without this weight would not represent the launch ensemble.

V. RECONSTRUCTED ENVIRONMENT AND LOCAL SCATTERING

A. A fixed joint host realization

The first calculation asks whether a published joint host realization can consistently supply the gravitational field and the distributions entering Eqs. (27) and (28). We select the NGC 3201 row with the largest normalized posterior quadrature weight in Eq. (6), retaining all thirteen coordinates. This deterministic selection has weight 0.001764; it is neither a maximum-likelihood estimate nor a posterior mode. A single selected row does not represent the full host posterior or establish a typical cluster environment.

We reconstruct the spherical, nonrotating, multimass lowered-isothermal model associated with the published fit [5–7]. The observation bundle supplies age 11.5 Gyr and metallicity [Fe/H] = −1.59. Historical stellar-evolution and distribution-function implementations are pinned with their mass-bin and retention conventions. Their exact identity with the original fitting run cannot be established from the released metadata. This is a reproducible compatible reconstruction, not an exact rerun of the original likelihood analysis.

Writing $f_j=m_jF_j$ for a mass-normalized distribution, the reconstructed family has the form

$$f_j(r,\mathbf{u}) = \mathcal{A}_j\exp\!\left[-\frac{L^2}{2r_{a,j}^2s_j^2}\right]e^{\mathcal{E}_j}P(g_{\rm t},\mathcal{E}_j),\qquad \mathcal{E}_j>0, \tag{36}$$

and is zero otherwise. Here $L=|\mathbf{r}\times\mathbf{u}|$, $\mathcal{E}_j=[\Psi(r)-u^2/2]/s_j^2$, and P is the regularized lower incomplete gamma function. The normalizations $\mathcal{A}_j$, velocity scales s_j, and species masses come from the same self-consistent solution. The local number distribution used in the collision integrals is explicitly $F_j=f_j/m_j$. The tangential second moment in the numerical host is the sum of two components, so $\beta_j=1-\langle u_t^2\rangle_j/(2\langle u_r^2\rangle_j)$.

The relative potential is referenced to the distribution's truncation radius r_t. For the isolated spherical continuation with physical potential zero at infinity,

$$\Psi(r) = \Phi_{\rm c}(r_t)-\Phi_{\rm c}(r), \tag{37}$$

$$\Phi_{\rm c}(r) = -\Psi(r)-\frac{GM}{r_t},\qquad r\le r_t, \tag{38}$$

$$u_{\rm cut}(r) = \sqrt{2\Psi(r)},\qquad v_{\rm esc,\infty}(r) = \sqrt{2\Psi(r)+2GM/r_t}. \tag{39}$$

Thus the field-star velocity cutoff is distinct from isolated escape to infinity, and neither is automatically a tidal escape criterion. We preserve the source model's $G = 0.004302\,\mathrm{pc\,(km\,s^{-1})^2}\,M_\odot^{-1}$ throughout its gravity and collision calculation. Its small difference from the catalog-display constant does not enter as an inconsistent force normalization.

The selected realization has $M = 1.7806\times10^5\,M_\odot$, $r_h = 10.325$ pc, and $r_t = 981.0$ pc. Its central density is $2.146\times10^3\,M_\odot\,\mathrm{pc}^{-3}$, with $u_{\rm cut}(0) = 17.410\,\mathrm{km\,s^{-1}}$ and $v_{\rm esc,\infty}(0) = 17.455\,\mathrm{km\,s^{-1}}$. The extended formal truncation radius is not adopted as an empirical boundary for NGC 3201. Section VI tests an external-field boundary and an accuracy domain; a physically calibrated outer transport model remains necessary. The separate $34.807\,\mathrm{km\,s^{-1}}$ parameter in the source stellar-evolution prescription is not the present reconstructed escape speed.

B. Independent numerical and projected checks

The reconstruction contains sixteen physical mass bins: nine main-sequence bins, five white-dwarf bins, one neutron-star bin, and one black-hole bin. No artificial tracer mass is appended for this observation bundle. We independently integrate density to recover mass and half-mass radius, compare the potential gradient with enclosed mass, and integrate the force to recover the relative potential. The virial check compares $2K+W$ with W, using density-weighted radial and both tangential velocity moments. Positive densities and pressures are interpolated in logarithms to resolve the steep tails of segregated species. Intervals whose cubic derivative reverses the nodal trend use a shape-preserving interpolant, preventing artificial outer-density shells while preserving continuity at mesh nodes.

Independent velocity quadrature recovers the density and raw second moments of every species at $r/r_h = 0, 0.1, 1, 3$. Projected luminous-tracer moments are calculated by line-of-sight integration, with the correct radial and tangential projection factors. Table IV reports the numerical errors. The adopted host checks require relative errors below 10^{-3}, or 2×10^{-3} for projected luminous-tracer quantities. These tolerances test implementation accuracy; they are not observational uncertainty intervals.

Figure 4 compares the reconstruction with the released projected posterior envelopes over $0.01\le R/r_h\le3$. At the 32 retained source radii, line-of-sight and radial proper-motion dispersions fall within the published central 95.45% envelopes.

TABLE IV. Numerical checks for the fixed NGC 3201 reconstruction. Errors are relative, except the potential error is normalized to its central value. Force checks cover $10^{-3} < r/r_h < 10$; DF checks use four radii. Projected quantities refer to the luminous tracer, not empty remnant tails. These checks establish numerical consistency, not physical calibration.

Check	Maximum error
Integrated mass	3.26×10^{-8}
Half-mass radius	1.20×10^{-6}
Force versus enclosed mass	5.72×10^{-5}
Relative potential	2.35×10^{-7}
Virial balance	9.28×10^{-8}
DF density and raw moments	1.66×10^{-8}
Projected luminous tracer	1.30×10^{-3}
Drift quadrature refinement	5.50×10^{-9}
Diffusion quadrature refinement	1.31×10^{-7}

Tangential proper motion lies inside at 27 radii, and the analogous total-density comparison lies inside at 25. A joint posterior draw need not lie within every pointwise interval; these counts are descriptive, not a goodness-of-fit probability. The comparison neither proves exact historical input compatibility nor replaces an observational refit.

C. Anisotropic coefficient evaluation

We evaluate the weak-scattering kernels without imposing a Maxwellian or a scalar velocity dispersion. To remove the integrable singularity from numerical quadrature, set $\mathbf{v} - \mathbf{u} = s\mathbf{n}$, with $s \geq 0$ and $|\mathbf{n}| = 1$. Equations (29) and (30) become

$$\partial_i h_j = -\int d\Omega \int ds\, F_j(\mathbf{v} - s\mathbf{n}) n_i, \tag{40}$$

$$\partial_i \partial_k g_j = \int d\Omega \int ds\, s F_j(\mathbf{v} - s\mathbf{n})(\delta_{ik} - n_i n_k). \tag{41}$$

The radial interval is the intersection of the ray with the field-star velocity ball $|\mathbf{u}| < u_{\rm cut}$. Positive quadrature weights preserve the positive-semidefinite diffusion tensor. An independent identity is

$$\mathrm{tr}\,\mathsf{D} = 8\pi G^2 \sum_j m_j^2 \ln\Lambda_j h_j. \tag{42}$$

Uniform-ball analytic limits, rotations of an anisotropic distribution, this trace identity, and quadrature refinement test the implementation.

The diagnostic remnant mass is $60\,M_\odot$, a controlled probe rather than an inferred merger remnant. We evaluate twenty local states: four radii $r/r_h = 0, 0.1, 1, 3$, zero speed, and radial or tangential velocities at 0.25 and 0.75 of the local stellar cutoff. Results are reported per unit common constant $\ln\Lambda$; no impact-parameter prescription is selected. With distance in pc and speed in $\mathrm{km\,s^{-1}}$, both raw drift and diffusion require division by 0.97779222 to express their time denominators in Myr. A noise amplitude would instead be obtained by factoring the converted diffusion tensor.

TABLE V. Local weak-scattering coefficients for a $60\,M_\odot$ diagnostic remnant with radial speed $0.25\sqrt{2\Psi(r)}$. Both drift magnitude $a_\parallel = -\mathbf{A}\cdot\widehat{\mathbf{v}}$ and diffusion trace are divided by a common constant $\ln\Lambda$. Units are $(\mathrm{km\,s^{-1}})$/Myr and $(\mathrm{km\,s^{-1}})^2$/Myr, respectively. The BH fraction is its contribution to the diffusion trace in the same host. No physical impact-parameter cutoff is selected.

r/r_h	$a_\parallel/\ln\Lambda$	$\mathrm{tr}\mathsf{D}/\ln\Lambda$	BH fraction
0	8.31×10^{-1}	1.04×10^{0}	8.95×10^{-1}
0.1	1.24×10^{-1}	4.87×10^{-2}	3.20×10^{-4}
1	7.13×10^{-4}	2.35×10^{-4}	4.19×10^{-35}
3	4.27×10^{-5}	8.77×10^{-6}	5.40×10^{-46}

Increasing the angular and ray quadrature orders from 40 to 64 changes drift by at most 5.50×10^{-9} in vector relative norm and diffusion by 1.31×10^{-7} in matrix relative norm on this grid. At zero speed, symmetry requires zero drift. Exactly antipodal quadrature directions are paired before summation, preserving this cancellation for even distributions without suppressing physical drift in a streaming distribution. All diffusion tensors are positive semidefinite, and the trace identity holds to 9.0×10^{-15} relative accuracy. This grid validates local evaluation at the tested states, not interpolation along arbitrary recoil trajectories.

D. Species contributions and the encounter regime

The selected host contains an expected 4.28 black holes, with total mass $61.2\,M_\odot$ in a bin of individual mass $14.31\,M_\odot$. These are model expectations, not observed object counts. At the center this species supplies approximately 27.7% of mass density but 95.8% of the formal diffusion trace at zero test velocity. At $0.1 r_h$, its diffusion fraction has fallen to 8.46×10^{-4}. At nonzero radial speed the decomposition remains strongly position dependent (Table V). Figure 5 compares density and diffusion contributions within this same host; it is not a counterfactual fit obtained by removing black holes.

The enhancement follows from the mass weighting and velocity dependence of Eq. (28). In a continuous description, the encounter contributions enter through the first two velocity-change moments. We use this description as the reference approximation and the species decomposition to identify where an encounter-level comparison is most informative. Finite changes in velocity, correlations between encounters and the response of the scattering population are treated as separate physical effects in Sec. XI. The coefficients here use the adopted host and common constant Coulomb-log normalization.

The reconstruction supplies the same species densities required for Step 2. The radial contrast therefore connects two parts of the problem: the local law changes the trajectory, and the trajectory determines which densities contribute to residence and exposure. Testing this connection requires timed histories under matched launch, encounter and boundary choices. The outer-force calculation below supplies one part of those choices.




FIG. 4. Projected kinematics of the fixed NGC 3201 reconstruction, compared with published posterior medians and central 95.45% model envelopes. These bands summarize correlated model realizations and are not independent measurements. The selected reconstruction retains one complete posterior row; it does not propagate the host posterior.




FIG. 5. Mass-density fractions in the reconstructed host (left) and species contributions to the formal diffusion trace at zero test velocity (right). The radial density curves describe the continuous model; the diffusion bars are four evaluated states. The adopted host has an expected BH count of 4.28. The decomposition identifies spatial regions for the matched encounter comparison in Sec. XI.

VI. OUTER VALIDITY AND TIDAL OPENINGS

A. Boundary definitions and observational scales

The outer boundary constrains the interpretation of the same host used for local scattering. A density-profile limit, a tidal opening, and a stop imposed by an approximation are distinct. Empirical cluster profiles supply fitted limiting radii [13]; tidal dynamics adds a dependence on the Galactic field and the cluster orbit [14]. Even above the critical energy of a stationary tidal problem, escape can be delayed [15]. A crossing cannot therefore be converted directly into a merger-retention probability.

For NGC 3201, Wan et al. [16] distinguish a King limiting radius of 36.1 pc from a cited Jacobi estimate of 83.46 pc. Our captured structural catalog lists 85.86 pc, with its definition referring to Eq. (8) of Webb et al. [17]: an eccentricity- and phase-dependent correction to a pericentric limiting scale. We do not invert that value to infer an instantaneous tidal tensor or average these radii as independent observations.

B. A controlled external-field calculation

We keep the reconstructed spherical mass distribution fixed and prescribe

$$\Phi_G(\mathbf{R}) = v_c^2 \ln(|\mathbf{R}|/R_0), \tag{43}$$

where R_0 sets an irrelevant potential constant. The baseline uses the catalog Galactocentric distance $R = 8.93\,\mathrm{kpc}$ and a flat-rotation benchmark $v_c = 220\,\mathrm{km\,s^{-1}}$. The latter is also used in the illustrative circular comparison of Ref. [16]. This is a specified environmental test, not a fitted Milky Way potential or a reconstruction of NGC 3201's eccentric orbit. The external distance is held separately from the selected host's inferred heliocentric distance; their joint uncertainty is not propagated.

In the translating nonrotating frame of Eq. (14), the linear tidal tensor is $\mathsf{T} = -\nabla\nabla\Phi_G$. Let $\widehat{\mathbf{R}}$ be the Galactic radial direction. For Eq. (43),

$$\mathsf{T} = \frac{v_c^2}{R^2}\left(2\widehat{\mathbf{R}}\widehat{\mathbf{R}}^{\mathsf{T}} - \mathsf{I}\right). \tag{44}$$

For a circular orbit, a frame rotating at $\Omega = v_c/R$ additionally has centrifugal acceleration. In axes along the Galactic radial, orbital tangential, and orbital normal directions, its effective tensor is

$$\mathsf{T}_{\mathrm{eff}} = \frac{v_c^2}{R^2}\,\mathrm{diag}(2, 0, -1). \tag{45}$$

The velocity-dependent Coriolis term still belongs to rotating-frame dynamics. It does not alter a stationary saddle, and neither it nor the centrifugal tensor is inserted into the nonrotating equation.

Balancing cluster gravity with the extensive effective direction gives the circular Hill estimate for an extended spherical host,

$$\frac{GM(<r_J)}{r_J^3} = \lambda_1, \qquad \lambda_1 = \frac{2v_c^2}{R^2}. \tag{46}$$

The point-mass result $(GM/\lambda_1)^{1/3}$ follows when all cluster mass lies inside the saddle. Here we retain the reconstructed enclosed mass. As an independent check, the exact circular inward and outward saddle distances satisfy

$$\frac{GM(<r_\pm)}{r_\pm^3} = \frac{v_c^2}{R(R \pm r_\pm)} + \frac{v_c^2}{R^2}, \tag{47}$$

with the minus sign toward the Galactic center. These conditions compare the tidal expansion with exact differential Galactic and centrifugal accelerations without integrating a trajectory.

C. An accuracy boundary for an isolated control

A tidal opening does not specify where neglecting the Galaxy becomes inaccurate. In the same logarithmic field, the exact maximum differential acceleration at fixed $0 < r < R$ is attained toward the Galactic center:

$$\max_{|\mathbf{r}|=r} |\mathbf{a}_{\mathrm{ext}}| = \frac{v_c^2 r}{R(R-r)}. \tag{48}$$

Dividing by the spherical cluster acceleration defines

$$\eta(r) = \frac{v_c^2 r^3}{GM(<r)R(R-r)}. \tag{49}$$

TABLE VI. Conditional boundary calculations in a spherical logarithmic Galactic field, keeping the NGC 3201 host fixed. r_J uses enclosed mass. $r_{1\%}$ bounds the maximum neglected Galactic acceleration by 1% of the cluster acceleration. Cases are sensitivity settings, not uncertainty intervals; the 13 kpc case is a circular surrogate.

R (kpc)	v_c (km/s)	r_J (pc)	$r_{1\%}$ (pc)	$\eta(3r_h)$
8.93	220	84.13	20.67	2.96%
8.93	198	90.44	22.39	2.40%
8.93	242	78.79	19.22	3.58%
13.00	220	108.79	27.40	1.40%

We propose a conditional accuracy boundary

$$r_\epsilon = \inf\{r > 0 : \eta(r) > \epsilon\}, \tag{50}$$

using $\epsilon = 0.01$ for an isolated-host control and comparing 0.03 and 0.1. The evaluated host has increasing η throughout the relevant interval, so its limiting radius is obtained from $\eta(r_\epsilon) = \epsilon$. This tolerance is an analysis choice, not a physical law. It bounds an instantaneous omitted force under the prescribed field; it does not bound accumulated orbit error, establish a tidally consistent equilibrium, or justify the collision closure.

D. Findings and implications for transport

The baseline gives $r_J = 84.13\,\mathrm{pc}$, compared with 85.77 pc for the point-mass approximation. Exact circular saddles are at 84.00 and 84.27 pc. Agreement in scale with the published 83.46 pc estimate does not independently validate the host or its velocity distribution. The radius is conditional on the adopted Galactic field and one host realization.

The force limits are $r_{1\%} = 20.67\,\mathrm{pc}$, $r_{3\%} = 31.12\,\mathrm{pc}$, and $r_{10\%} = 47.94\,\mathrm{pc}$. At the existing coefficient radii r_h and $3r_h$, the maximum omitted force is 0.175% and 2.962% of cluster gravity, respectively. Thus the outermost local kernel remains a defined integral of its specified distribution, but its use in an isolated trajectory exceeds the proposed 1% force budget. Table VI varies v_c by ±10% and includes a separate 13 kpc circular surrogate discussed in Ref. [16]. These scenarios are assumption tests, not an orbit track or confidence intervals.

The unmodified host places 5.63% of its mass beyond the baseline saddle, while its formal truncation is 981 pc. This is a consistency warning about the outer distribution, not a predicted escaped fraction. Simply removing this material would not yield a self-consistent tidal distribution function. An exterior boundary requires a host appropriate to the imposed tide as well as a remnant event rule.

In the conservative circular Hill problem, the critical specific energy is $E_{J,c} = \Phi_{\mathrm{cl}}(r_J) - \lambda_1 r_J^2/2$. The corresponding central opening speed,

$$v_{\mathrm{open}}(0) = \sqrt{2[E_{J,c} - \Phi_{\mathrm{cl}}(0)]}, \tag{51}$$

is $16.68\,\mathrm{km\,s^{-1}}$. It refers to the remnant's full rotating-frame speed at the center, not its recoil alone. It is neither a measured escape speed nor a retention threshold with friction and




FIG. 6. Baseline boundary diagnostics for the fixed NGC 3201 realization. Left: the maximum differential Galactic force relative to cluster gravity, with the chosen 1% tolerance and two existing local-coefficient evaluation radii. Right: enclosed mass in the unmodified host, compared with the circular extended-host saddle and formal distribution-function truncation. Exterior mass is not classified as escaped by this calculation.

diffusion. Noncircular tides and collisional changes remove the conservation on which this energy argument rests.

Boundary roots have relative residuals below 9.5×10^{-13}. Independent density integration agrees with the enclosed mass at the tested boundaries to 1.2×10^{-7}; thinning the radial mesh changes inferred radii by less than 1.5×10^{-7}. Analytic point-mass and Plummer limits, the exact angular force maximum, and force–potential consistency give additional checks. These errors describe numerics, not Galactic or host uncertainty.

For an isolated control launched strictly inside $r_{1\%}$, its first outward crossing is proposed as $\tau_{\rm valid}$ in Eq. (33). A launch already outside has $\tau_{\rm valid} = 0$; a boundary launch needs a declared side convention. This stop would censor the trajectory and accumulated exposure. It would not be labeled escape or permanent loss of access to the center. A trajectory including the external force has a different validity criterion. Extending the calculation to the tidal outskirts requires a time-dependent Galactic trajectory, an appropriate outer host, and explicit records of departures and possible returns. The sparse-remnant issue identified in Sec. V remains a separate prerequisite for physical collisional transport.

VII. GALACTIC ORBIT AND TIME-DEPENDENT HOST SUPPORT

A. Observed phase point and defined reference orbits

The static boundary calculation does not specify how the tide changes along NGC 3201's orbit. We therefore integrate the cluster's Galactic reference motion and apply the resulting field to the same host as a diagnostic. This calculation evolves neither a kicked remnant nor the cluster's internal distribution.

The database's orbital table supplies a Galactocentric Cartesian point

$$\mathbf{R}_0 = (7.59, -4.65, 0.71)\,\mathrm{kpc},$$
$$\mathbf{V}_0 = (-250.21, -200.78, 149.10)\,\mathrm{km\,s^{-1}}. \quad (52)$$

Its axes point toward the Sun, along solar motion, and north of the Galactic plane. We preserve the published point and its rounding. The table also provides heliocentric distance $D = 4.74 \pm 0.04\,\mathrm{kpc}$, proper motions $(\mu_{\alpha*}, \mu_\delta) = (8.350, -1.976)\,\mathrm{mas\,yr^{-1}}$, and radial velocity $495.38 \pm 0.06\,\mathrm{km\,s^{-1}}$. Both proper-motion errors are $0.005\,\mathrm{mas\,yr^{-1}}$, with correlation 0.18; a full six-dimensional covariance is not supplied. These compiled constraints retain the distance and astrometric sources in Refs. [10, 12].[2]

We adopt Model I of Irrgang et al. [18], a fixed axisymmetric bulge–disk–halo potential, as a reproducible reference used by the catalog's modeling framework. Its Table 1 parameters and exact potential unit $100\,\mathrm{km^2\,s^{-2}}$ are retained. This is not a uniquely inferred Galactic potential. No bar, spiral structure, secular evolution, or Galactic-parameter posterior is added.

Distance changes must alter transverse velocity as well as position. With line-of-sight unit vector $\mathbf{n}$ and proper-motion tangent vectors $\mathbf{e}_\alpha, \mathbf{e}_\delta$ in the same axes,

$$\Delta\mathbf{R} = \Delta D\,\mathbf{n},$$
$$\Delta\mathbf{V} = k_\mu \Delta D(\mu_{\alpha*}\mathbf{e}_\alpha + \mu_\delta \mathbf{e}_\delta), \quad (53)$$

where $k_\mu = 4.74047046$ in the units of Eq. (2). We compare the catalog distance, its quoted error shifts, and the selected

[2] The orbital input uses the database's /newdata/ orbital table, without merging it with the root-path version.

host's 4.54335 kpc distance. The latter retains that host's distance coordinate but is not a joint astrometric–host posterior. The catalog Cartesian anchor preserves its solar-frame transformation; missing cross-covariances and solar-frame uncertainties are not invented.

Each reference orbit obeys $\ddot{\mathbf{R}} = -\nabla\Phi_G$ for 1 Gyr forward in the specified potential. Spherical radial turning points and Galactic-plane crossings are detected as events. This horizon samples repeated phases; it is not a forecast that the internal host remains unchanged for a gigayear. Energy and axial angular momentum are conserved in the adopted Galactic model. Their relative numerical drifts remain below 10^{-8}, and refined integration and time reversal meet the declared position and velocity tolerances.

B. A sufficient isolated-force domain along the orbit

Write $\mathsf{T}(\mathbf{R}) = -\nabla\nabla\Phi_G(\mathbf{R})$. For a bound $B(\mathbf{R}, r)$ satisfying

$$\|\mathsf{T}(\mathbf{R}+\mathbf{s})\|_2 \le B(\mathbf{R}, r) \quad \text{for } |\mathbf{s}| \le r, \tag{54}$$

the exact differential Galactic force obeys

$$|\mathbf{a}_{\rm ext}| \le rB(\mathbf{R}, r). \tag{55}$$

This follows by integrating the force derivative along the segment from $\mathbf{R}$ to $\mathbf{R}+\mathbf{r}$. Appendix A supplies an analytic componentwise bound for the adopted potential. It is valid for all directions in the ball and does not infer a supremum from finitely sampled angles.

A sufficient radius for a chosen omitted-force tolerance solves

$$\eta_B(r,t) = \frac{B(\mathbf{R}(t), r)r^3}{GM(<r)} = \epsilon, \qquad \epsilon = 0.01. \tag{56}$$

This generalizes the static calculation without identifying the radius with escape. It controls an instantaneous neglected acceleration under the supplied host; it is neither a cumulative trajectory-error guarantee nor evidence that the stellar distribution is tidally self-consistent.

C. Frozen-frame scales and forcing timescales

For comparison, define a frame whose radial direction follows $\mathbf{R}$ with minimal instantaneous rotation,

$$\boldsymbol{\omega} = \frac{\mathbf{R}\times\mathbf{V}}{|\mathbf{R}|^2},$$
$$\mathsf{T}_f = \mathsf{T} + |\boldsymbol{\omega}|^2\mathsf{I} - \boldsymbol{\omega}\boldsymbol{\omega}^{\mathsf{T}}. \tag{57}$$

We report r_f from $GM(<r_f)/r_f^3 = \lambda_{\max}(\mathsf{T}_f)$. On this inclined eccentric orbit, r_f is only a frozen-frame diagnostic. The Euler acceleration $-\dot{\boldsymbol{\omega}}\times\mathbf{r}$ is omitted from that definition, and a stationary Jacobi surface does not follow. Actual rotating-frame dynamics would also require Coriolis acceleration. The transport formulation in nonrotating axes continues to use the exact differential Galactic force.

An additional indicator compares internal dynamical time to changes in forcing:

$$\mathcal{A}(r,t) = \sqrt{\frac{r^3}{GM(<r)}}\frac{\|\dot{\mathsf{T}}\|_2}{\|\mathsf{T}\|_2}. \tag{58}$$

Time units are converted consistently. This quantity includes tensor orientation changes in the fixed axes. We separately evaluate variation of the tensor norm alone. Neither indicator is an inferred heating rate or a calibrated adiabatic correction.

D. Findings and limits of the frozen host

The catalog-reference orbit has mean pericenter 8.333 kpc and apocenter 25.362 kpc, compared with catalog estimates 8.33 and 25.82 kpc. The apocenter difference is 1.77%. This is a descriptive scale check; the original averaging procedure and complete model configuration are not reconstructed.

For the selected-host-distance orbit, mean pericenter and apocenter are 8.215 and 23.810 kpc. The sufficient 1% force radius varies from approximately 8.2 to 39.8 pc. Halving output spacing changes its minimum by 0.0674%, giving 8.173 pc; rounding downward to a whole parsec motivates an 8 pc fixed isolated-control domain for this specific calculation. It does not transfer unchanged to another host or Galactic model. Initial points outside a control domain are invalid at time zero; interior trajectories are censored on validity exit, without being labeled escaped.

The bounds on neglected force reach 1.73% at r_h and 29.5% at $3r_h$. The static logarithmic-field limit of 20.67 pc therefore cannot be carried unchanged through the orbit. A calculation retaining exact Galactic forcing does not require this omitted-force stop, but still requires an adequate host and collision model.

The frozen-frame scale spans 62.1–183.9 pc, and the unmodified host puts 1.37–8.74% of its mass beyond it. These are neither stripped-mass estimates nor successive equilibria. The ratio $|\dot{\boldsymbol{\omega}}|/\lambda_{\max}(\mathsf{T}_f)$ reaches 0.67, reinforcing the distinction from a stationary tidal-opening calculation. At r_h, $\mathcal{A}$ reaches 1.16; norm variation alone gives 1.14. Thus rapid forcing is not only a tensor-orientation effect. The indicators identify phases where a slowly adjusting host approximation needs testing rather than supplying that test's outcome.

Self-consistent nonspherical equilibria can be constructed for circular orbits in a frozen tidal field [19]; that construction does not establish equilibrium under an eccentric orbit's changing tide. A consistent evolving host instead requires species equations of the form

$$\partial_t F_j + \mathbf{u}\cdot\nabla_{\mathbf{r}}F_j + \mathbf{a}_g\cdot\nabla_{\mathbf{u}}F_j = C_j[\{F_k\}], \tag{59}$$

coupled to Eq. (11), appropriate boundaries, and specified collisional physics. We have not solved this host-evolution problem. The orbit supplies its external forcing and tests the domain of the fixed-host approximation; it does not repair the outer distribution by deleting exterior stars. The central sparse-remnant scattering issue remains independent of this environmental calculation.

TABLE VII. Galactic reference-orbit diagnostics over the defined 1 Gyr horizon in Irrgang Model I. The host mass profile is fixed. D is heliocentric distance; mean pericenter and apocenter are spherical Galactocentric radii from detected turning events. The intervals show orbit-phase variation of the sufficient 1% isolated-force radius r_ϵ and frozen-frame diagnostic r_f, not posterior uncertainties.

Case	D (kpc)	$\langle R_p \rangle$ (kpc)	$\langle R_a \rangle$ (kpc)	r_ϵ (pc)	r_f (pc)
Catalog distance	4.740	8.33	25.36	9.1–41.8	63.0–193.6
Selected-host distance	4.543	8.22	23.81	8.2–39.8	62.1–183.9
Catalog distance minus quoted error	4.700	8.31	25.03	9.0–41.4	63.0–191.6
Catalog distance plus quoted error	4.780	8.36	25.70	9.1–42.3	63.1–195.7




FIG. 7. The reference orbit using the selected host's distance, with the internal host held fixed only for the diagnostic. (a) Galactic radius and absolute height. (b) Sufficient isolated-force radius and frozen-frame scale; neither is a measured cluster edge. (c) Bounds on the omitted Galactic force at the existing local-coefficient radii. (d) Changes in forcing relative to host dynamical time. The proposed fixed 8 pc control is conditional on this orbit and host, and does not classify escape.

VIII. COMPARISON ACROSS SIX CANDIDATE HOSTS

The same reconstruction and local-coefficient calculation is applied to all six shortlisted clusters. Each reference preserves the original selection rule: the complete joint row with the largest normalized nested-sampling quadrature weight. This is a reproducible conditional reference, not a posterior mode or a representative host. Every reference is checked for mass, force, potential, virial balance, distribution-function moments and projected luminous-tracer consistency before its twenty local scattering states are evaluated. The probe mass, velocity fractions, directions and constant Coulomb-log normalization remain those of Sec. V.

Table VIII separates the selected-host results from empirical summaries of the released derived-model realizations. There are 4,000 such realizations per cluster. The historical compatible analysis code forms these products from equal-weight chains, but the generating seed and mapping to raw nested rows are unavailable. We therefore do not join the derived arrays to nested rows by position. Their quantiles describe the released conditional model product, not a new observational fit or a direct census of individual black holes. In

TABLE VIII. Conditional standard-fit comparison. $N_{\rm BH}^{\rm ref}$ is the expected count in the selected maximum-quadrature-weight joint row. Published intervals are empirical 16th–84th percentiles of 4,000 released derived model realizations per cluster. f_D is the BH contribution to the local diffusion trace at zero remnant velocity, with a common constant Coulomb logarithm. The probe mass is $60\,M_\odot$. These are model quantities, not directly observed counts or residence fractions. The NGC 6397 standard fit is a sensitivity case; its alternative no-BH chain is unavailable.

Cluster	$N_{\rm BH}^{\rm ref}$	Published $N_{\rm BH}$	$m_{\rm r}/M_{\rm BH}^{\rm ref}$	$f_D(0)$ (%)	$f_D(0.1r_h)$ (%)
47 Tuc	65.9	61.9 [51.2, 81.1]	0.1357	81.579	0.01328
NGC 1851	16	15 [13.1, 17.6]	0.3034	99.542	0.0004788
NGC 3201	4.28	0.926 [0, 4.13]	0.981	95.836	0.0846
ω Cen	12,100	12,400 [12,100, 13,000]	0.0003391	99.999	98.52
M4	5.33	6.17 [3.99, 9.65]	0.8782	99.431	0.02131
NGC 6397 (standard)	2.94	3.08 [2.84, 3.31]	3.041	97.793	$< 10^{-4}$

particular, 1,899 of the 4,000 NGC 3201 realizations contain no BH component. An absent subsystem has zero BH scattering contribution, but its half-mass radius and the ratio of probe mass to subsystem mass are undefined.

A separate sensitivity pilot selects eight weighted joint rows per cluster, using categorical probabilities proportional to the released nested-sampling weights and a fixed reproducible seed. Sampling is with replacement and preserves all parameter correlations. Every attempt remains in the record, including an initially failed numerical check; a failed draw is not replaced by a more convenient host. Forty-seven of the 48 attempts satisfy the host checks. One M4 joint row fails to converge to a finite model extent in the compatible reconstruction; it is retained as unresolved, contributes no scattering estimate, and is not replaced. These attempts are a pilot, not a converged posterior estimate of transport uncertainty. Nine zero-speed radii, $r/r_h = 0, 0.001, 0.003, 0.01, 0.03, 0.1, 0.3, 1, 3$, resolve the compact central contribution more closely. At zero velocity, an independent axisymmetric integral evaluates the diffusion trace, checked under 64-to-128 quadrature refinement.

The reference calculations distinguish population size from spatial influence. The BH contribution is substantial at the center in both low-count and massive-reservoir models. In the references for 47 Tuc, NGC 1851, NGC 3201 and M4 it falls sharply by $0.1r_h$, while the ω Cen reference retains a large contribution there. In the NGC 6397 standard fit it falls even more rapidly. A central diffusion fraction cannot establish the importance of BH scattering along a recoil orbit: the trajectory, crossing duration and repeated encounters remain necessary. The fixed probe also has different ratios to the host and BH-reservoir masses, so the comparison does not isolate either ratio as a causal parameter.

The comparison adopts the published standard-fit, spherical, nonrotating multimass distributions [7]. NGC 3201 supplies a conditional sparse-population example, while ω Cen supplies a larger reservoir under the same model family. The NGC 6397 calculation uses its available standard-fit chain; the distinct no-BH alternative is not evaluated because its chain is absent from the inspected release. Rotation, flattening and additional population structure are outside these adopted backgrounds. Complete joint rows preserve the parameter dependence within each distribution. These choices define the physical settings of the controlled transport comparison.

Environmental controls retain a distinct phase-space source row for each cluster. Four distance cases per cluster use the same Galactic potential and coordinate transformation; three static circular controls use each cluster's own catalog Galactocentric radius and common circular speeds of 198, 220 and 242 km s^{-1}. Full phase-space and Galactic-model covariances remain unavailable. Integration and diagnostic-cadence refinement retain the original acceptance limits and record earlier failures. All 24 reference orbits pass their numerical checks. M4 nevertheless differs from the catalog extrema beyond the descriptive 5% threshold: its catalog-distance mean pericenter is 0.772 kpc against 0.64 kpc, a 20.6% difference; its apocenter differs by 2.54%. This discrepancy remains unresolved, and numerical convergence does not validate the assumed Galactic potential or reproduce the catalog averaging procedure. The resulting force domains are conditional instantaneous approximation bounds, not remnant escape surfaces or a proof that the host remains in equilibrium.

IX. STEP 2: RESIDENCE AND ENVIRONMENTAL EXPOSURE

A. Finite-region entry and re-entry

Specify an operational region by a level-set function,

$$C(t) = \{\mathbf{r} : g(\mathbf{r}, t) < 0\}. \tag{60}$$

The function is negative inside, positive outside, and zero on the boundary. This g describes geometry and is distinct from the velocity-space potentials g_j in Eq. (26). A sphere may be represented by $g = |\mathbf{r}| - r_c(t)$, but neither a shape nor a radius is selected. A projected observational core radius is not automatically the required spatial boundary.

For an initially interior launch, define first exit and subsequent re-entry as

$$\tau_{\rm out} = \inf\{t > 0 : g(\mathbf{r}_t, t) > 0\}, \tag{61}$$

$$\tau_{\rm re} = \inf\{t > \tau_{\rm out} : g(\mathbf{r}_t, t) < 0\}. \tag{62}$$

The strict opposite-side conditions exclude an isolated tangential contact. For a continuous path, the infimum identifies the boundary time associated with a sign-changing crossing. An observed re-entry delay is $\tau_{\rm re} - \tau_{\rm out}$.




FIG. 8. BH fraction of the local diffusion trace at zero remnant velocity. Blue curves show the six selected references; gray curves show numerically accepted joint-row pilot realizations. Lines connect evaluated radii and are neither credible bands nor an adopted trajectory interpolator. The horizontal scale is linear near zero and logarithmic beyond $10^{-3} r_h$. All species share the same constant Coulomb-log normalization. The NGC 6397 panel describes the available standard-fit sensitivity case.

For an initially exterior launch, define first entry by

$$\tau_{\mathrm{ent}} = \inf\{t > 0 : g(\mathbf{r}_t, t) < 0\}. \tag{63}$$

A boundary launch requires a declared initial-side convention. Classification may use the first definite side reached, while recording the contact separately. It must not create an artificial return event. A path that never exits has no observed re-entry, and the convention $\inf \varnothing = +\infty$ does not establish that a censored event is physically impossible.

For a differentiable boundary,

$$\frac{d}{dt} g(\mathbf{r}_t, t) = \partial_t g + \mathbf{v}_t \cdot \nabla g. \tag{64}$$

Boundary evolution can change membership without inward remnant motion. Both the crossing and its origin must remain identifiable. Sign changes remain the definition even when the derivative vanishes. Separate inner and outer event surfaces can control numerical recrossings only if their effects are reported; they do not silently redefine $C(t)$.

Only events observed before the modeled endpoint are recorded as completed. Simultaneous crossing and stopping events require an ordering convention. Return flags and termination causes are separate fields: a path can re-enter the region and later escape.

B. Path-integrated observables

The residence accumulated before termination is

$$C_T = \int_0^{T_*} \mathbf{1}_{\{\mathbf{r}_t \in C(t)\}}\, dt. \tag{65}$$

It has units of time and obeys $0 \le C_T \le T_* \le T$. The ratio C_T/T uses a common horizon across realizations. If a description stops before T because subsequent motion is unresolved, this ratio is not a complete-horizon physical residence fraction. The alternative C_T/T_*, for $T_* > 0$, describes the modeled interval and answers a different question.

For each species, define mass-density exposure

$$J_{j,T} = \int_0^{T_*} \rho_j(\mathbf{r}_t, t)\, dt. \tag{66}$$

Its dimensions are mass times inverse volume times time. A normalized exposure additionally requires a declared positive reference:

$$I_{j,T} = \int_0^{T_*} \frac{\rho_j(\mathbf{r}_t, t)}{\rho_{\mathrm{ref},j}(t)}\, dt, \tag{67}$$

$\rho_{\mathrm{ref},j}(t) > 0.$

It has units of time. The reference is a convention, not an automatically inferred central density. Unless it bounds the sampled density, $I_{j,T}/T$ need not lie between zero and one.

Density may vary even while a trajectory remains inside $C(t)$; density outside that region may also be nonzero. Region-restricted exposure can be defined by inserting the same indicator into Eq. (66), but it is a distinct observable. None of these integrals is an encounter count.

C. Time-augmented occupation

Preserve the chronological relation between motion and environment through

$$\mu_T(\mathcal{A}) = \int_0^{T_*} \mathbf{1}_{\{(t,\mathbf{r}_t,\mathbf{v}_t)\in\mathcal{A}\}}\, dt, \tag{68}$$

where $\mathcal{A}$ is a measurable subset of $[0,T]\times\mathbb{R}^3\times\mathbb{R}^3$. The measure has total mass T_* and units of time. Equations (65) and (66) are weighted integrals of that same measure:

$$C_T = \int \mathbf{1}_{\{g(\mathbf{r},t)<0\}}\, \mu_T(dt, d^3r, d^3v), \tag{69}$$

$$J_{j,T} = \int \rho_j(\mathbf{r},t)\, \mu_T(dt, d^3r, d^3v). \tag{70}$$

Removing time labels can destroy the association with an evolving host; removing velocity can destroy information needed for a later dynamical kernel. Aggregated exposure therefore accompanies the trajectory and event record rather than replacing them.

D. Ensemble averages and incomplete histories

At fixed Q, nonnegative path integrands give the identities

$$\mathbb{E}[C_T \mid Q] = \int_0^T dt \int d^3r\, d^3v \times \mathbf{1}_{\{g(\mathbf{r},t)<0\}}\, p_{\mathrm{alive}}(\mathbf{r},\mathbf{v},t \mid Q), \tag{71}$$

$$\mathbb{E}[J_{j,T} \mid Q] = \int_0^T dt \int d^3r\, d^3v \times \rho_j(\mathbf{r},t)\, p_{\mathrm{alive}}(\mathbf{r},\mathbf{v},t \mid Q). \tag{72}$$

If a density conditional on survival is used instead, it must be multiplied by $S(t \mid Q)$. Renormalizing the remaining trajectories silently changes the population being averaged. Equations (71) and (72) provide a future consistency comparison between trajectory accumulation and a transport-density solution.

The event record must distinguish no exit observed, entry or re-entry observed, finite-horizon censoring, declared escape, unresolved strong interactions, and loss of model validity. Escape can compete with return; it is not interchangeable with numerical failure. A spatial boundary crossing counts as escape only under the declared escape rule.

After an unresolved stop, complete-horizon residence and exposure remain unknown. The nonnegative accumulated integrals describe the modeled interval, while extrapolation needs a continuation model. Censoring can depend on the path and environment. Treating it as independent would be an additional statistical assumption requiring justification.

X. CONDITIONING AND CALCULATION REQUIREMENTS

A. Physical and inferential uncertainty

Host uncertainty, launch uncertainty, and physical scattering are different sources of variation. For a trajectory observable O, their joint accounting takes the form

$$\mathbb{E}[O \mid \mathcal{D},\mathcal{M}] = \int d\Theta\, p(\Theta \mid \mathcal{D},\mathcal{M}) \times \int dz_0\, \pi_0(z_0 \mid \Theta,\mathcal{D},\mathcal{M}) \times \mathbb{E}_{\mathrm{path}}[O \mid \Theta, z_0, \mathcal{D},\mathcal{M}]. \tag{73}$$

The inner expectation includes the Wiener forcing and any additional randomness in a supplied stopping-event model. This factorization preserves conditional dependence. Merger selection can change host weights as well as launch properties. Alternative host families require separate conditioning or explicit model probabilities. Region definitions, reference densities, and any selection-dependent horizon must also be specified consistently.

Observed stellar kinematics constrain a host through an inference problem; they do not directly provide the entire potential, hidden remnant populations, or collision coefficients. The posterior summaries in Sec. III quantify dependence within published candidate fits. They do not determine probabilities for competing host families or their merger-conditioned weights.

For a fixed observed cluster c, a compatible reconstructed host law $\Pi_c(d\Theta \mid \mathcal{D}_c)$ would enter a trajectory-observable distribution as

$$\Pr_c(O \in A \mid \mathcal{D}_c,\lambda) = \int \Pr(O \in A \mid \Theta,\lambda)\, \Pi_c(d\Theta \mid \mathcal{D}_c). \tag{74}$$

Here A is a measurable set of observable values, not the drift vector $\mathbf{A}$, and λ retains the declared launch, recoil, collision, boundary, and horizon choices. The integrand includes the launch and path averaging in Eq. (73). This expression is a conditional modeling prescription, not an evaluated prediction. A merger-selected population additionally needs the host reweighting in Eq. (73); the six clusters are not assigned universal population weights.

Each host realization must supply the same F_j to Eqs. (9)–(13) and (25)–(28). The density used for exposure must come from that realization as well. Section V constructs those local functions for one fixed realization and checks its projected predictions. Section VIII extends the calculation to six references and a joint-row pilot; each trajectory comparison will

first be conditioned on one such realization, then propagated through complete joint rows. We use present-day snapshots as specified environments; reconstructing a particular historical merger environment would require its evolution and formation history.

B. Information passed between the two steps

A calculation becomes defined only when its physical inputs and event conventions are supplied. These include the units and frame; species masses and distributions; internal and external gravity with boundary conditions; the joint launch law; collision closure and scale cutoffs; physical horizon; region geometry; and escape, strong-event, and validity criteria. An unspecified input is not a zero-valued effect.

Step 1 must preserve physical times, positions, velocities, remnant-mass labels, host and launch identifiers, original realization weights, and termination causes. A density-based calculation must preserve its conditioning and absorbed or killed probability. Reproducibility requires the source of each input and enough numerical information to reconstruct coefficient evaluation, stochastic realizations, and error control.

Step 2 attaches crossing events, residence, species exposures, and a time-augmented occupation measure. Every summary retains its denominator convention, modeled endpoint, and status. A path that becomes unresolved is not filled by an assumed central rest state. The same host and epoch must supply the density used for exposure and the distribution used to drive transport.

C. Acceptance criteria before physical interpretation

The host and local-coefficient checks in Sec. V establish dimensional consistency, force–potential agreement, number-to-mass normalization, and positive-semidefinite diffusion for the stated realization and evaluation grid. Section XI D verifies conservative trajectories and finite-region occupation. Section XI E adds a conditional stochastic pilot with independent moment, stopping and null controls. Further convergence and calibration requirements remain. The implementation must preserve the Itô convention and account for probability lost at each boundary or stop. Appropriate conservative and analytically justified stochastic limits provide independent checks, including the energy identity in Eq. (32).

Entry and residence calculations must distinguish tangencies from crossings, handle interior, exterior, and boundary launches, and account for moving regions. Brief passages must remain resolved under a declared refinement procedure. Convergence is needed for the relevant distributions and accumulated observables, not solely for a few representative trajectories. Numerical and sampling errors must be separated from uncertainty in physical inputs.

Where both representations exist, path averages and the survival-weighted identities in Eqs. (71) and (72) should agree within declared numerical accuracy. Comparisons testing a physical simplification must hold the conditional population, horizon, region, and remaining inputs fixed. A useful result could identify a regime in which a reduction preserves the observables, or a discrepancy establishing which additional information is required.

Internal agreement does not calibrate the collision model. Comparison with independently resolved dynamics appropriate to the host and encounter regime is needed before interpreting predictions as quantitatively reliable. Host and coefficient quadrature controls are specified for the completed calculation. A deterministic orbit solver and diagnostic launches are supplied in Sec. XI D. The matched pilot supplies independent stochastic ensembles and explicit precision targets. A physically calibrated encounter partition and a resolved-dynamics calibration test remain to be supplied.

XI. RESIDENCE HYPOTHESIS AND CONTROLLED ENCOUNTER COMPARISON

A. The observable and the hypothesis

The theoretical target is the accuracy of a transport approximation for a specified observable. At fixed host, launch law, region, external field and horizon, we compare a local encounter process with a diffusion process constructed from the same encounter law. The hypothesis is that finite encounter increments become important for residence where they substantially change the expected remaining occupation, and where the remnant actually samples those states. This is more specific than associating accuracy with a total BH count: it depends on the encounter-rate distribution, the geometry of returning trajectories and the time available for subsequent excursions.

The current six-cluster profiles locate candidate regions for this comparison. The compact central contribution in several adopted hosts and its greater radial extent in the ω Cen reference motivate different launch radii, speeds and directions. Zero-BH realizations supply a control with no BH encounter contribution. These findings select physical regimes; the hypothesis requires the matched calculation below.

B. A common encounter law and its diffusion limit

Let $x = (\mathbf{r}, \mathbf{v})$, and let $\mathcal{B}$ denote the species whose encounter representation is varied. The initial comparison takes $\mathcal{B}$ to be the BH component. Other species retain the same treatment in both calculations. For local independent encounters with single objects, adopt the differential rate

$$d\Gamma_j = F_j(\mathbf{r}, \mathbf{u}, t)\, w\, d^3u\, b\, db\, d\varphi, \qquad w = |\mathbf{v} - \mathbf{u}|, \tag{75}$$

where b is the impact parameter of the asymptotic relative orbit and $0 \le \varphi < 2\pi$. A declared finite impact-parameter domain makes the rate finite and specifies the spatial scale on which the local description is used. For an elastic Newtonian

two-body encounter,

$$b_{90,j}(w) = \frac{G(m_{\rm r}+m_j)}{w^2}, \qquad \vartheta_j = 2\arctan\frac{b_{90,j}}{b},$$
$$\boldsymbol{\xi}_j = \frac{m_j}{m_{\rm r}+m_j}(\mathbf{w}'-\mathbf{w}), \qquad |\mathbf{w}'| = w, \tag{76}$$

where $\mathbf{w} = \mathbf{v} - \mathbf{u}$ and $\mathbf{w}'$ is rotated by the deflection angle ϑ_j and impact azimuth. This changes the remnant velocity by $\boldsymbol{\xi}_j$ while conserving the pair's energy and momentum in the isolated scattering problem. The allowed closest approaches must be declared; binaries, relativistic capture and merger transitions require additional interaction laws. The impulse description assumes an encounter duration short compared with the resolved orbital and background-evolution times.

The velocity-jump rate measure is the image of Eq. (75) under Eq. (76):

$$\nu_{t,x}(d\boldsymbol{\xi}) = \sum_{j\in\mathcal{B}}\int d\Gamma_j\,\delta_{\boldsymbol{\xi}_j}(d\boldsymbol{\xi}). \tag{77}$$

Its first two moments define

$$\mathbf{A}_\nu = \int \boldsymbol{\xi}\,\nu_{t,x}(d\boldsymbol{\xi}),$$
$$\mathsf{D}_\nu = \int \boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\nu_{t,x}(d\boldsymbol{\xi}). \tag{78}$$

For a test function f, write the two backward generators as

$$\mathcal{L}_J f = \mathcal{L}_0 f + \int [\bar{f}(\mathbf{r},\mathbf{v}+\boldsymbol{\xi}) - f(x)]\,\nu_{t,x}(d\boldsymbol{\xi}),$$
$$\mathcal{L}_D f = \mathcal{L}_0 f + \mathbf{A}_\nu\cdot\nabla_{\mathbf{v}} f + \tfrac{1}{2}\mathsf{D}_\nu : \nabla^2_{\mathbf{v}} f. \tag{79}$$

The common operator $\mathcal{L}_0$ contains the smooth gravity, external force and unchanged species contributions. A bar denotes zero continuation into any declared killed state; both processes use the same physical stopping rules. The labels J and D refer to jump and diffusion dynamics and are distinct from the density-exposure symbol $J_{j,T}$.

This construction holds the first two encounter moments fixed while testing the effect of finite transitions. The reported unit-Coulomb-log kernels supply the weak-deflection reference and numerical checks; they are not automatically the moments of a specified finite-angle encounter domain. Those moments must be evaluated from Eq. (78). If a practical calculation resolves only an inner impact-parameter range, its diffusion coefficients integrate only the complementary range. A matching scale that depends on relative speed stays inside the encounter integrals. Varying that partition tests consistency without counting an encounter in both terms.

C. Finite-domain encounter calculations

We evaluate a common finite-angle encounter law in the six reference backgrounds, retaining their local anisotropic BH number distributions. These calculations specify both the transition sampler and its first two moments. They use a $60\,M_\odot$ probe and a frozen, independently sampled BH bath; lighter species are outside this particular collision calculation.

Let r_ρ be the radius at which the total BH mass density is half its central value. Define

$$L(r) = \min\left(r_\rho, \frac{\rho_{\rm BH}(r)}{|\partial_r\rho_{\rm BH}(r)|}\right), \qquad b_{\max} = \epsilon L(r), \tag{80}$$

with $L = r_\rho$ when the derivative vanishes and $\epsilon = 0.05, 0.1, 0.2$. Density-length prescriptions motivate the environmental scale [20, 21]; the finite central cap and these three fractions are declared controls, not calibrated cutoffs. We evaluate $r/r_\rho = 0, 1, 2$, zero probe speed, and radial and tangential speeds equal to the local number-weighted BH dispersion $\sigma_{\rm BH}$. Here $3\sigma^2_{\rm BH}$ is the number-weighted mean squared speed; this scalar control does not isotropize the distribution in the integrals.

To define an elastic Newtonian domain, set $M_j = m_{\rm r} + m_j$ and $\eta_j = m_{\rm r}m_j/M_j^2$. The leading parabolic radiation-loss estimate [22] gives the capture scale

$$r_{{\rm cap},j} = \frac{GM_j}{c^2}\left[\frac{85\pi}{6\sqrt{2}}\eta_j\left(\frac{c}{w}\right)^2\right]^{2/7},$$
$$r_{{\rm min},j} = \max\left(100\frac{GM_j}{c^2}, 10r_{{\rm cap},j}\right),$$
$$b^2_{{\rm min},j} = r^2_{{\rm min},j} + \frac{2GM_j r_{{\rm min},j}}{w^2}. \tag{81}$$

The factors 100 and 10 are explicit weak-field and radiative-exclusion conventions. Empty annuli contribute zero allowed flux. Incoming flux below this floor is recorded separately for a prospective common unresolved-event hazard; no capture outcome or trajectory stopping event is calculated here.

The impact-parameter and azimuth integrals can be performed analytically. At fixed $\mathbf{w}$, write $B = b_{90,j}$, $\zeta_j = m_j/M_j$, $\mathbf{n} = \mathbf{w}/w$, and

$$Q = \ln\frac{b^2_{\max}+B^2}{b^2_{\min}+B^2},$$
$$H = \frac{1}{b^2_{\min}+B^2} - \frac{1}{b^2_{\max}+B^2}. \tag{82}$$

Per unit incoming number density, the integrated rate and moments are

$$\gamma = \pi w(b^2_{\max} - b^2_{\min}),$$
$$\mathbf{a} = -2\pi\zeta_j w^2 B^2 Q\,\mathbf{n},$$
$$d_\parallel = 4\pi\zeta_j^2 w^3 B^4 H,$$
$$d_\perp = 2\pi\zeta_j^2 w^3 B^2(Q - B^2 H),$$
$$\mathsf{d} = d_\parallel\mathbf{n}\mathbf{n}^{\mathsf{T}} + d_\perp(\mathsf{I} - \mathbf{n}\mathbf{n}^{\mathsf{T}}). \tag{83}$$

Integrating these expressions against $F_j\,d^3u$ and summing over BH species gives Γ, $\mathbf{A}_\nu$ and D_ν. The zero-relative-speed limit is evaluated without division by w. The sampler draws field velocities with weights proportional to number density, relative speed and allowed annular area, then draws b^2 uniformly and a uniform impact azimuth. Its conditional first and second moments are therefore $\mathbf{A}_\nu/\Gamma$ and D_ν/Γ.

TABLE IX. Finite-angle BH encounters in the adopted reference hosts. At radius r_ρ, BH mass density is half its central value; this is not a half-mass radius. Each row uses $r = r_\rho$, radial remnant speed $v = \sigma_{\rm BH}$, remnant mass $60\,M_\odot$ and $b_{\rm max} = 0.1L$. The one-dimensional scale $\sigma_{\rm BH}$ is number weighted, and L is the smaller of r_ρ and the local BH density scale. $\Gamma_{\rm el}$ and $\langle|\Delta\mathbf{v}|^2\rangle_{\rm el}^{1/2}$ refer to the allowed elastic domain. f_{90} is the fraction of its diffusion trace arising from relative velocities with $b_{90} > L$. The latter is a scale diagnostic, not an error bound or a residence prediction.

Cluster	r_ρ (pc)	$\Gamma_{\rm el}$ (Myr^{-1})	$\langle\|\Delta\mathbf{v}\|^2\rangle_{\rm el}^{1/2}/\sigma_{\rm BH}$	f_{90} (%)
47 Tuc	0.1022	4.229	0.2201	0.02
NGC 1851	0.02245	1.784	0.7885	8.63
NGC 3201	0.2073	0.03278	0.8414	3.35
ω Cen	0.02478	170	0.4128	0.01
M4	0.03293	0.2063	0.8244	55.50
NGC 6397 (standard)	0.004386	1.161	0.4667	99.70




FIG. 9. Finite-domain BH encounter controls at $r = r_\rho$ and radial $v = \sigma_{\rm BH}$. The event rate and the fraction of diffusion trace contributed by $b_{90,j} > L$ are evaluated from the same allowed elastic law as the matched moments. Changing ϵ changes that law. The scale fraction diagnoses which deflection lengths contribute; it is not a residence error or an established validity threshold.

All 162 local settings meet the declared numerical criteria. Successive velocity-quadrature orders apply a 0.5% criterion to scaled drift and relative diffusion and rate refinements. The drift scale is $\max(|\mathbf{A}_v|, \mathrm{tr}\,\mathsf{D}_v/\sigma_{\rm BH})$; number-density normalization is checked to 0.2%. The separate scale-fraction diagnostic uses an absolute 0.025 refinement criterion. Its numerator is integrated over the exact relative-velocity ball $w < \sqrt{GM_j/L}$ for each species, avoiding an unresolved discontinuous mask. These are numerical checks, not physical calibration. At one specified state per host, 10^5 independent allowed increments reproduce the three first and six distinct raw-second moment components, with all discrepancies below 2.58 estimated standard errors across the six samples.

Table IX separates event frequency from the size of an individual velocity change. These are different physical ingredients of the residence hypothesis. A substantial per-event deflection can be infrequently sampled, while a larger rate can produce many smaller changes. Their consequences require the remnant's time-dependent sampling of the host. The cutoff controls and the recorded close-passage channel specify can-

didate matched comparisons; the calculation does not evolve a remnant or assign a residence-accuracy class.

D. Collisionless residence and path-integrated encounters

We next connect the local encounter law to timed motion in each of the six adopted potentials. The reference calculation suppresses collisions for every species while retaining their full smooth gravitational field. It measures the zero-collision occupation and evaluates encounter budgets along the resulting paths. The budgets do not feed back on the motion. This control isolates the orbital and occupation calculations needed before comparing collisional descriptions.

With the BH density radius defined above, set

$$v_{\rm dyn} = \sqrt{-r_\rho \psi'(r_\rho)}, \qquad t_{\rm dyn} = r_\rho / v_{\rm dyn},$$
$$r_{\rm i} = 0.5 r_\rho, \qquad v_{\rm launch} = \sqrt{2[\psi(r_{\rm i}) - \psi(2r_\rho)]}. \tag{84}$$

For each host, one launch is radially outward and one is tangential, with the same speed and energy. The radial apocenter is $2r_\rho$; the tangential turning points follow from its nonzero angular momentum. These are conditional total velocities, not inferred recoil velocities or a merger-selected population. The horizon is $T = 10 t_{\rm dyn}$, not ten orbital periods, and the region is $r < r_\rho$.

The Cartesian equations use $\ddot{\mathbf{r}} = \psi'(r)\mathbf{r}/r$ from the same potential as the launch energy. A path stops at the horizon or the first outward crossing of the smaller of the host support and its recorded minimum one-percent isolated-force domain. The latter is a validity boundary, not an escape surface or an accumulated orbit-error bound. All twelve reference paths reach the horizon inside their control domains. Every core visit contributes to the collisionless residence $C_T^{(0)}$; no remnant is brought to rest after returning.

Along each path, the actual velocity is rotated into the local radial/tangential frame to evaluate the anisotropic BH distribution. The finite-angle law uses $b_{\rm max} = 0.1 L(r)$ and Eq. (81). Define the dimensionless path budgets

$$\mathcal{H}_{\rm el} = \int_0^{T_*} \Gamma_{\rm el}(X_t^{(0)})\, dt, \qquad \mathcal{H}_{\rm exc} = \int_0^{T_*} \Gamma_{\rm exc}(X_t^{(0)})\, dt,$$
$$B_A = \frac{1}{v_{\rm launch}} \int_0^{T_*} |\mathbf{A}_\nu(X_t^{(0)})|\, dt,$$
$$B_D = \frac{1}{v_{\rm launch}^2} \int_0^{T_*} \mathrm{tr}\, \mathsf{D}_\nu(X_t^{(0)})\, dt. \tag{85}$$

Here $X_t^{(0)}$ is the prescribed collisionless path. These quantities describe its accumulated encounter intensity and moment scales. They are not realized encounter counts, capture probabilities, velocity changes, or corrections to residence. In particular, B_A discards vector cancellation and neither moment budget accounts for the altered path after an encounter. We also integrate $\rho_{\rm BH}(r(t))/\rho_{\rm BH}(0)$ and divide by the declared horizon to retain the same exposure convention in both launch directions.

An independent calculation uses the spherical radial equation,

$$\dot{r}^2 = 2[E + \psi(r)] - \ell^2/r^2, \tag{86}$$

with specific energy E and angular-momentum magnitude ℓ. Integrating between its independently determined turning points gives residence at the actual launch phase and finite horizon. A sine-squared radial substitution resolves the endpoint roots. This check does not use the Cartesian crossing times. The Cartesian calculation is separately refined in step size and tolerance, and energy and angular momentum are monitored. All twelve controls pass: the independent residence discrepancy divided by the horizon is below 8×10^{-10}, and the maximum energy drift divided by $v_{\rm dyn}^2$ is below 8×10^{-9}. These are numerical agreement measures for the adopted potential, not uncertainty estimates for the physical host.

The path integrals split at region crossings and radial turning points, including central passages. Their positive velocity quadrature is centered on relative velocity, with the field distribution evaluated at $\mathbf{u} = \mathbf{v} - \mathbf{w}$. The integration ball contains its full shifted support. This resolves the close-passage flux near $w = 0$ without altering the encounter law. Separate time and velocity refinement use a 0.5% criterion for the encounter budgets and local moments; normalized density-exposure refinement is below 10^{-5}. Allowed and excluded rates recover the parent incoming flux.

Table X and Fig. 10 show that equal energy does not fix finite-time residence. The ordering of radial and tangential occupation changes across these conditional settings; it depends on angular momentum and the phase reached at the finite horizon. This is not a universal directional preference or an encounter-induced effect. The residence fractions span 0.2625–0.3161, while $\mathcal{H}_{\rm el}$ spans approximately 0.0248–3.53. The common dimensionless horizon corresponds to 0.0208–1.67 Myr across hosts; the encounter budgets reflect both their local rates and these different physical durations. These controls supply verified crossing and occupation machinery and select the high-intensity setting examined in Sec. XI E.

E. Matched collisional residence in a conditional host

We now activate BH encounters in the ω Cen reference, selected before the collisional calculation because it has the largest allowed encounter integral among the collisionless controls. The remnant mass, launch radius, two equal-energy velocity directions, region and horizon remain those of Sec. XI D. All physical host species retain their smooth gravity. Collisions with lighter species are disabled in both arms, so this comparison isolates the reduction of the specified BH encounter law to its first two moments. The host and its BH reservoir do not respond to the remnant. These choices define a conditional transport experiment, not a prediction for an observed merger or a calibrated relaxation model for the whole cluster.

The adopted scales are $r_\rho = 0.02478\,\mathrm{pc}$, $v_{\rm dyn} = 11.675\,\mathrm{km\,s^{-1}}$ and $t_{\rm dyn} = 0.002076\,\mathrm{Myr}$. The launch speed is

TABLE X. Collisionless reference residence and BH encounter integrals. Each launch begins at $0.5r_\rho$ with the speed giving a radial apocenter of $2r_\rho$; the tangential launch has the same energy. Residence counts every visit to $r < r_\rho$ during $T = 10t_{\rm dyn}$. The gravitational field includes all physical host species. The allowed BH elastic rate uses $b_{\rm max} = 0.1L$ and the declared close-passage exclusion, evaluated along each prescribed path. Its time integral is a diagnostic that does not alter the orbit. These are conditional collisionless controls, not collisional residence predictions or merger probabilities.

		$C_T^{(0)}/T$		$\int_0^{T_*} \Gamma_{\rm el}\, dt$	
Cluster	T (Myr)	Radial	Tangential	Radial	Tangential
47 Tuc	0.19	0.3025	0.2783	0.998	0.834
NGC 1851	0.0818	0.2627	0.2753	0.142	0.137
NGC 3201	1.67	0.3161	0.2910	0.0728	0.0599
ω Cen	0.0208	0.2719	0.2887	3.53	3.44
M4	0.242	0.2625	0.2750	0.0488	0.0469
NGC 6397 (standard)	0.0224	0.2636	0.2765	0.0258	0.0248




FIG. 10. Collisionless residence and the allowed BH encounter rate integrated along the same reference paths. Launches have equal energy within each host but different angular momenta. The horizon retains their initial phases. The lower panel measures a path diagnostic; collisions are not applied to the trajectories.

19.884 km s^{-1}. We retain $b_{\rm max} = 0.1L$ and the same excluded close-passage channel as the local encounter calculation. Its flux defines a killing rate in both arms; the stop records an unresolved interaction, without assigning a capture outcome.

a. A common numerical encounter law. We tabulate the entire positive encounter-rate measure in radius, speed and pitch $p = 2\arctan(v_t/|v_r|)/\pi$, using the continuous endpoint conventions $p = 0$ for radial motion and $p = 1$ for tangential motion. At zero speed the pitch is set to zero; at the center the radial frame is aligned with the velocity. The coarse and refined tables have 550 and 3591 vertices, respectively, with every coarse interval bisected in all three coordinates. Non-negative multilinear weights $w_i(x)$ and the local frame trans-

formation R_x define

$$\nu_h(x, d\boldsymbol{\delta}) = \sum_i w_i(x)(R_x)_* \nu_i(d\boldsymbol{\delta}),$$
$$\Gamma_h(x) = \sum_i w_i(x)\Gamma_i,$$
$$\mathbf{A}_h(x) = R_x \sum_i w_i(x)\mathbf{A}_i,$$
$$\mathbf{D}_h(x) = R_x\left[\sum_i w_i(x)\mathbf{D}_i\right]R_x^{\mathsf{T}}. \tag{87}$$

Here $(R_x)_*$ transforms the increment measure. An event selects vertex i with probability $w_i\Gamma_i/\Gamma_h$, then samples that vertex's allowed flux. The sampled law and its moments have identical symmetry averages and quadrature. In particular, $\mathbf{D}_h$ is the raw second jump moment per unit time, not the covariance of a single conditional event. Thus interpolation preserves the matching algebraically; accuracy relative to the underlying local distribution remains a separate numerical question.

Every table vertex passes the declared velocity-quadrature and number-density checks. At 72 local comparisons, including symmetry limits, refining the table lowers the maximum relative diffusion-tensor discrepancy from 9.24% to 1.97%; the maximum scaled drift discrepancy falls from 4.57% to 0.963%. These finite checks are not uniform error bounds on the encounter law. A further six-state sampler check uses 40,000 independent increments per state on each grid. All sampled means and raw second moments lie within six estimated Monte Carlo standard errors of their matched values; the largest standardized discrepancy is 3.02. This checks implementation consistency, not the residence distribution.

b. Transport and stopping. The collision law is frozen at the start of each step. An exponential clock with the local excluded rate shortens the step to duration δ when necessary. Two velocity-Verlet gravity half-flows surround a velocity update: a Gaussian increment with mean $\mathbf{A}_h\delta$ and covariance $\mathbf{D}_h\delta$, or the sum of $N \sim \mathrm{Poisson}(\Gamma_h\delta)$ independent finite increments from ν_h/Γ_h. The two unrestricted frozen updates have exactly the same mean and covariance. Multiple events are permitted, and their summed increment is applied at the midpoint. Individual event times within that split update are not resolved. Consequently equality of substep moments is not a claim of identical evolving paths or of time-step convergence.

Occupation inside $r < r_\rho$ is integrated exactly along each straight drift segment of this scheme; density exposure uses its midpoint density. A spatial table exit is located on the drift chord. Speed-domain exits are checked at the resolved gravity and collision endpoints, with their resolution assessed by step refinement. The numerical domain is $r < 4r_\rho$, $|\mathbf{v}| < 4v_{\mathrm{dyn}}$; its spatial edge lies well inside the inherited isolated-force domain. Neither table boundary is a physical escape surface. The first resolved validity stop or excluded event ends accumulation, and every path retains the full $T = 10t_{\mathrm{dyn}}$ denominator.

There are 4096 independent paths per arm, direction, grid and time step, giving 65,536 trajectories across 16 settings. The steps are $0.04t_{\mathrm{dyn}}$ and $0.02t_{\mathrm{dyn}}$. Separate random streams drive Gaussian increments, excluded clocks, Poisson counts and finite events. No paired-sample reduction of uncertainty is assumed. Null-collision controls reproduce the previously calculated residence and exposure within 1.30×10^{-5} after normalization, with smaller discrepancies at the finer step. An independent all-species density integral checks the gravitational acceleration to relative discrepancy below 5.46×10^{-4}. Analytical tests check anisotropic compound-Poisson and Gaussian moments, multiple events, exponential killing and competing stops.

c. Interpretation and numerical precision. For a numerical-domain stop at τ, omitted normalized residence is bounded pathwise by $(T-\tau)/T$. We retain this bound and the probability of a numerical stop, separately from the excluded physical channel. Mean contrasts use independent-ensemble standard errors and descriptive 95% normal intervals. Stop probabilities use approximate Wilson intervals, including nonzero upper limits for zero observed stops. The CDF-supremum discrepancy uses a union-DKW envelope: its half-width is the sum of the two single-ensemble bounds, here 0.0463. These intervals are not simultaneous across every comparison in the pilot.

The registered mean-residence target is 0.01, with a 0.005 numerical sensitivity budget; the distribution target is a CDF-supremum discrepancy of 0.05. A separate 0.002 budget concerns missing residence after numerical-domain stops. Grid and time-step contrasts carry their own sampling uncertainty. Finite refinement tests do not supply a rigorous continuum-error bound, and a sampled interval containing zero does not establish numerical convergence. The pilot tests the encounter endpoint and identifies the precision needed for the occupation-response calculation; it does not evaluate that response predictor.

d. Measured endpoint differences. Table XI and Fig. 11 compare mean occupation and residence distributions. On the refined grid at $\Delta t = 0.02t_{\mathrm{dyn}}$, mean residence fractions are 0.3040 (jump) and 0.3116 (diffusion) for the radial launch, and 0.3243 and 0.3292 for the tangential launch. Both arms exceed their collisionless controls, 0.2719 and 0.2887, respectively. This is the combined effect of the specified collision law; the experiment does not separately attribute it to drift and random deflections.

The signed jump-minus-diffusion contrasts are -0.00757 and -0.00499, with descriptive 95% intervals $[-0.01126, -0.00389]$ and $[-0.00952, -0.00045]$. The radial interval spans the 0.01 absolute-difference target; the tangential interval lies below that target at this numerical setting. In contrast, the residence-CDF discrepancies are 0.1265 and 0.1055, with marginal union-DKW intervals $[0.0802, 0.1727]$ and $[0.0592, 0.1517]$. The residence-distribution discrepancy exceeds the declared 0.05 target at this resolution even though the mean shifts are smaller. All eight numerical comparisons show empirical CDF distances between 0.0969 and 0.1265; no simultaneous confidence claim over these comparisons is made.

The refined-step mean BH exposures are 0.4084 versus 0.4132 for radial motion and 0.4146 versus 0.4166 for tangential motion. The signed exposure contrasts have inter-

TABLE XI. Conditional matched encounter pilot in the ω Cen reference. Each entry compares independent ensembles of 4096 paths per arm over $T = 10t_{\rm dyn}$. The jump and diffusion collision increments have the same local mean and covariance. Only BH collisions are active; all physical species contribute smooth gravity. Means include stopped paths and use the full horizon. The signed contrast $\delta_C = (\langle C_T\rangle_J - \langle C_T\rangle_D)/T$ has a descriptive 95% normal interval; the CDF discrepancy has a 95% union-DKW interval. These are conditional numerical results, not calibrated cluster predictions.

Launch	Grid	$\Delta t/t_{\rm dyn}$	$\langle C_T/T\rangle_J$	$\langle C_T/T\rangle_D$	δ_C [95% interval]	Δ_F [95% interval]
Radial	Coarse	0.04	0.3045	0.3111	-0.0066 $[-0.0102, -0.0029]$	0.097 [0.051, 0.143]
Radial	Coarse	0.02	0.3064	0.3110	-0.0046 $[-0.0085, -0.0008]$	0.112 [0.066, 0.158]
Radial	Fine	0.04	0.3040	0.3164	-0.0124 $[-0.0162, -0.0086]$	0.117 [0.071, 0.164]
Radial	Fine	0.02	0.3040	0.3116	-0.0076 $[-0.0113, -0.0039]$	0.126 [0.080, 0.173]
Tangential	Coarse	0.04	0.3229	0.3319	-0.0091 $[-0.0137, -0.0044]$	0.121 [0.075, 0.168]
Tangential	Coarse	0.02	0.3210	0.3280	-0.0069 $[-0.0114, -0.0025]$	0.117 [0.071, 0.163]
Tangential	Fine	0.04	0.3246	0.3292	-0.0045 $[-0.0091, +0.0001]$	0.098 [0.052, 0.145]
Tangential	Fine	0.02	0.3243	0.3292	-0.0050 $[-0.0095, -0.0005]$	0.105 [0.059, 0.152]

vals $[-0.00735, -0.00225]$ and $[-0.00483, 0.00083]$, respectively. Thus the tangential exposure contrast is unresolved by its sampling interval even though its residence CDF differs. A descriptive event-count diagnostic finds 88 radial and 130 tangential jump paths without any resolved event or stop. They retain the collisionless numerical residence, contributing 2.15% and 3.17% of their ensembles. This discrete component illustrates information discarded by a continuous approximation, but its size alone cannot account for the full observed CDF discrepancy.

The refinement evidence prevents a continuum accuracy verdict. Some absolute mean-refinement intervals extend above the 0.005 budget, reaching 0.00912. At the refined step no radial numerical stop is observed; the tangential diffusion arm has one spatial stop and its jump counterpart has none. The tangential empirical missing-residence bound is 5.39×10^{-5}. Nevertheless, summing the two marginal Wilson upper endpoints gives 0.00232, exceeding the separate 0.002 missing-residence budget; that sum is not a joint 95% bound. Across the full pilot, ten spatial stops and one excluded-channel stop are retained, with no speed-domain stops. The result therefore establishes a difference between these matched numerical residence distributions, while further numerical precision and physical calibration are required to transfer that statement to the continuous host model or to a real cluster. The first-response test below uses independently generated predictions and validation trajectories.

F. From local increments to expected residence

For a nonnegative occupation weight $q(t, x)$, define the expected remaining accumulation in the diffusion process,

$$U_D(t, x) = \mathbb{E}_{D,t,x}\left[\int_t^{T\wedge\tau_D} q(s, X_s)\, ds\right]. \tag{88}$$

The residence weight is $q = \mathbf{1}_{C(t)}$; a species exposure uses $q = \rho_j$. With zero terminal and killed-state payoff, the backward equation is

$$(\partial_t + \mathcal{L}_D)U_D = -q. \tag{89}$$

Thus local encounters affect residence through the dependence of U_D on the remnant's velocity as well as its position. A velocity change can alter future crossings even when its instantaneous position is unchanged.

Let $O_q = \int_0^{T_*} q(s, X_s)\, ds$. For the same initial law in both processes, stopped generator calculus gives

$$\mathbb{E}_J O_q - \mathbb{E}_D O_q = \mathbb{E}_J \int_0^{T\wedge\tau_J} \mathcal{R}_D(s, X_s)\, ds, \tag{90}$$

$$\begin{aligned}\mathcal{R}_D &= (\mathcal{L}_J - \mathcal{L}_D)U_D, \\ &= \int \Big[\bar{U}_D(\mathbf{r}, \mathbf{v} + \boldsymbol{\xi}) - U_D(x) - \boldsymbol{\xi}\cdot\nabla_{\mathbf{v}} U_D \\ &\qquad - \tfrac{1}{2}(\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}) : \nabla^2_{\mathbf{v}} U_D\Big]\, \nu_{t,x}(d\boldsymbol{\xi}).\end{aligned} \tag{91}$$

Time arguments on U_D in the last expression are suppressed. To obtain Eq. (90), apply the stopped Dynkin identity [23] to U_D along the jump process, substitute Eq. (89), and use the zero endpoint payoff. This is an application of standard generator calculus, supplying a theoretical relation rather than a numerical result.

The identity requires integrability and a common generator domain, or an appropriate extended-domain formulation. It can first be applied to a smooth occupation weight; the sharp-region limit and absorbing-boundary regularity require separate justification. The full zero-continued residual must be retained for jumps that cross a killed boundary. A global Taylor bound is not assumed there. When a finite uniform residual bound exists for residence, it supplies the sufficient mean-error bound

$$\frac{|\mathbb{E}_J C_T - \mathbb{E}_D C_T|}{T} \le \sup_{t,x} |\mathcal{R}_D(t, x)|. \tag{92}$$

A path-weighted estimate can be sharper than this supremum. Neither expression fixes the sign of the residence difference, and cancellation in the mean does not establish agreement of the full distribution.

Equations (90)–(91) identify the physical mechanism to examine: finite velocity changes matter where the expected remaining occupation is not well represented by its quadratic velocity expansion. The central diffusion fraction sets a local




FIG. 11. Conditional residence under the matched numerical encounter laws. Left: mean residence with descriptive 95% sampling intervals, compared with the collisionless value (dashed). C and F denote the coarse and refined encounter tables. Right: cumulative residence distributions for the refined table and $\Delta t = 0.02 t_{\rm dyn}$. Stopped paths remain in both distributions. A close mean does not require similar occupation distributions.

second-moment contribution; the residual adds the encounter distribution and the future orbital response needed for an accuracy statement. No residual or residence-error bound has yet been evaluated for the adopted hosts.

G. A prediction that can be tested independently

The residual identity is an analytical relation. A further, falsifiable proposal is that a response estimate computed using the diffusion process predicts the sign and size of the residence change in a specified regime. Define an auxiliary family

$$\mathcal{L}_a = (1-a)\mathcal{L}_D + a\mathcal{L}_J, \qquad 0 \le a \le 1. \tag{93}$$

This is a diagnostic interpolation of generators, not a change to the adopted cluster population. Every member retains the same first two encounter moments. Under the regularity conditions for differentiating the stopped expectation, its first response is

$$\frac{\partial}{\partial a}\mathbb{E}_a C_T\Big|_{a=0} = \mathbb{E}_D \int_0^{T\wedge\tau_D} \mathcal{R}_D(s, X_s)\, ds,$$
$$\widehat{\delta}_C = \frac{1}{T}\mathbb{E}_D \int_0^{T\wedge\tau_D} \mathcal{R}_D(s, X_s)\, ds. \tag{94}$$

The signed quantity $\widehat{\delta}_C$ is a proposed predictor for the finite change at $a = 1$. Replacing the jump-process expectation in Eq. (90) by a diffusion-process expectation is not exact at that endpoint. Comparing the prediction with encounter ensembles at increasing a tests where the first response remains useful and where repeated finite transitions require the full process. A zero-BH realization gives a null control when only BH encounters are varied.

The hypothesis can fail if the first response misses the sign, magnitude or prescribed accuracy class of the full residence change. An endpoint remainder resolved beyond the declared tolerance, after controlling sampling and numerical errors, establishes a nonlinear response over this interpolation even

though the first two local moments match. An imprecise estimate does not establish that failure. Prediction accuracy and diffusion accuracy are separate tests: a response estimate can correctly predict a large residence change for which diffusion is inadequate. Establishing diffusion adequacy requires the inferred total jump–diffusion discrepancy, with its uncertainty, to satisfy the residence tolerance. Neither predictive accuracy nor such adequacy has been established in the current calculations.

A numerical route to this first response uses a single transition replacement. For N time steps, let P_D and P_J be the corresponding diffusion and encounter transition kernels, including their integrated occupation rewards, and define $P_a = (1-a)P_D + aP_J$. Choose a step K uniformly. Follow diffusion except for one encounter update at K, then resume diffusion. If $O^{(K)}$ and O^D are the two accumulated observables, differentiation of the finite product gives

$$\frac{d}{da}\mathbb{E}_a O\Big|_{a=0} = N\,\mathbb{E}\Big[O^{(K)} - O^D\Big]. \tag{95}$$

The relation includes any change in the immediate reward. Independent random streams isolate each pair, with shared continuation noise within the pair. Exact finite-state enumeration and stochastic checks verify this discrete construction. Connecting it to Eq. (94) requires time-step convergence, and prediction of the full encounter endpoint remains a separate test. The following calculation evaluates this discrete response; its continuous-time interpretation requires additional convergence evidence.

H. A controlled test of the residence response

The test asks whether the first response predicts the finite change in normalized residence, rather than whether the matched moments reproduce every trajectory statistic. Write $Y = C_T/T$, $\mu_h(a) = \mathbb{E}_{a,h}Y$ and $S_h = \mu_h'(0)$ for the complete numerical transition family. The subscript h denotes both the encounter table and time step. Equation (95) gives an unbiased estimator of S_h for this finite product. It does not identify S_h with the continuous-time response without convergence, nor imply $\mu_h(1) - \mu_h(0) = S_h$.

a. Conditional replacement and coupling. A uniformly selected scheduled step K is retained even if the diffusion prefix has already stopped. Such paths contribute zero; discarding them would condition on survival. At an active insertion, condition on the shared excluded-channel clock and let δ be the resulting step duration. The frozen allowed-event mean and probabilities are

$$\lambda = \Gamma_h(X_K)\delta, \qquad p_0 = e^{-\lambda}, \qquad p_+ = 1 - p_0. \tag{96}$$

The encounter transition is decomposed into a zero-event branch and a branch with a Poisson count conditioned to be positive. The latter includes multiple encounters; it is not a forced single collision. Two diffusion branches use increments $\mathbf{A}_h\delta \pm \sqrt{\delta}\,F_hZ$, where $F_hF_h^{\mathsf{T}} = \mathbf{D}_h$, Z is standard normal and the drift is unchanged in both. All four branches include the same full gravity, reward and stopping operations, then resume diffusion. Subsequent normal and exponential variates are shared by aligned path rows; after separation each branch evaluates its own state-dependent hazard. This coupling preserves each marginal law without imposing equal stopping times.

Let Y_0, Y_+, Y_{D+}, Y_{D-} be the four full-horizon rewards. The weighted score is

$$\widehat{S}_i = N\left[p_0Y_0 + p_+Y_+ - \frac{Y_{D+} + Y_{D-}}{2}\right], \qquad \mathbb{E}\widehat{S}_i = S_h. \tag{97}$$

Conditioning the Poisson update and pairing opposite normal increments changes sampling variance, not the derivative being estimated. Its efficiency must be measured against the direct replacement-pair estimator. Constant-rate impulse controls permit an independent polynomial-moment calculation: a sixth-degree terminal reward has a nonzero quadratic remainder in a, distinguishing a correctly estimated derivative from an exact endpoint prediction.

b. Independent prediction and validation. The host, remnant mass, encounter domain, region, horizon and stopping rules remain those of Sec. XI E. In addition to the radial and tangential launches, a velocity at 45° to the outward radial direction has the same initial position and speed. The first two endpoint settings were previously examined; the oblique direction supplies a previously unused launch, not an independently calibrated host. The primary setting uses the refined table and $\Delta t = 0.01t_{\mathrm{dyn}}$. The coarse and refined tables at $0.02t_{\mathrm{dyn}}$ supply separate grid and time sensitivity comparisons.

An independent pilot of 1024 samples per estimator and launch compares variance and cost. Only its weighted-score variance selects the primary prediction count: the smallest power of two meeting a nominal simultaneous-interval half-width of 0.0005, between 8192 and 65,536 samples. The cap is a resource limit, not a precision guarantee. Control settings use 8192 samples. Prediction samples are separate from the pilot, and all predictions are recorded before validation trajectories are generated.

Validation uses 8192 independent pairs at each setting and interpolation value, with a diffusion control paired to each mixture path. At the primary resolution $a = 0.05, 0.1, 0.25, 0.5, 1$; control resolutions use $a = 1$. An independent Bernoulli choice selects the entire encounter or diffusion transition at every scheduled step. Here a is a diagnostic weight between transition laws; the host population and physical encounter parameters remain fixed. Define the prediction residual

$$R_h(a) = \mu_h(a) - \mu_h(0) - aS_h. \tag{98}$$

Separate prediction and validation samples permit their variance contributions to be added. Normal-approximation intervals use a predeclared Bonferroni family size of 64, covering 63 response, validation and refinement statements. Rare influential scores remain visible through their extrema and contribution diagnostics; these intervals are not exact finite-sample coverage guarantees. At the two smallest values, $R_h(a)/a$ assesses departure from the first-order response at finite a. An

interval containing zero alone does not demonstrate equivalence or establish the derivative limit.

At $a = 1$, an error interval wholly inside $[-0.001, 0.001]$ supports the declared finite-resolution prediction accuracy; an interval wholly outside establishes disagreement, and overlap is inconclusive. The separate numerical-sensitivity budget is 0.0005. Numerical-domain termination is also retained: with $B_b = (T - \tau_b)/T$ for a numerical stop in branch b, and zero otherwise, the per-score continuation allowance is

$$B_{S,i} = N\left[p_0 B_0 + p_+ B_+ + \frac{B_{D+} + B_{D-}}{2}\right]. \tag{99}$$

The missing-residence budget is 0.0002. A finite refinement contrast is not a rigorous continuum-error bound, and zero observed stops do not establish zero stopping probability. Passing a sampling criterion therefore does not by itself establish continuum or physical-cluster accuracy. Signed score contributions by insertion time and radius, normalized by all samples, decompose the estimated response. When that estimate is unresolved, the bin sums do not identify a physical region of dominant sensitivity without their own uncertainty analysis. They do not predict a residence distribution, capture rate or merger probability.

c. Pilot precision. The weighted-score sample variances are 132.6, 1890.9 and 1762.0 for the radial, tangential and oblique launches, respectively. The corresponding direct-pair variances are 259.2, 2392.9 and 2127.3. The measured variance reductions are therefore modest: factors of 1.96, 1.27 and 1.21. They do not establish a reduction in computational cost, which also includes constructing the local encounter samplers. The variance-only selection rule reaches the 65,536-sample cap for all three primary predictions. Consequently, the requested precision is not guaranteed by the prespecified sample allocation; it must be assessed from the independent production samples. These pilot scores are excluded from every production response estimate.

d. Local coupling diagnostic. An exploratory deterministic check, specified after examining the pilot, distinguishes the marginal diffusion law from the coupling used to estimate its response. For $D = V\Lambda V^{\mathsf{T}}$, both $F(D) = V\sqrt{\Lambda}$ and the symmetric root $G(D) = V\sqrt{\Lambda}V^{\mathsf{T}}$ reproduce covariance D. However, common standard-normal noise Z gives

$$\mathbb{E}\left\| \sqrt{\Delta t}\,[F(D_1) - F(D_0)]Z\right\|^2 = \Delta t\,\|F(D_1) - F(D_0)\|_F^2. \tag{100}$$

Eigenvector signs and basis changes can therefore weaken the coupling even when $D_1 - D_0$ is small. For example, the retained factors of $D_\pm = \mathrm{diag}(1 \pm \epsilon, 1 \mp \epsilon, 2)$ differ in Frobenius norm by 2 as $\epsilon \to 0$, whereas their symmetric roots approach each other. In the radial launch state, changing the dimensionless transverse velocity by 10^{-8} changes the local tensor by only 4.13×10^{-10} in relative Frobenius norm. The noise-distance squared, divided by $\Delta t\ \mathrm{tr}\, D_0$, is nevertheless 1.26 for the retained eigenbasis factors, compared with 4.33×10^{-20} for the symmetric roots. Both choices retain the correct marginal covariance. These local checks do not attribute the measured residence-score variance to this mechanism, establish a better trajectory estimator, or alter the prespecified runs. They identify the coupling choice evaluated in Sec. XI I.

e. Independent validation and scope of the conclusion. All nine prediction batches were completed and hash-registered before the 21 independent validation batches. Together with the six excluded pilot batches, the calculation retains 423,936 independent pairs or quartets, comprising 1,345,536 branch trajectories. The three primary predictions each use 65,536 weighted samples. Their simultaneous-interval half-widths are 0.1564, 0.5726, 0.5490, in radial, tangential and oblique order, compared with the 0.0005 target. The sample cap therefore does not resolve the requested response precision.

Table XII and Fig. 12 compare these predictions with the independently measured shifts. All three primary endpoint residual intervals overlap the declared tolerance boundary, yielding inconclusive accuracy classifications. This is a limitation of the inference at the achieved precision; it establishes neither predictive agreement nor a resolved nonlinear response. All six small-a scaled-residual intervals include zero. These finite-a intervals do not establish an accurate derivative limit.

None of the twelve response and endpoint refinement contrasts has its full interval within the 0.0005 sensitivity budget. The primary finite-sample continuation allowances are 0.1134, 0.0504, 0.0450. These are conservative bounds on possible missing residence in the retained samples, not measurements of an omitted physical effect or population confidence bounds. The weighted response can amplify rare numerical-domain stops through its factor of N. Sampling, refinement and continuation therefore remain separate obstacles to a continuum accuracy classification. The prespecified time and radius bins supply descriptive score decompositions; the unresolved response does not permit a physical interpretation of their apparent peaks.

I. Precision of the residence-response estimator

The unresolved prediction motivates a controlled precision comparison at the same fine-table setting. Its objective is to reduce the sampling variance of S_h, keeping the conditional host, encounter law, complete transition, occupation reward and stopping conventions fixed. The three methods use the weighted quartet of Eq. (97): eigenbasis coupling with uniform insertion (E), symmetric-root coupling with uniform insertion (S), and symmetric-root coupling with proportional insertion-time stratification (T). Their marginal transition laws and response target are identical; their joint branch couplings and sampling designs differ.

a. Mathematical controls. The symmetric positive-semidefinite root $G(D)$ removes dependence on arbitrary eigenvector signs and choices within a degenerate eigenspace. Both $F(D)$ and $G(D)$ map a standard normal vector to the required covariance, but shared-noise discrepancies depend on the chosen factors. Matrix-factor and Gaussian-transport results distinguish covariance preservation from optimal coupling [24]. The symmetric root is continuous on the positive-semidefinite cone; a uniform Lipschitz bound cannot be assumed near zero eigenvalues. Its local regularity does

TABLE XII. First-response prediction and independent endpoint validation in the conditional ω Cen reference, using the refined encounter table and $\Delta t = 0.01 t_{\rm dyn}$. All quantities concern normalized residence C_T/T. Each response uses 65,536 independent weighted samples; each endpoint shift uses 8192 mixture/diffusion pairs independent of the prediction. Intervals are normal approximations with Bonferroni family size 64. The endpoint error is the measured shift minus S_h; its classification uses the declared 0.001 absolute tolerance. These sampling intervals exclude numerical refinement and continuation uncertainty, so a finite-resolution classification is not a continuum or physical-cluster verdict.

Launch	S_h [interval]	Endpoint shift [interval]	Prediction error [interval]	Test
Radial	−0.0368 [−0.1932, +0.1196]	−0.0078 [−0.0124, −0.0032]	+0.0290 [−0.1275, +0.1855]	Inconclusive
Tangential	−0.1421 [−0.7147, +0.4305]	−0.0067 [−0.0122, −0.0012]	+0.1354 [−0.4373, +0.7081]	Inconclusive
Oblique	−0.0617 [−0.6107, +0.4874]	−0.0043 [−0.0092, +0.0006]	+0.0574 [−0.4917, +0.6064]	Inconclusive

not prove optimal coupling of an entire stopped residence history. Numerical controls verify marginal drift and covariance, degenerate-eigenspace invariance, stopped rewards and exact recovery of the original algorithm in method E.

Time stratification has a separate variance guarantee. Partition the $N = 1000$ scheduled insertion steps into four equal intervals, with probabilities $w_s = 1/4$. Let m_s and v_s denote the conditional mean and variance of the symmetric-factor score within interval s. Independent insertion draws give

$$S_h = \sum_s w_s m_s, \qquad \widehat{S}_{\rm T} = \sum_s w_s \overline{S}_s,$$
$$\mathrm{Var}(\widehat{S}_{\rm T}) = \sum_s \frac{w_s^2 v_s}{n_s}. \tag{101}$$

With proportional allocation $n_s = n w_s$, the law of total variance gives

$$\mathrm{Var}(\widehat{S}_{\rm S}) - \mathrm{Var}(\widehat{S}_{\rm T}) = \frac{1}{n}\sum_s w_s (m_s - S_h)^2 \geq 0. \tag{102}$$

This identity compares S and T under the same score law. It does not guarantee a substantial reduction, lower cost, or a corresponding comparison with E. Sample variance estimates can fluctuate around this population relation. Paths stopped before insertion retain zero scores and their assigned strata; neither tail trimming nor survival conditioning is used.

b. Fixed comparison and reporting. The design fixes four independent batches of 1024 quartets for each method and each of the radial, tangential and oblique launches: 36,864 quartets and 147,456 correlated branch trajectories in total. The methods have independent random streams. Each T batch contains 256 draws per interval; uncertainty is computed from the within-interval variances in Eq. (101). The complete protocol and implementation were recorded before execution. Launch and method order rotate between batches. Sample counts and allocations are not adapted to outcomes.

The primary efficiency ratios compare mean-estimator variances E/S and S/T at equal sample count. Values greater than one favor the denominator. Independent within-design percentile bootstraps use 8192 replicates, with nominal 95% Bonferroni coverage across six ratios. These intervals are exploratory approximations and cannot account for unseen rare scores. Means and independent-arm differences have descriptive standard errors, without an equivalence claim. A secondary ratio compares variance multiplied by recorded process CPU time, conditional on the machine and cache state. Bootstrap intervals for that ratio hold timing fixed and exclude timing uncertainty; equal quartet counts do not imply equal computing budgets.

c. Measured precision and remaining uncertainty. Table XIII and Fig. 13 show E/S variance ratios of 112.5, 1133.6 and 1885.3 for radial, tangential and oblique launches. All three exploratory intervals lie above one. The twelve individual batch comparisons also favor S in point estimate, with ratios ranging from 45.4 to 3256.8; this diagnostic is not an additional confidence test. At equal sample count, the standard errors decrease by factors of 10.6, 33.7 and 43.4. The corresponding recorded CPU-adjusted ratios are 110.0, 1183.6 and 1928.8. The improvement is therefore substantial under both reported metrics in this conditional calculation. It supports the relevance of covariance-factor continuity to the estimator, without asserting optimality or a universal efficiency factor.

For S/T, the estimated variance ratios are 0.876, 1.099 and 0.978, and every exploratory interval includes one. No additional gain from the four fixed time strata is resolved. Estimates below one do not contradict Eq. (102): independently estimated variances fluctuate, and the guaranteed population reduction can be small. The largest one percent of absolute scores contributes 33–41% of the total absolute score across the six symmetric-factor ensembles. Influential tails therefore remain important even after the large coupling improvement. A further descriptive check, specified after the pilot, decomposes each estimated variance into its centered-square contributions. One radial S score supplies 61.1% of that sample variance; the largest 41 contributions supply 92.9%, 89.6% and 84.0% for the three S launches. These are finite-sample concentration diagnostics, not estimates of unobserved tail probabilities.

The symmetric/uniform response estimates are −0.02160, 0.00540 and −0.01163, with descriptive standard errors 0.01850, 0.01914 and 0.01304. These errors still exceed the earlier 0.0005 simultaneous-half-width target before any multiplicity adjustment. The associated finite-sample numerical-continuation allowances are zero, 3.85×10^{-5} and zero; zeros in a finite sample do not bound the population contribu-




FIG. 12. Independent mixture validation of the first-response prediction in the conditional ω Cen reference. Top: measured normalized-residence shifts and the prediction aS_h at the primary resolution. Bottom: their residuals and the declared 0.001 error band. Error bars and prediction bands use normal approximations with family size 64. Broad prediction intervals make the endpoint test inconclusive; they do not demonstrate agreement. Numerical refinement and continuation allowances are assessed separately.

tion. Refinement and continuation have not been resolved by the coupling comparison. The evidence supports S as the reference estimator for a subsequent independently sampled response test. Earlier endpoint validation is not reused to certify the selected estimator's prediction accuracy. The response hypothesis remains open, while this calculation removes a demonstrated source of avoidable sampling variance from its numerical test.

J. Separating finite transitions from population response

The first comparison assumes that successive encounters sample a supplied local distribution. A subsequent finite-population comparison resolves individual BH positions, velocities and reciprocal forces, testing repeated encounters, correlations and exchange of energy with the reservoir. A noninteger expected BH count does not define such a realization. An explicit integer-count and mass-sampling law must reproduce the declared ensemble moments and any imposed joint constraints; pair correlations and binaries must be specified separately.

In that calculation the smooth BH contribution to gravity is replaced by the resolved BH forces, and its collision term is removed from the diffusion operator. The remnant acts back on the resolved objects. Retaining the original full smooth BH field or its full diffusion term in addition would double-count their contribution. The lighter species can initially retain their common prescribed background; relaxing that choice defines a further comparison. A local jump process with independently resampled encounters and a mutually interacting finite population answer different questions, so their results remain distinguishable.

The diagnostic variables include $m_{\rm r}/m_j$, the distribution of $b_{90,j}/b_{{\rm max},j}$, the radial extent of the BH component, and the remnant mass relative to the reservoir encountered within the chosen region. These are tested alongside the encounter cor-

TABLE XIII. Precision comparison at the same fine-table setting and $\Delta t = 0.01 t_{\rm dyn}$. Each method uses 4096 independent response quartets per launch, divided into four execution batches. E denotes eigenbasis coupling with uniform insertion, S symmetric-root coupling with uniform insertion, and T symmetric-root coupling with proportional time stratification. Standard errors concern the estimated response slope. Variance ratios above one favor the denominator method. Bracketed intervals are exploratory percentile bootstrap intervals adjusted over six ratios; they do not cover unseen rare scores, numerical bias or physical-model uncertainty.

Launch	SE(E)	SE(S)	SE(T)	Variance E/S [interval]	Variance S/T [interval]
Radial	0.19622	0.01850	0.01976	112.50 [33.35, 517.46]	0.88 [0.13, 6.38]
Tangential	0.64449	0.01914	0.01826	1.13×10^3 [639.71, 2.37×10^3]	1.10 [0.41, 3.02]
Oblique	0.56597	0.01304	0.01318	1.89×10^3 [1.07×10^3, 3.86×10^3]	0.98 [0.41, 2.52]




FIG. 13. Efficiency of response sampling in the conditional ω Cen reference. E/S compares eigenbasis and symmetric-root coupling under uniform insertion; S/T compares uniform and proportionally stratified insertion under symmetric-root coupling. Ratios above one favor the denominator. Panel (a) uses variance at equal quartet count; panel (b) multiplies it by recorded CPU time. The six exploratory variance-ratio intervals use within-design resampling; the right panel rescales these intervals by the observed cost ratios. Timing uncertainty, unseen rare scores, numerical convergence and physical calibration are not covered.

relation time and the time on which the local background changes. An averaging interval must contain enough small, decorrelated increments while resolving changes in the orbit and background. The fixed $60\,M_\odot$ probe used here does not hold these ratios equal across hosts.

K. Matched launches, tides and measurable outcomes

Within each adopted host, both encounter descriptions use the same merger-conditioned launch distribution, remnant mass, physical horizon and region. The initial study can condition on specified launch states and then average them under a declared joint launch law. Positions, progenitor velocities and recoil directions retain their correlations. Central radial launches are one control, not the full ensemble.

For comparisons across hosts, use

$$\widetilde{C} = \frac{C_T}{t_0}, \qquad \widetilde{J}_j = \frac{J_{j,T} r_h^3}{M t_0}. \tag{103}$$

Matching $\mathbf{r}_{\rm m}/r_h$, velocities in units of v_0, T/t_0 and the scaled region separates host structure from the imposed geometric and temporal scales. Remnant-to-host and remnant-to-scatterer mass ratios are explicit controls; the present common-mass calculations do not match all of them. The

comparison within one host isolates encounter representation before cross-host differences are interpreted.

The Galactic reference orbit and differential force are also held fixed between the two descriptions. An isolated control is restricted to its declared omitted-force domain; a calculation with the differential Galactic force uses a separate physical escape convention. The fixed-host response indicator already calculated identifies phases for a later frozen-versus-evolving-host comparison. That comparison changes the host treatment while preserving launch and encounter choices, thereby separating tidal response from the approximation of encounters.

When finite two-body encounters are resolved, the corresponding τ_{strong} stop in Eq. (33) is removed in both matched descriptions. Otherwise the comparison would terminate before the effect under study. Interactions still outside the supplied physics use the same declared event or killing rule in both descriptions. The residence reported after such a stop is the accumulated modeled residence; a complete-horizon continuation is not inferred.

Two complementary residence discrepancies are

$$\Delta_C = \frac{|\mathbb{E}_J C_T - \mathbb{E}_D C_T|}{T},$$
$$\Delta_F = \sup_{0\le c\le 1} \left| \Pr_J(C_T/T \le c) - \Pr_D(C_T/T \le c) \right|. \tag{104}$$

Exposure distributions, entry-time distributions and the probabilities of each termination cause accompany these quantities. Censored paths remain in the accounting; conditioning on survivors would define a different comparison. The backward residual constrains a mean, while Δ_F tests changes in distribution shape and tails.

Numerical, ensemble and smoothing errors must be reduced below predeclared tolerances for the compared observables. An uncertainty interval for a nonnegative discrepancy lying entirely below its tolerance supports adequacy in that setting; one lying entirely above establishes a material difference. An interval spanning the tolerance is inconclusive. Coverage accounts for the launch settings and observables examined. A resolved discrepancy is then attributed through separate finite-transition, finite-population and host-evolution comparisons. A result remains informative whether it identifies a required additional dynamical variable or a regime in which the simpler description preserves residence. Section XI E declares tolerances, encounter cutoffs, diagnostic launches and independent ensembles for one conditional setting. A comparison across the six hosts and a merger-conditioned population require their own declared inputs. The pilot intervals are marginal; a simultaneous accuracy claim requires coverage accounting for the comparisons used.

XII. DISCUSSION

The research objective is to connect encounter physics to the accuracy of finite-time remnant residence. The adopted cluster models supply the potentials, mass spectra and local phase-space distributions for this problem. Their numerical reconstructions and scattering integrals establish a consistent starting point across six environments. The source audit supplies unit conventions, tracer definitions and complete joint parameter rows; these remain the conditioning of the transport calculation.

The completed results show why the calculation needs both spatial and velocity information. A substantial central BH contribution can be concentrated in a small region or persist farther through the host. The remnant's excursions determine how those regions are sampled. Population size, local second moments and spatial concentration therefore enter different parts of the physical question. The comparison is organized around these quantities rather than assigning a single accuracy criterion from a cluster label or a global BH count.

The matched generators isolate the reduction of finite encounter increments to their first two moments. The backward occupation equation then relates that reduction to the expected remaining residence or exposure. This is the analytical bridge between the calculated local coefficients and the trajectory observables. Its nonlocal residual provides a concrete quantity to evaluate, together with direct comparisons of means, distributions and termination probabilities. The identity does not itself establish the size of the residual in a physical cluster.

Finite-population response and host evolution are separate extensions of this comparison. Resolving reciprocal encounters tests memory and reservoir response; retaining time-dependent Galactic forcing and evolving the host tests the background through which the remnant moves. Keeping these comparisons distinct makes an observed difference interpretable and identifies which added physics is required for a specified accuracy.

The current evidence consists of host reconstructions, local kernels, joint-row sensitivity, environmental diagnostics, finite-angle encounter calculations, collisionless controls, a conditional matched collisional pilot and an independently sampled discrete response test. The encounter law establishes event frequency and deflection amplitude separately, with a shared sampler and moment law. The control paths show how angular momentum and launch phase affect finite-time residence at fixed energy, and connect local encounter rates to the environments actually sampled. The single-transition replacement now supplies a measured discrete response with predictions recorded before independent validation. Its achieved precision is insufficient to assess the declared endpoint accuracy, so the physical response hypothesis remains open. The exact response identity and the direct numerical endpoint differences retain their separate evidential roles. The fixed covariance-coupling comparison substantially reduces response-estimator variance at similar measured cost, while four-interval time stratification provides no resolved additional benefit. This supplies a better reference estimator for an independent response test; it does not establish the response hypothesis or eliminate influential tails. Resolving the response, its numerical refinement and its boundary continuation is necessary before testing an observable-specific accuracy criterion across these adopted environments. Timed trajectories from a validated calculation could subsequently support companion-access physics with an additional target population and interaction law; density exposure by itself remains

distinct from an encounter count or merger probability.

Appendix A: Conservative Galactic-force bound

All lengths in this derivation share one unit. Let $s = |\mathbf{R}|$, $\varpi = \sqrt{X^2 + Y^2}$, and ℓ be the radius of the spatial ball used for the force bound. Define

$$\begin{aligned} s_- &= s - \ell, \qquad \varpi_- = \max(\varpi - \ell, 0), \\ \zeta_- &= \sqrt{\max(|Z| - \ell, 0)^2 + b_d^2}, \\ d_- &= \sqrt{\varpi_-^2 + (a_d + \zeta_-)^2}. \end{aligned} \tag{A1}$$

For Irrgang Model I, write the potential amplitudes as $C_b = GM_b$, $C_d = GM_d$, and $C_h = GM_h$, with scale lengths b_b, a_d, b_d, a_h and halo cutoff Λ. If $s_- > 0$ and $s + \ell < \Lambda$, sufficient component bounds are

$$\begin{aligned} B_b &= \frac{2C_b}{(s_-^2 + b_b^2)^{3/2}}, \\ B_d &= \frac{C_d}{d_-^3} \max\left(2, 1 + \frac{a_d b_d^2}{\zeta_-^3}\right), \\ B_h &= \frac{C_h}{a_h s_-(a_h + s_-)}, \qquad B = B_b + B_d + B_h. \end{aligned} \tag{A2}$$

The bulge and $\gamma = 2$ halo expressions bound their radial and tangential Hessian eigenvalues. For the disk, let $\zeta = \sqrt{Z^2 + b_d^2}$, $d^2 = X^2 + Y^2 + (a_d + \zeta)^2$, and

$$\begin{aligned} \mathbf{q} &= (X, Y, (a_d + \zeta)Z/\zeta), \\ \mathsf{J} &= \mathrm{diag}(1, 1, 1 + a_d b_d^2/\zeta^3). \end{aligned} \tag{A3}$$

Its Hessian is $C_d(\mathsf{J}/d^3 - 3\mathbf{q}\mathbf{q}^\mathsf{T}/d^5)$. Since $\mathsf{J} \succeq \mathsf{I}$ and $|\mathbf{q}| \le d$, its eigenvalues lie between $-2C_d/d^3$ and $C_d J_{zz}/d^3$. The conservative spatial minima in Eq. (A1) then imply B_d. Summing operator norms bounds the full tensor, so Eq. (55) applies to the exact force difference throughout the ball, not just its linear tidal approximation.

[1] M. N. Rosenbluth, W. M. MacDonald, and D. L. Judd, Fokker-planck equation for an inverse-square force, Phys. Rev. **107**, 1 (1957).

[2] K. Tep, J.-B. Fouvry, and C. Pichon, Non-resonant relaxation of anisotropic globular clusters (2022), arXiv:2201.03985 [astro-ph.GA].

[3] J. J. Webb, N. W. C. Leigh, A. Singh, K. E. S. Ford, B. McKernan, and J. Bellovary, The evolution of kicked stellar-mass black holes in star cluster environments (2018), arXiv:1711.09100 [astro-ph.GA].

[4] B. Bar-Or, G. Kupi, and T. Alexander, Stellar energy relaxation around a massive black hole, Astrophys. J. **764**, 52 (2013).

[5] M. Gieles and A. Zocchi, A family of lowered isothermal models, Mon. Not. R. Astron. Soc. **454**, 576 (2015), arXiv:1508.02120 [astro-ph.IM].

[6] N. Dickson, V. Hénault-Brunet, H. Baumgardt, M. Gieles, and P. Smith, Multimass modelling of Milky Way globular clusters—I. implications on their stellar initial mass function above 1 $m_\odot$ (2023), arXiv:2303.01637 [astro-ph.GA].

[7] N. Dickson, P. J. Smith, V. Hénault-Brunet, M. Gieles, and H. Baumgardt, Multimass modelling of Milky Way globular clusters—II. present-day black hole populations (2024), arXiv:2308.13037 [astro-ph.GA].

[8] M. Libralato, A. Bellini, E. Vesperini, *et al.*, The Hubble Space Telescope UV legacy survey of galactic globular clusters. XXIII. proper-motion catalogs and internal kinematics (2022), arXiv:2206.09924 [astro-ph.GA].

[9] M. Häberle, N. Neumayer, C. Clontz, *et al.*, oMEGACat. VI. analysis of the overall kinematics of Omega Centauri in 3D: velocity dispersion, kinematic distance, anisotropy, and energy equipartition, Astrophys. J. **983**, 95 (2025), arXiv:2503.04903.

[10] E. Vasiliev and H. Baumgardt, Gaia EDR3 view on galactic globular clusters, Mon. Not. R. Astron. Soc. **505**, 5978 (2021), arXiv:2102.09568.

[11] H. Baumgardt and M. Hilker, A catalogue of masses, structural parameters and velocity dispersion profiles of 112 Milky Way globular clusters, Mon. Not. R. Astron. Soc. **478**, 1520 (2018), arXiv:1804.08359.

[12] H. Baumgardt and E. Vasiliev, Accurate distances to galactic globular clusters through a combination of Gaia EDR3, HST, and literature data, Mon. Not. R. Astron. Soc. **505**, 5957 (2021), arXiv:2105.09526.

[13] I. King, The structure of star clusters. I. an empirical density law, Astron. J. **67**, 471 (1962).

[14] F. Renaud, M. Gieles, and C. M. Boily, Evolution of star clusters in arbitrary tidal fields, Mon. Not. R. Astron. Soc. **418**, 759 (2011).

[15] T. Fukushige and D. C. Heggie, The time-scale of escape from star clusters, Mon. Not. R. Astron. Soc. **318**, 753 (2000).

[16] Z. Wan, W. H. Oliver, H. Baumgardt, G. F. Lewis, M. Gieles, V. Hénault-Brunet, T. de Boer, E. Balbinot, G. S. Da Costa, and D. Mackey, The dynamics of the globular cluster NGC 3201 out to the Jacobi radius, Mon. Not. R. Astron. Soc. **502**, 4513 (2021).

[17] J. J. Webb, W. E. Harris, A. Sills, and J. R. Hurley, The influence of orbital eccentricity on tidal radii of star clusters, Astrophys. J. **764**, 124 (2013).

[18] A. Irrgang, B. Wilcox, E. Tucker, and L. Schiefelbein, Milky Way mass models for orbit calculations, Astron. Astrophys. **549**, A137 (2013).

[19] G. Bertin and A. L. Varri, The construction of nonspherical models of quasi-relaxed stellar systems, Astrophys. J. **689**, 1005 (2008).

[20] A. Just, F. M. Khan, P. Berczik, A. Ernst, and R. Spurzem, Dynamical friction of massive objects in galactic centres, Mon. Not. R. Astron. Soc. **411**, 653 (2011).

[21] J. A. Petts, A. Gualandris, and J. I. Read, A semi-analytic dynamical friction model that reproduces core stalling (2015),

arXiv:1509.07871.
[22] R. M. O'Leary, B. Kocsis, and A. Loeb, Gravitational waves from scattering of stellar-mass black holes in galactic nuclei (2009), arXiv:0807.2638.
[23] E. B. Dynkin, Infinitesimal operators of Markov processes, Theory Probab. Appl. **1**, 34 (1956).
[24] R. Bhatia, T. Jain, and Y. Lim, On the Bures-Wasserstein distance between positive definite matrices (2017), arXiv:1712.01504 [math.FA].