

## 1. Lepton Flavor Violation

Table 1: HFAG Winter 2012 upper limit for the lepton flavor violating  $\tau$  decay modes. L means lepton number violation as well as the lepton flavor violation.

Decay mode	Category	Upper limit at 90%		Exp.	Ref.
$\Gamma_{156} = e^- \gamma$	$l\gamma$	$< 12.$	$\cdot 10^{-8}$	Belle	[1]
		$< 3.3$	$\cdot 10^{-8}$	Babar	[2]
$\Gamma_{157} = \mu^- \gamma$		$< 4.5$	$\cdot 10^{-8}$	Belle	[1]
		$< 4.4$	$\cdot 10^{-8}$	Babar	[2]
$\Gamma_{158} = e^- \pi^0$	$lP^0$	$< 2.2$	$\cdot 10^{-8}$	Belle	[3]
		$< 13.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{159} = \mu^- \pi^0$		$< 2.7$	$\cdot 10^{-8}$	Belle	[3]
		$< 11.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{162} = e^- \eta$		$< 4.4$	$\cdot 10^{-8}$	Belle	[3]
		$< 16.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{163} = \mu^- \eta$		$< 2.3$	$\cdot 10^{-8}$	Belle	[3]
		$< 15.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{172} = e^- \eta'(958)$		$< 3.6$	$\cdot 10^{-8}$	Belle	[3]
		$< 24.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{173} = \mu^- \eta'(958)$		$< 3.8$	$\cdot 10^{-8}$	Belle	[3]
		$< 14.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{160} = e^- K_S^0$		$< 2.6$	$\cdot 10^{-8}$	Belle	[5]
		$< 3.3$	$\cdot 10^{-8}$	Babar	[6]
$\Gamma_{161} = \mu^- K_S^0$		$< 2.3$	$\cdot 10^{-8}$	Belle	[5]
		$< 4.0$	$\cdot 10^{-8}$	Babar	[6]
$\Gamma_{174} = e^- f_0(958)$	$lS^0$	$< 3.2$	$\cdot 10^{-8}$	Belle	[7]
$\Gamma_{175} = \mu^- f_0(958)$		$< 3.4$	$\cdot 10^{-8}$	Belle	[7]
$\Gamma_{164} = e^- \rho^0$	$lV^0$	$< 1.8$	$\cdot 10^{-8}$	Belle	[8]
		$< 4.6$	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{165} = \mu^- \rho^0$		$< 1.2$	$\cdot 10^{-8}$	Belle	[8]
		$< 2.6$	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{168} = e^- K^*(892)^0$		$< 3.2$	$\cdot 10^{-8}$	Belle	[8]
		$< 5.9$	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{169} = \mu^- K^*(892)^0$		$< 7.2$	$\cdot 10^{-8}$	Belle	[8]
		$< 17.$	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{170} = e^- \bar{K}^*(892)^0$		$< 3.4$	$\cdot 10^{-8}$	Belle	[8]
		$< 4.6$	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{171} = \mu^- \bar{K}^*(892)^0$		$< 7.0$	$\cdot 10^{-8}$	Belle	[8]
		$< 7.3$	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{176} = e^- \phi$		$< 3.1$	$\cdot 10^{-8}$	Belle	[8]
		$< 3.1$	$\cdot 10^{-8}$	Babar	[9]

$\Gamma_{177} = \mu^- \phi$		$< 8.4$	$\cdot$	$10^{-8}$	Belle	[8]
		$< 19.$	$\cdot$	$10^{-8}$	Babar	[9]
$\Gamma_{166} = e^- \omega$		$< 4.8$	$\cdot$	$10^{-8}$	Belle	[8]
		$< 11.$	$\cdot$	$10^{-8}$	Babar	[10]
$\Gamma_{167} = \mu^- \omega$		$< 4.7$	$\cdot$	$10^{-8}$	Belle	[8]
		$< 10.$	$\cdot$	$10^{-8}$	Babar	[10]
<hr/>						
$\Gamma_{178} = e^- e^+ e^-$	$lll$	$< 2.7$	$\cdot$	$10^{-8}$	Belle	[11]
		$< 2.9$	$\cdot$	$10^{-8}$	Babar	[12]
$\Gamma_{181} = \mu^- e^+ e^-$		$< 1.8$	$\cdot$	$10^{-8}$	Belle	[11]
		$< 2.2$	$\cdot$	$10^{-8}$	Babar	[12]
$\Gamma_{179} = e^- \mu + \mu^-$		$< 2.7$	$\cdot$	$10^{-8}$	Belle	[11]
		$< 3.2$	$\cdot$	$10^{-8}$	Babar	[12]
$\Gamma_{183} = \mu^- \mu + \mu^-$		$< 2.1$	$\cdot$	$10^{-8}$	Belle	[11]
		$< 3.3$	$\cdot$	$10^{-8}$	Babar	[12]
$\Gamma_{182} = e^- \mu + e^-$		$< 1.5$	$\cdot$	$10^{-8}$	Belle	[11]
		$< 1.8$	$\cdot$	$10^{-8}$	Babar	[12]
$\Gamma_{180} = \mu^- e^+ \mu^-$		$< 1.7$	$\cdot$	$10^{-8}$	Belle	[11]
		$< 2.6$	$\cdot$	$10^{-8}$	Babar	[12]
<hr/>						
$\Gamma_{184} = e^- \pi^+ \pi^-$	$lhh$	$< 2.3$	$\cdot$	$10^{-8}$	Belle	[13]
		$< 12.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{186} = \mu^- \pi^+ \pi^-$		$< 2.1$	$\cdot$	$10^{-8}$	Belle	[13]
		$< 29.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{188} = e^- \pi^+ K^-$		$< 3.7$	$\cdot$	$10^{-8}$	Belle	[13]
		$< 32.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{194} = \mu^- \pi^+ K^-$		$< 8.6$	$\cdot$	$10^{-8}$	Belle	[13]
		$< 26.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{189} = e^- K^+ \pi^-$		$< 3.1$	$\cdot$	$10^{-8}$	Belle	[13]
		$< 17.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{195} = \mu^- K^+ \pi^-$		$< 4.5$	$\cdot$	$10^{-8}$	Belle	[13]
		$< 32.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{192} = e^- K^+ K^-$		$< 3.4$	$\cdot$	$10^{-8}$	Belle	[13]
		$< 14.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{198} = \mu^- K^+ K^-$		$< 4.4$	$\cdot$	$10^{-8}$	Belle	[13]
		$< 25.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{191} = e^- K_S^0 K_S^0$		$< 7.1$	$\cdot$	$10^{-8}$	Belle	[5]
$\Gamma_{197} = \mu^- K_S^0 K_S^0$		$< 8.0$	$\cdot$	$10^{-8}$	Belle	[5]
$\Gamma_{185} = e^+ \pi^- \pi^-$	(L)	$< 2.0$	$\cdot$	$10^{-8}$	Belle	[13]
	(L)	$< 27.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{187} = \mu^+ \pi^- \pi^-$	(L)	$< 3.9$	$\cdot$	$10^{-8}$	Belle	[13]
	(L)	$< 7.0$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{190} = e^+ \pi^- K^-$	(L)	$< 3.2$	$\cdot$	$10^{-8}$	Belle	[13]
	(L)	$< 18.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{196} = \mu^+ \pi^- K^-$	(L)	$< 4.8$	$\cdot$	$10^{-8}$	Belle	[13]

$\Gamma_{193} = e^+ K^- K^-$	(L)	$< 22.$	$\cdot$	$10^{-8}$	Babar	[14]
	(L)	$< 3.3$	$\cdot$	$10^{-8}$	Belle	[13]
$\Gamma_{199} = \mu^+ K^- K^-$	(L)	$< 15.$	$\cdot$	$10^{-8}$	Babar	[14]
	(L)	$< 4.7$	$\cdot$	$10^{-8}$	Belle	[13]
	(L)	$< 48.$	$\cdot$	$10^{-8}$	Babar	[14]
$\Gamma_{211} = \pi^- \Lambda$	$\Lambda h$	$< 3.0$	$\cdot$	$10^{-8}$	Belle	[15]
		$< 5.8$	$\cdot$	$10^{-8}$	Babar	[16]
$\Gamma_{212} = \pi^- \bar{\Lambda}$		$< 2.8$	$\cdot$	$10^{-8}$	Belle	[15]
		$< 5.9$	$\cdot$	$10^{-8}$	Babar	[16]
$\Gamma_{xx} = K^- \Lambda$		$< 4.2$	$\cdot$	$10^{-8}$	Belle	[15]
		$< 15.$	$\cdot$	$10^{-8}$	Babar	[16]
$\Gamma_{xx} = K^- \bar{\Lambda}$		$< 3.1$	$\cdot$	$10^{-8}$	Belle	[15]
		$< 7.2$	$\cdot$	$10^{-8}$	Babar	[16]

## References

- [1] K. Hayasaka, et al., Phys.Lett. B666 (2008) 16–22.
- [2] B. Aubert, et al., Phys.Rev.Lett. 104 (2010) 021802.
- [3] K. Hayasaka, J.Phys.:Conf.Ser. 335 (2011) 012029. See <http://iopscience.iop.org/1742-6596/335/1/012029/>.
- [4] B. Aubert, et al., Phys.Rev.Lett. 98 (2007) 061803.
- [5] Y. Miyazaki, Phys.Lett. B692 (2010) 4–9.
- [6] B. Aubert, et al., Phys.Rev. D79 (2009) 012004.
- [7] Y. Miyazaki, et al., Phys. Lett. B672 (2009) 317–322.
- [8] Y. Miyazaki, Phys.Lett. B699 (2011) 251–257.
- [9] B. Aubert, et al., Phys.Rev.Lett. 103 (2009) 021801.
- [10] B. Aubert, et al., Phys.Rev.Lett. 100 (2008) 071802.
- [11] K. Hayasaka, K. Inami, Y. Miyazaki, et al., Phys. Lett. B687 (2010) 139–143.
- [12] J. Lees, et al., Phys.Rev. D81 (2010) 111101.
- [13] Y. Miyazaki, Search for Lepton-Flavor-Violating and Lepton-Number-Violating  $\tau^- \rightarrow l h h'$  Decays Mode, 2011. A talk presented at International Workshop on  $e^+e^-$  collisions from phi to psi (PHIPSI11), Novosibirsk, 19-22 Sep 2011, <http://phipsi11.inp.nsk.su/talks/phipsi11.hayasaka.pdf>.
- [14] B. Aubert, et al., Phys.Rev.Lett. 95 (2005) 191801.
- [15] K. Hayasaka, Recent LFV results on tau lepton from Belle, 2011. A talk presented at International Workshop on  $e^+e^-$  collisions from phi to psi (PHIPSI11), Novosibirsk, 19-22 Sep 2011. <http://phipsi11.inp.nsk.su/talks/phipsi11.hayasaka.pdf>.
- [16] G. Lafferty, Nucl.Phys.Proc.Suppl. 169 (2007) 186–191.