

1. Lepton Flavor Violation

Table 1: HFAG Winter 2012 upper limit for the lepton flavor violating τ decay modes. L means lepton number violation as well as the lepton flavor violation.

Decay mode	Category	Upper limit at 90%		Exp.	Ref.
$\Gamma_{156} = e^- \gamma$	$l\gamma$	$< 12.$	$\cdot 10^{-8}$	Belle	[1]
		< 3.3	$\cdot 10^{-8}$	Babar	[2]
$\Gamma_{157} = \mu^- \gamma$		< 4.5	$\cdot 10^{-8}$	Belle	[1]
		< 4.4	$\cdot 10^{-8}$	Babar	[2]
$\Gamma_{158} = e^- \pi^0$	lP^0	< 2.2	$\cdot 10^{-8}$	Belle	[3]
		$< 13.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{159} = \mu^- \pi^0$		< 2.7	$\cdot 10^{-8}$	Belle	[3]
		$< 11.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{162} = e^- \eta$		< 4.4	$\cdot 10^{-8}$	Belle	[3]
		$< 16.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{163} = \mu^- \eta$		< 2.3	$\cdot 10^{-8}$	Belle	[3]
		$< 15.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{172} = e^- \eta'(958)$		< 3.6	$\cdot 10^{-8}$	Belle	[3]
		$< 24.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{173} = \mu^- \eta'(958)$		< 3.8	$\cdot 10^{-8}$	Belle	[3]
		$< 14.$	$\cdot 10^{-8}$	Babar	[4]
$\Gamma_{160} = e^- K_S^0$		< 2.6	$\cdot 10^{-8}$	Belle	[5]
		< 3.3	$\cdot 10^{-8}$	Babar	[6]
$\Gamma_{161} = \mu^- K_S^0$		< 2.3	$\cdot 10^{-8}$	Belle	[5]
		< 4.0	$\cdot 10^{-8}$	Babar	[6]
$\Gamma_{174} = e^- f_0(958)$	lS^0	< 3.2	$\cdot 10^{-8}$	Belle	[7]
$\Gamma_{175} = \mu^- f_0(958)$		< 3.4	$\cdot 10^{-8}$	Belle	[7]
$\Gamma_{164} = e^- \rho^0$	lV^0	< 1.8	$\cdot 10^{-8}$	Belle	[8]
		< 4.6	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{165} = \mu^- \rho^0$		< 1.2	$\cdot 10^{-8}$	Belle	[8]
		< 2.6	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{168} = e^- K^*(892)^0$		< 3.2	$\cdot 10^{-8}$	Belle	[8]
		< 5.9	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{169} = \mu^- K^*(892)^0$		< 7.2	$\cdot 10^{-8}$	Belle	[8]
		$< 17.$	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{170} = e^- \bar{K}^*(892)^0$		< 3.4	$\cdot 10^{-8}$	Belle	[8]
		< 4.6	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{171} = \mu^- \bar{K}^*(892)^0$		< 7.0	$\cdot 10^{-8}$	Belle	[8]
		< 7.3	$\cdot 10^{-8}$	Babar	[9]
$\Gamma_{176} = e^- \phi$		< 3.1	$\cdot 10^{-8}$	Belle	[8]
		< 3.1	$\cdot 10^{-8}$	Babar	[9]

$\Gamma_{177} = \mu^- \phi$		< 8.4	\cdot	10^{-8}	Belle	[8]
		$< 19.$	\cdot	10^{-8}	Babar	[9]
$\Gamma_{166} = e^- \omega$		< 4.8	\cdot	10^{-8}	Belle	[8]
		$< 11.$	\cdot	10^{-8}	Babar	[10]
$\Gamma_{167} = \mu^- \omega$		< 4.7	\cdot	10^{-8}	Belle	[8]
		$< 10.$	\cdot	10^{-8}	Babar	[10]
<hr/>						
$\Gamma_{178} = e^- e^+ e^-$	lll	< 2.7	\cdot	10^{-8}	Belle	[11]
		< 2.9	\cdot	10^{-8}	Babar	[12]
$\Gamma_{181} = \mu^- e^+ e^-$		< 1.8	\cdot	10^{-8}	Belle	[11]
		< 2.2	\cdot	10^{-8}	Babar	[12]
$\Gamma_{179} = e^- \mu + \mu^-$		< 2.7	\cdot	10^{-8}	Belle	[11]
		< 3.2	\cdot	10^{-8}	Babar	[12]
$\Gamma_{183} = \mu^- \mu + \mu^-$		< 2.1	\cdot	10^{-8}	Belle	[11]
		< 3.3	\cdot	10^{-8}	Babar	[12]
$\Gamma_{182} = e^- \mu + e^-$		< 1.5	\cdot	10^{-8}	Belle	[11]
		< 1.8	\cdot	10^{-8}	Babar	[12]
$\Gamma_{180} = \mu^- e^+ \mu^-$		< 1.7	\cdot	10^{-8}	Belle	[11]
		< 2.6	\cdot	10^{-8}	Babar	[12]
<hr/>						
$\Gamma_{184} = e^- \pi^+ \pi^-$	lhh	< 2.3	\cdot	10^{-8}	Belle	[13]
		$< 12.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{186} = \mu^- \pi^+ \pi^-$		< 2.1	\cdot	10^{-8}	Belle	[13]
		$< 29.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{188} = e^- \pi^+ K^-$		< 3.7	\cdot	10^{-8}	Belle	[13]
		$< 32.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{194} = \mu^- \pi^+ K^-$		< 8.6	\cdot	10^{-8}	Belle	[13]
		$< 26.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{189} = e^- K^+ \pi^-$		< 3.1	\cdot	10^{-8}	Belle	[13]
		$< 17.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{195} = \mu^- K^+ \pi^-$		< 4.5	\cdot	10^{-8}	Belle	[13]
		$< 32.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{192} = e^- K^+ K^-$		< 3.4	\cdot	10^{-8}	Belle	[13]
		$< 14.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{198} = \mu^- K^+ K^-$		< 4.4	\cdot	10^{-8}	Belle	[13]
		$< 25.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{191} = e^- K_S^0 K_S^0$		< 7.1	\cdot	10^{-8}	Belle	[5]
$\Gamma_{197} = \mu^- K_S^0 K_S^0$		< 8.0	\cdot	10^{-8}	Belle	[5]
$\Gamma_{185} = e^+ \pi^- \pi^-$	(L)	< 2.0	\cdot	10^{-8}	Belle	[13]
	(L)	$< 27.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{187} = \mu^+ \pi^- \pi^-$	(L)	< 3.9	\cdot	10^{-8}	Belle	[13]
	(L)	< 7.0	\cdot	10^{-8}	Babar	[14]
$\Gamma_{190} = e^+ \pi^- K^-$	(L)	< 3.2	\cdot	10^{-8}	Belle	[13]
	(L)	$< 18.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{196} = \mu^+ \pi^- K^-$	(L)	< 4.8	\cdot	10^{-8}	Belle	[13]

$\Gamma_{193} = e^+ K^- K^-$	(L)	$< 22.$	\cdot	10^{-8}	Babar	[14]
	(L)	< 3.3	\cdot	10^{-8}	Belle	[13]
$\Gamma_{199} = \mu^+ K^- K^-$	(L)	$< 15.$	\cdot	10^{-8}	Babar	[14]
	(L)	< 4.7	\cdot	10^{-8}	Belle	[13]
	(L)	$< 48.$	\cdot	10^{-8}	Babar	[14]
$\Gamma_{211} = \pi^- \Lambda$	Λh	< 3.0	\cdot	10^{-8}	Belle	[15]
		< 5.8	\cdot	10^{-8}	Babar	[16]
$\Gamma_{212} = \pi^- \bar{\Lambda}$		< 2.8	\cdot	10^{-8}	Belle	[15]
		< 5.9	\cdot	10^{-8}	Babar	[16]
$\Gamma_{xx} = K^- \Lambda$		< 4.2	\cdot	10^{-8}	Belle	[15]
		$< 15.$	\cdot	10^{-8}	Babar	[16]
$\Gamma_{xx} = K^- \bar{\Lambda}$		< 3.1	\cdot	10^{-8}	Belle	[15]
		< 7.2	\cdot	10^{-8}	Babar	[16]

References

- [1] K. Hayasaka, et al., Phys.Lett. B666 (2008) 16–22.
- [2] B. Aubert, et al., Phys.Rev.Lett. 104 (2010) 021802.
- [3] K. Inami, Updated Search for Lepton-Flavor-Violating τ^- Decays into $\ell^- \pi^0$, $\ell^- \eta$ and $\ell^- \eta'$, 2011. A talk presented at the 11th International Workshop on Tau-Lepton Physics (TAU2010), Manchester, 13-17 Sep 2010, http://belle.kek.jp/belle/talks/tau/13.Sep/tau2010_inami-5.pdf.
- [4] B. Aubert, et al., Phys.Rev.Lett. 98 (2007) 061803.
- [5] Y. Miyazaki, Phys.Lett. B692 (2010) 4–9.
- [6] B. Aubert, et al., Phys.Rev. D79 (2009) 012004.
- [7] Y. Miyazaki, et al., Phys. Lett. B672 (2009) 317–322.
- [8] Y. Miyazaki, Phys.Lett. B699 (2011) 251–257.
- [9] B. Aubert, et al., Phys.Rev.Lett. 103 (2009) 021801.
- [10] B. Aubert, et al., Phys.Rev.Lett. 100 (2008) 071802.
- [11] K. Hayasaka, K. Inami, Y. Miyazaki, et al., Phys. Lett. B687 (2010) 139–143.
- [12] J. Lees, et al., Phys.Rev. D81 (2010) 111101.
- [13] Y. Miyazaki, Search for Lepton-Flavor-Violating and Lepton-Number-Violating $\tau^- \rightarrow l h h'$ Decays Mode, 2011. A talk presented at International Workshop on $e^+ e^-$ collisions from phi to psi (PHIPSI11), Novosibirsk, 19-22 Sep 2011, http://phipsi11.inp.nsk.su/talks/phipsi11_hayasaka.pdf.
- [14] B. Aubert, et al., Phys.Rev.Lett. 95 (2005) 191801.
- [15] K. Hayasaka, Recent LFV results on tau lepton from Belle, 2011. A talk presented at International Workshop on $e^+ e^-$ collisions from phi to psi (PHIPSI11), Novosibirsk, 19-22 Sep 2011. http://phipsi11.inp.nsk.su/talks/phipsi11_hayasaka.pdf.
- [16] G. Lafferty, Nucl.Phys.Proc.Suppl. 169 (2007) 186–191.