## Optimization Homework 1

Hector G. Flores Rodriguez (Id. 25721714)

October 11, 2017

## 1. Problem 1

**1-(b)** The Fibonnacci method was used to optimized the functions:

$$f_1(x) = x^2 + 2x$$
  
 $f_2(x) = x^3 - 13x^2 + 36$ 

Prior to optmizing the provided functions, an initial interval for candidate starting points was determined by visually inspecting the graphs of the functions (shown in Figure 1).




Figure 1: Quadratic and non-quadratic functions for optmization.

The estimated value for  $x_1^*$  lies near -1, whereas the estimated value for  $x_2^*$  lies near 8. Hence, we can sample from a normal distribution for both functions:

$$X_1 \sim \mathcal{N}(\mu_1 = -1, \, \sigma_1 = 1)$$
  
 $X_2 \sim \mathcal{N}(\mu_2 = 8, \, \sigma_2 = 1)$ 

Table 1: Results for  $f_1(x) = x^2 + 2x$ 

Table 1. Results for $f_1(x) = x + 2x$					
	$x^*$	$f_1(x^*)$	function_calls	clock_time	
	-1.003858	-0.999985	13.0	0.000258	
	-0.999863	-1.000000	12.0	0.000121	
	-0.458484	-0.706761	12.0	0.000142	
	-0.133349	-0.248917	12.0	0.000113	
	-0.995906	-0.999983	12.0	0.000107	
	-1.794326	-0.369046	12.0	0.000126	
	-0.998712	-0.999998	12.0	0.000121	
	-0.675915	-0.894969	12.0	0.000112	
	-0.849852	-0.977456	12.0	0.000097	
	-0.998391	-0.999997	12.0	0.000108	
	-1.006449	-0.999958	12.0	0.000099	
	-1.009986	-0.999900	12.0	0.000097	
	-0.998913	-0.999999	12.0	0.000097	
	-1.293503	-0.913856	12.0	0.000130	
	-0.989732	-0.999895	12.0	0.000165	
	-0.493534	-0.743493	12.0	0.000159	
	-1.455641	-0.792392	12.0	0.000164	
	-0.779196	-0.951246	12.0	0.000162	
	-1.010637	-0.999887	12.0	0.000120	
	-0.995907	-0.999983	12.0	0.000100	
Avg:	-0.947108	-0.879886	12.050	0.000158	
Std:	0.351413	0.216419	0.224	0.000056	

Table 2: Results for  $f_2(x) = x^3 - 13x^2 + 36$ 

$x^*$	$f_2(x^*)$	function_calls	clock_time	
7.699171	314.553840	13.0	0.000316	
7.630426	305.599131	12.0	0.000154	
7.940157	347.457533	12.0	0.000132	
7.030768	235.247719	12.0	0.000120	
7.778227	325.086647	12.0	0.000136	
7.676198	311.540361	12.0	0.000131	
7.471320	285.591838	12.0	0.000118	
9.357814	592.745461	12.0	0.000137	
7.949493	348.780494	12.0	0.000145	
7.497876	288.862239	12.0	0.000126	
7.477308	286.326859	12.0	0.000160	
6.089015	150.528652	12.0	0.000166	
8.458776	426.580208	12.0	0.000126	
7.625984	305.026966	12.0	0.000138	
8.655849	459.757240	12.0	0.000249	
7.073400	239.804382	12.0	0.000280	
7.573978	298.386878	12.0	0.000229	
7.160760	249.349189	12.0	0.000226	
7.293491	264.392311	12.0	0.000221	
7.571402	298.060772	12.0	0.000213	
7.650571	316.683936	12.050	0.000176	
0.664623	92.297258	0.224	0.000059	
	x* 7.699171 7.630426 7.940157 7.030768 7.778227 7.676198 7.471320 9.357814 7.949493 7.497876 7.477308 6.089015 8.458776 7.625984 8.655849 7.073400 7.573978 7.160760 7.293491 7.571402 7.650571	$x^*$ $f_2(x^*)$ 7.699171314.5538407.630426305.5991317.940157347.4575337.030768235.2477197.778227325.0866477.676198311.5403617.471320285.5918389.357814592.7454617.949493348.7804947.497876288.8622397.477308286.3268596.089015150.5286528.458776426.5802087.625984305.0269668.655849459.7572407.073400239.8043827.573978298.3868787.160760249.3491897.293491264.3923117.571402298.0607727.650571316.683936	$x^*$ $f_2(x^*)$ function_calls7.699171314.55384013.07.630426305.59913112.07.940157347.45753312.07.030768235.24771912.07.778227325.08664712.07.676198311.54036112.07.471320285.59183812.09.357814592.74546112.07.497876288.86223912.07.477308286.32685912.06.089015150.52865212.08.458776426.58020812.07.625984305.02696612.08.655849459.75724012.07.073400239.80438212.07.573978298.38687812.07.160760249.34918912.07.293491264.39231112.07.571402298.06077212.07.650571316.68393612.050	