

FY2045 Quantum Mechanics I

Fall 2023

Henning Goa Hugdal

Week 10

Time-independent perturbation theory

General formulation

In most situations, a system is defined by a Hamiltonian H for which the Schrödinger equation cannot be solved exactly. However, if $H = H_0 + V$, where V is a small, time-independent term — a **perturbation** — and we know the solutions to

$$H_0|n\rangle = E_n^0|n\rangle, \quad (1)$$

we can find **approximate** solutions of


$$H|\psi_n\rangle = E_n|\psi_n\rangle, \quad (2)$$

using **time-independent perturbation theory**.

Examples of perturbations

Anharmonic oscillator:

$$H = H_0 + \lambda x^4$$



CC BY-NC; Ümit Kaya via LibreTexts

Zeemann effect, $H_Z = -\boldsymbol{\mu}_S \cdot \mathbf{B}$.

Relativistic energy correction

$$E = \sqrt{m^2 c^4 + c^2 p^2} \approx mc^2 + \frac{p^2}{2m} - \frac{p^4}{m^3 c^2} + \dots$$

Spin-orbit coupling

$$H_{SO} = \xi \mathbf{L} \cdot \mathbf{S}.$$

Hyperfine splitting of atomic levels due to spin-spin interactions between electrons and nucleus.

Non-degenerate perturbation theory

Write

$$H = H_0 + \lambda V,$$

with “book keeping device” $\lambda \in [0, 1]$.

Analogy — Roots of polynomials


$$x^2 + \lambda x - c = 0.$$

If λ is small, we use ansatz

$$x = x_0 + \lambda x_1 + \lambda^2 x_2 + \dots,$$

to get approximate solution

$$x \approx \pm\sqrt{c} - \frac{\lambda}{2} \pm \frac{\lambda^2}{8\sqrt{c}} + \mathcal{O}(\lambda^3).$$



Non-degenerate perturbation theory

We write $H = H_0 + \lambda V$, where $\lambda \in [0, 1]$. When $\lambda \rightarrow 0$, we should have

$$E_n \rightarrow E_n^0 \text{ and } |\psi_n\rangle \rightarrow |n\rangle.$$

We therefore write

$$E_n = E_n^0 + \lambda E_n^{(1)} + \lambda^2 E_n^{(2)} + \dots, \quad (3)$$

$$|\psi_n\rangle = |n\rangle + \lambda |n^{(1)}\rangle + \lambda^2 |n^{(2)}\rangle + \dots, \quad (4)$$

where we have assumed **non-degenerate** states. Inserting into the SE, eq. (2), and collecting like powers of λ , we get

$$\lambda^0 : \quad H_0 |n\rangle = E_n^0 |n\rangle,$$

$$\lambda^1 : \quad H_0 |n^{(1)}\rangle + V |n\rangle = E_n^0 |n^{(1)}\rangle + E_n^{(1)} |n\rangle, \quad (5)$$

$$\lambda^2 : \quad H_0 |n^{(2)}\rangle + V |n^{(1)}\rangle = E_n^0 |n^{(2)}\rangle + E_n^{(1)} |n^{(1)}\rangle + E_n^{(2)} |n\rangle. \quad (6)$$

Non-degenerate perturbation theory

First-order corrections

$$\underline{E_n^{(1)} = \langle n|V|n\rangle},$$

$$\underline{|n^{(1)}\rangle = \sum_{m \neq n} \frac{\langle m|V|n\rangle}{E_n^0 - E_m^0} |m\rangle}.$$

Non-degenerate perturbation theory

First-order corrections

$$E_n^{(1)} = \langle n|V|n\rangle,$$

$$|n^{(1)}\rangle = \sum_{m \neq n} \frac{\langle m|V|n\rangle}{E_n^0 - E_m^0} |m\rangle.$$

Second-order energies

$$E_n^{(2)} = \sum_{m \neq n} \frac{|\langle m|V|n\rangle|^2}{E_n^0 - E_m^0}.$$

Summary

The solution to the eigenvalue problem

$$(H_0 + \lambda V)|\psi_n\rangle = E_n|\psi_n\rangle$$

is

$$|\psi_n\rangle = |n\rangle + \sum_{m \neq n} \frac{\langle m|\lambda V|n\rangle}{E_n^0 - E_m^0} |m\rangle + \mathcal{O}(\lambda^2),$$

$$E_n = E_n^0 + \langle n|\lambda V|n\rangle + \sum_{m \neq n} \frac{|\langle m|\lambda V|n\rangle|^2}{E_n^0 - E_m^0} + \mathcal{O}(\lambda^3).$$

Example — Perturbed harmonic oscillator


A harmonic oscillator is perturbed by force F in positive x -direction:

$$V = -Fx.$$

The perturbed Hamiltonian is

$$H = H_0 + V = \frac{\hat{p}_x^2}{2m} + \frac{1}{2}m\omega^2x^2 - Fx.$$

What are the corrections to the energies and eigenstates?



Example — Perturbed harmonic oscillator

To second order in F we find the energies

$$\begin{aligned} E_n &= E_n^0 + E_n^{(1)} + E_n^{(2)} \\ &= \hbar\omega \left[n + \frac{1}{2} \right] - \frac{F^2}{2m\omega^2}, \end{aligned}$$

and to first order the state vectors

$$\begin{aligned} |\psi_n\rangle &= \left(1 - \frac{ix_0\hat{p}_x}{\hbar} \right) |n\rangle, \\ \Rightarrow \psi_n(x) &= \left(1 - x_0 \frac{d}{dx} \right) \psi_n^{(0)}(x), \end{aligned}$$

with $x_0 = \frac{F}{m\omega^2}$.

Example — Perturbed harmonic oscillator

Exact solution

Rewrite perturbed Hamiltonian

$$H = \frac{\hat{p}_x^2}{2m} + \frac{1}{2}m\omega^2(x - x_0)^2 - \frac{F^2}{2m\omega^2}.$$

Hence, we have exact solution

$$E_n = \hbar\omega \left[n + \frac{1}{2} \right] - \frac{F^2}{2m\omega^2},$$

$$\psi_n(x) = \psi_n^{(0)}(x - x_0) = \psi_n^{(0)}(x) - x_0 \frac{d}{dx} \psi_n^{(0)}(x) \dots,$$

which agrees exactly with the second order energy correction, and to first order with the wavefunction correction.

