

FY2045 Quantum Mechanics I

Fall 2023

Henning Goa Hugdal

Week 3

General formulation of QM

Dirac's $\langle \text{bra} | \text{ket} \rangle$ notation

State vector

A quantum mechanical state of a system is described by a state vector

$$|\psi\rangle$$

in a complex, linear vector space \mathcal{H} — Hilbert space.

Dual vector

For each vector $|a\rangle$ we define the dual vector $\langle a|$ in the dual space \mathcal{H}^* ,

$$|a\rangle \xleftrightarrow{\text{dual}} \langle a|,$$

so that we can define the scalar (inner) product of vectors $|a\rangle$ and $|b\rangle$

$$\langle a||b\rangle \equiv \langle a|b\rangle \in \mathbb{C},$$

with $\langle a|b\rangle = \langle b|a\rangle^*$.

Example

Probability amplitude for particle arriving at point x :


$$\langle \text{Particle arrives at } x | \text{particle leaves } s \rangle$$

or simply

$$\langle x | s \rangle.$$

Can go through either slit 1 or 2:

$$\langle x | s \rangle = \langle x | 1 \rangle \langle 1 | s \rangle + \langle x | 2 \rangle \langle 2 | s \rangle.$$



Based on Ch. 3 of Vol. III in the Feynman Lectures.

Interpretation

The wavefunction is probability amplitude of finding state $|\psi\rangle$ at point x :

$$\psi(x) = \langle x|\psi\rangle.$$

The momentum wavefunction is probability amplitude of finding state $|\psi\rangle$ with momentum p :

$$\phi(p) = \langle p|\psi\rangle.$$

The probability amplitude of finding state $|\psi\rangle$ with energy E_n :

$$\langle \psi_n|\psi\rangle.$$

Completeness

n linearly independent vectors $|1\rangle, |2\rangle, |3\rangle, \dots$ span \mathcal{H} if $\forall |\psi\rangle \in \mathcal{H}$ we have

$$|\psi\rangle = \sum_{k=1}^n c_n |k\rangle.$$

Assuming orthonormality $\langle m|k\rangle = \delta_{mk}$,

$$\Rightarrow |\psi\rangle = \sum_k \langle k|\psi\rangle |k\rangle = \sum_k |k\rangle \langle k|\psi\rangle,$$

meaning we have the completeness relation

$$\sum_k |k\rangle \langle k| = 1.$$

Operators

An operator \hat{A} applied to a vector $|a\rangle \in \mathcal{H}$ results in a new vector $|c\rangle \in \mathcal{H}$,

$$\hat{A}|a\rangle = |c\rangle.$$

Adjoint or Hermitian conjugate of operator:

$$\langle a|\hat{A}^\dagger|b\rangle = \langle b|\hat{A}|a\rangle^* \quad \forall |a\rangle, |b\rangle \in \mathcal{H}.$$

meaning that we have the dual vector

$$\hat{A}|a\rangle \xleftrightarrow{\text{dual}} \langle a|\hat{A}^\dagger.$$

Properties

$$(\hat{A}^\dagger)^\dagger = \hat{A},$$

$$(\alpha\hat{A})^\dagger = \alpha^*\hat{A}^\dagger,$$

$$(\hat{A}\hat{B})^\dagger = \hat{B}^\dagger\hat{A}^\dagger.$$

For a Hermitian (self-adjoint) matrix

$$\hat{A}^\dagger = \hat{A}.$$

Eigenvectors and eigenvalues

The eigenvectors $|\alpha\rangle$ and eigenvalues λ_α of an operator \hat{A} are defined by

$$\hat{A}|\alpha\rangle = \lambda_\alpha|\alpha\rangle.$$

The set of eigenvectors $\{|\alpha\rangle\}$ corresponding to physical quantities is assumed to be complete — they form a basis set that span \mathcal{H} .

The eigenvectors of a **Hermitian** operator are **real**.

Examples

Energy:

$$\hat{H}|n\rangle = E_n|n\rangle.$$

Position:

$$\hat{x}|x'\rangle = x'|x'\rangle.$$

Momentum:

$$\hat{p}|p'\rangle = p'|p'\rangle.$$

Postulates in general formulation

Postulate A

Each observable quantity F corresponds to a linear, Hermitian operator \hat{F} in Hilbert space. The operators for a generalized coordinate q_n and generalized momentum p_n fulfill

$$[\hat{q}_n, \hat{p}_n] = \hat{q}_n \hat{p}_n - \hat{p}_n \hat{q}_n = i\hbar.$$

Postulate C

The expectation value of an observable F , given the state $|\Psi\rangle$, is

$$\langle F \rangle = \langle \Psi | \hat{F} | \Psi \rangle.$$

Postulate B

Each state of a physical system is represented by a state vector $|\Psi, t\rangle$ in Hilbert space with length 1, $\langle \Psi, t | \Psi, t \rangle = 1$. The vector fulfills the time-dependent SE

$$i\hbar \frac{\partial}{\partial t} |\Psi, t\rangle = \hat{H} |\Psi, t\rangle.$$

Postulate D

The measurement of an observable F yields one of the eigenvalues f_n of the corresponding operator \hat{F} .

Position representation

Eigenvectors $|x'\rangle$ of operator \hat{x} :

$$\hat{x}|x'\rangle = x'|x'\rangle,$$

with eigenvalue x' taking continuous values.

$|x'\rangle$ is δ -function normalized

$$\langle x''|x'\rangle = \delta(x'' - x'),$$

with completeness relation

$$\int dx' |x'\rangle \langle x'| = \mathbb{1}.$$

Position space wavefunction

Given a state vector $|\psi\rangle$, the position space wavefunction is given by

$$\psi(x') = \langle x'|\psi\rangle,$$

the projection of $|\psi\rangle$ on the position basis vector $|x'\rangle$.

Operators

For an operator \hat{F} , which is a function of \hat{x} and \hat{p} , we have

$$\langle x''|F(\hat{p}, \hat{x})|x'\rangle = F\left(\frac{\hbar}{i} \frac{\partial}{\partial x''}, x''\right) \delta(x'' - x').$$

Momentum formulation

Eigenvectors $|p\rangle$ of operator \hat{p} :

$$\hat{p}|p\rangle = p|p\rangle,$$

with eigenvalue p taking continuous values.
 $|p\rangle$ is δ -function normalized

$$\langle p'|p\rangle = \delta(p' - p),$$

with completeness relation

$$\int dp |p\rangle \langle p| = \mathbb{1}.$$

Momentum space wavefunction

Given a state vector $|\psi\rangle$, the momentum space wavefunction is given by

$$\phi(p) = \langle p|\psi\rangle,$$

the projection of $|\psi\rangle$ on the momentum basis vector $|p\rangle$.

Relation between $|p\rangle$ and $|x\rangle$

Projecting $|p\rangle$ on the position basis $|x\rangle$, we get the momentum eigenfunctions in the position formulation,

$$\langle x|p\rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{ipx/\hbar}.$$