

DataCORE

Parallel Tracks – Core Analysis & Multiple Regression

Afternoon: DataCORE

1. **Multiple Regression + Interpretation** (1:00–2:45)
2. **Scientific Storytelling — Figure + Interpretation Pairing** (2:45–3:00)



Multiple regression with 2+ predictors, full output table, assumption checks, and a "**holding constant**" interpretation.

Shared dataset: the wage survey — education, gender, region, experience_years, hourly_wage.

1:00 – 2:45

Why **multiple** regression?

- Simple regression: one predictor (wage ~ experience).
- But wage also depends on **education**. If experienced workers are also more educated, simple regression **confounds** the two.
- Multiple regression estimates each predictor's effect **holding the others constant**.

DV = hourly wage · **IVs** = years of experience **and** education level.

$$= \text{wage} = \beta_0 + \beta_1 \cdot \text{experience} + \beta_2 \cdot \text{education}$$

Each β is a *partial* slope — the effect of that variable with the rest held fixed.

Fit OLS with 2+ predictors

Python (statsmodels)

R (base)

```
model = smf.ols("hourly_wage ~ experience_years + C(education)", data=wages).fit()
print(model.summary().tables[1])
```

```
=====
              coef      std err          t      P>|t|      [0.025      0.975]
-----
Intercept          19.7837         0.620      31.911      0.000      18.562      21.006
C(education)[T.Bachelor]         4.1288         0.625       6.601      0.000       2.896       5.362
C(education)[T.Graduate]         9.7785         0.814      12.010      0.000       8.174      11.383
C(education)[T.High School]      -2.5176         0.643       -3.913      0.000      -3.786      -1.250
experience_years         0.4177         0.033      12.535      0.000       0.352       0.483
=====
```

 `C(education)` / a factor adds one row **per category** (vs. a baseline). Each coefficient is that group's gap from the baseline, **holding experience constant**.

Read the full output

Python R

```
print(f"R-squared      = {model.rsquared:.3f}")
```

R-squared = 0.695

```
print(f"Adj. R-squared= {model.rsquared_adj:.3f}")
```

Adj. R-squared= 0.689

```
print(f"F-statistic    = {model.fvalue:.1f}    (p = {model.f_pvalue:.2e})")
```

F-statistic = 122.2 (p = 3.27e-54)

```
print(f"N              = {int(model.nobs)}")
```

N = 220

 **Each coefficient:** the effect of that variable. **Std err / t / p:** is it distinguishable from 0? **R²:** variation explained. **F-stat:** is the model as a whole useful?

Say it the careful way

“ ***Holding education constant***, each additional year of experience is associated with a ***\\$0.42*** change in hourly wage ($p < 0.001$).”

The template:

*"Holding **[controls]** constant, a one-unit increase in **[IV]** is associated with a **[coef]** unit change in **[DV]** ($p = __$)."*

"Associated with," not "causes."

Even with controls, this is observational — unmeasured factors may remain.


Check the assumptions

Residuals · Python

Q-Q · Python

R

```
plt.scatter(model.fittedvalues, model.resid, alpha=0.5, color="#4a7fb5")
plt.axhline(0, color="#D55E00", lw=2)
plt.xlabel("Fitted values"); plt.ylabel("Residuals")
plt.title("Residuals vs. fitted")
plt.tight_layout(); plt.show()
```



 **Residuals vs. fitted:** want a shapeless cloud around 0 (no funnel, no curve). **Q-Q plot:** points near the line = roughly normal residuals.

Scholars Mission Sheet

Complete the **Session 9 Scholars Mission Sheet** on your own data.

- 1 Choose a **variable pair** (DV + key IV) plus at least one **control**.
- 2 State the **relationship found** – coefficient, sign, significance.
- 3 Write the **plain-language** "holding constant" interpretation.
- 4 Name **one causal limitation** – what could still confound it?

45:00

2:45 – 3:00 · Scientific Storytelling

Make them a matched pair

A great results slide is a **chart + one sentence** that fit together perfectly.

1

Write a **1-sentence interpretation** of your strongest chart.

2

Read it aloud **without** showing the chart — does it stand on its own?

3

Now show the chart — does it **prove** the sentence?

The interpretation tells the audience what to see; the chart proves it.

Example: the sentence "*Wages rise steadily with experience*" paired with a scatter that clearly trends up.

15:00

