

COMPSCI 514: ALGORITHMS FOR DATA SCIENCE

Andrew McGregor

Lecture 14

Next Few Classes: Low-rank approximation, the SVD, and principal component analysis (PCA).

- Reduce d -dimensional data points to a smaller dimension m .
- Like JL, **compression is linear**, i.e., by applying a matrix.
- Chose matrix taking into account **structure of dataset**.
- Can give better compression than random projection.


Next Few Classes: Low-rank approximation, the SVD, and principal component analysis (PCA).

- Reduce d -dimensional data points to a smaller dimension m .
- Like JL, **compression is linear**, i.e., by applying a matrix.
- Chose matrix taking into account **structure of dataset**.
- Can give better compression than random projection.

Will be using a fair amount of linear algebra: orthogonal basis, column/row span, eigenvectors, etc,


EMBEDDING WITH ASSUMPTIONS

Assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie in any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



EMBEDDING WITH ASSUMPTIONS

Assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie in any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .




Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all \vec{x}_i, \vec{x}_j :

$$\|\mathbf{V}^T \vec{x}_i - \mathbf{V}^T \vec{x}_j\|_2 = \|\vec{x}_i - \vec{x}_j\|_2.$$

EMBEDDING WITH ASSUMPTIONS

Assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie in any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all \vec{x}_i, \vec{x}_j :

$$\|\mathbf{V}^T \vec{x}_i - \mathbf{V}^T \vec{x}_j\|_2 = \|\vec{x}_i - \vec{x}_j\|_2.$$

- $\mathbf{V}^T \in \mathbb{R}^{k \times d}$ is a linear embedding of $\vec{x}_1, \dots, \vec{x}_n$ into k dimensions with **no distortion**.

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T y\|_2 = \|y\|_2.$$

Proof:

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T y\|_2 = \|y\|_2.$$

Proof:

- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V} \vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T y\|_2 = \|y\|_2.$$

Proof:

- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V} \vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$
- $\|\vec{y}\|_2^2 = \vec{y}^T \vec{y}$

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T y\|_2 = \|y\|_2.$$

Proof:

- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V}\vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$
- $\|\vec{y}\|_2^2 = \vec{y}^T \vec{y} = (\mathbf{V}\vec{c})^T (\mathbf{V}\vec{c})$

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T y\|_2 = \|y\|_2.$$

Proof:

- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V}\vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$
- $\|\vec{y}\|_2^2 = \vec{y}^T \vec{y} = (\mathbf{V}\vec{c})^T (\mathbf{V}\vec{c}) = \vec{c}^T \mathbf{V}^T \mathbf{V} \vec{c}$

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T \vec{y}\|_2 = \|\vec{y}\|_2.$$

Proof:

- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V} \vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$
- $\|\vec{y}\|_2^2 = \vec{y}^T \vec{y} = (\mathbf{V} \vec{c})^T (\mathbf{V} \vec{c}) = \vec{c}^T \mathbf{V}^T \mathbf{V} \vec{c}$
- $\|\mathbf{V}^T \vec{y}\|_2^2 = (\mathbf{V}^T \vec{y})^T (\mathbf{V}^T \vec{y}) = \vec{y}^T \mathbf{V} \mathbf{V}^T \vec{y}$

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T \vec{y}\|_2 = \|\vec{y}\|_2.$$

Proof:

- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V} \vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$
- $\|\vec{y}\|_2^2 = \vec{y}^T \vec{y} = (\mathbf{V} \vec{c})^T (\mathbf{V} \vec{c}) = \vec{c}^T \mathbf{V}^T \mathbf{V} \vec{c}$
- $\|\mathbf{V}^T \vec{y}\|_2^2 = (\mathbf{V}^T \vec{y})^T (\mathbf{V}^T \vec{y}) = \vec{y}^T \mathbf{V} \mathbf{V}^T \vec{y} = \vec{c}^T \mathbf{V}^T \mathbf{V} \mathbf{V}^T \mathbf{V} \vec{c}$

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T y\|_2 = \|y\|_2.$$

Proof:

- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V}\vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$
- $\|\vec{y}\|_2^2 = \vec{y}^T \vec{y} = (\mathbf{V}\vec{c})^T (\mathbf{V}\vec{c}) = \vec{c}^T \mathbf{V}^T \mathbf{V} \vec{c}$
- $\|\mathbf{V}^T \vec{y}\|_2^2 = (\mathbf{V}^T \vec{y})^T (\mathbf{V}^T \vec{y}) = \vec{y}^T \mathbf{V} \mathbf{V}^T \vec{y} = \vec{c}^T \mathbf{V}^T \mathbf{V} \mathbf{V}^T \mathbf{V} \vec{c}$
- But $\mathbf{V}^T \mathbf{V} = I$

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T \vec{y}\|_2 = \|\vec{y}\|_2.$$

Proof:

- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V} \vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$
- $\|\vec{y}\|_2^2 = \vec{y}^T \vec{y} = (\mathbf{V} \vec{c})^T (\mathbf{V} \vec{c}) = \vec{c}^T \mathbf{V}^T \mathbf{V} \vec{c}$
- $\|\mathbf{V}^T \vec{y}\|_2^2 = (\mathbf{V}^T \vec{y})^T (\mathbf{V}^T \vec{y}) = \vec{y}^T \mathbf{V} \mathbf{V}^T \vec{y} = \vec{c}^T \mathbf{V}^T \mathbf{V} \mathbf{V}^T \mathbf{V} \vec{c}$
- But $\mathbf{V}^T \mathbf{V} = I$ since

$$[\mathbf{V}^T \mathbf{V}]_{i,j} = \vec{v}_i^T \vec{v}_j = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}$$

DOT PRODUCT TRANSFORMATION

Claim: Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns. For all $\vec{y} \in \mathcal{V}$:

$$\|\mathbf{V}^T \vec{y}\|_2 = \|\vec{y}\|_2.$$

Proof:


- If $\vec{y} = \sum_i c_i \vec{v}_i$ then $\vec{y} = \mathbf{V} \vec{c}$ where $\vec{c}^T = (c_1, \dots, c_k)$
- $\|\vec{y}\|_2^2 = \vec{y}^T \vec{y} = (\mathbf{V} \vec{c})^T (\mathbf{V} \vec{c}) = \vec{c}^T \mathbf{V}^T \mathbf{V} \vec{c}$
- $\|\mathbf{V}^T \vec{y}\|_2^2 = (\mathbf{V}^T \vec{y})^T (\mathbf{V}^T \vec{y}) = \vec{y}^T \mathbf{V} \mathbf{V}^T \vec{y} = \vec{c}^T \mathbf{V}^T \mathbf{V} \mathbf{V}^T \mathbf{V} \vec{c}$
- But $\mathbf{V}^T \mathbf{V} = I$ since

$$[\mathbf{V}^T \mathbf{V}]_{i,j} = \vec{v}_i^T \vec{v}_j = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}$$

- So $\|\vec{y}\|_2^2 = \vec{c}^T \vec{c} = \|\mathbf{V}^T \vec{y}\|_2^2$.


EMBEDDING WITH ASSUMPTIONS

Now assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie **close to** any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .




EMBEDDING WITH ASSUMPTIONS

Now assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie close to any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



EMBEDDING WITH ASSUMPTIONS


Now assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie close to any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



Letting $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns, $\mathbf{V}^T \vec{x}_i \in \mathbb{R}^k$ is still a good embedding for $x_i \in \mathbb{R}^d$.

EMBEDDING WITH ASSUMPTIONS


Now assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie close to any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



Letting $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns, $\mathbf{V}^T \vec{x}_i \in \mathbb{R}^k$ is still a good embedding for $x_i \in \mathbb{R}^d$. The key idea behind low-rank approximation and principal component analysis (PCA).

EMBEDDING WITH ASSUMPTIONS

Now assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie close to any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



Letting $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns, $\mathbf{V}^T \vec{x}_i \in \mathbb{R}^k$ is still a good embedding for $x_i \in \mathbb{R}^d$. The key idea behind low-rank approximation and principal component analysis (PCA).

- How do we find \mathcal{V} and \mathbf{V} ?
- How good is the embedding?

LOW-RANK FACTORIZATION

Claim: $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace $\mathcal{V} \Leftrightarrow$ the data matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$ has rank $\leq k$.


$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK FACTORIZATION

Claim: $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace $\mathcal{V} \Leftrightarrow$ the data matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$ has rank $\leq k$.

- Letting $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} , can write any \vec{x}_i as:

$$\vec{x}_i = \mathbf{V}\vec{c}_i = c_{i,1} \cdot \vec{v}_1 + c_{i,2} \cdot \vec{v}_2 + \dots + c_{i,k} \cdot \vec{v}_k.$$



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.


LOW-RANK FACTORIZATION

Claim: $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace $\mathcal{V} \Leftrightarrow$ the data matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$ has rank $\leq k$.

- Letting $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} , can write any \vec{x}_i as:

$$\vec{x}_i = \mathbf{V}\vec{c}_i = c_{i,1} \cdot \vec{v}_1 + c_{i,2} \cdot \vec{v}_2 + \dots + c_{i,k} \cdot \vec{v}_k.$$

- So $\vec{v}_1, \dots, \vec{v}_k$ span the rows of \mathbf{X} and thus $\text{rank}(\mathbf{X}) \leq k$.



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Claim: $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ lie in a k -dimensional subspace $\mathcal{V} \Leftrightarrow$ the data matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$ has rank $\leq k$.


- Every data point \vec{x}_i (row of \mathbf{X}) can be written as $\vec{x}_i = \mathbf{V}\vec{c}_i = c_{i,1} \cdot \vec{v}_1 + \dots + c_{i,k} \cdot \vec{v}_k$.

$\vec{x}_1, \dots, \vec{x}_n$: data points (in \mathbb{R}^d), \mathcal{V} : k -dimensional subspace of \mathbb{R}^d , $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Claim: $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ lie in a k -dimensional subspace $\mathcal{V} \Leftrightarrow$ the data matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$ has rank $\leq k$.

- Every data point \vec{x}_i (row of \mathbf{X}) can be written as

$$\vec{x}_i = \mathbf{V} \vec{c}_i = c_{i,1} \cdot \vec{v}_1 + \dots + c_{i,k} \cdot \vec{v}_k.$$




$\vec{x}_1, \dots, \vec{x}_n$: data points (in \mathbb{R}^d), \mathcal{V} : k -dimensional subspace of \mathbb{R}^d , $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Claim: $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ lie in a k -dimensional subspace $\mathcal{V} \Leftrightarrow$ the data matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$ has rank $\leq k$.

- Every data point \vec{x}_i (row of \mathbf{X}) can be written as

$$\vec{x}_i = \mathbf{V} \vec{c}_i = c_{i,1} \cdot \vec{v}_1 + \dots + c_{i,k} \cdot \vec{v}_k.$$




- \mathbf{X} can be represented by $(n + d) \cdot k$ parameters vs. $n \cdot d$.

$\vec{x}_1, \dots, \vec{x}_n$: data points (in \mathbb{R}^d), \mathcal{V} : k -dimensional subspace of \mathbb{R}^d , $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Claim: $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ lie in a k -dimensional subspace $\mathcal{V} \Leftrightarrow$ the data matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$ has rank $\leq k$.

- Every data point \vec{x}_i (row of \mathbf{X}) can be written as

$$\vec{x}_i = \mathbf{V} \vec{c}_i = c_{i,1} \cdot \vec{v}_1 + \dots + c_{i,k} \cdot \vec{v}_k.$$




- \mathbf{X} can be represented by $(n + d) \cdot k$ parameters vs. $n \cdot d$.
- The rows of \mathbf{X} are spanned by k vectors: the columns of $\mathbf{V} \Rightarrow$ the columns of \mathbf{X} are spanned by k vectors: the columns of \mathbf{C} .

$\vec{x}_1, \dots, \vec{x}_n$: data points (in \mathbb{R}^d), \mathcal{V} : k -dimensional subspace of \mathbb{R}^d , $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK FACTORIZATION


Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as $\mathbf{X} = \mathbf{CV}^T$.



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK FACTORIZATION

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as $\mathbf{X} = \mathbf{C}\mathbf{V}^T$.




Exercise: What is this coefficient matrix \mathbf{C} ? **Hint:** Use that $\mathbf{V}^T\mathbf{V} = \mathbf{I}$.

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK FACTORIZATION

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as $\mathbf{X} = \mathbf{C}\mathbf{V}^T$.




Exercise: What is this coefficient matrix \mathbf{C} ? **Hint:** Use that $\mathbf{V}^T\mathbf{V} = \mathbf{I}$.

$$\bullet \quad \mathbf{X} = \mathbf{C}\mathbf{V}^T \implies \mathbf{X}\mathbf{V} = \mathbf{C}\mathbf{V}^T\mathbf{V}$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK FACTORIZATION

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as $\mathbf{X} = \mathbf{C}\mathbf{V}^T$.




Exercise: What is this coefficient matrix \mathbf{C} ? **Hint:** Use that $\mathbf{V}^T\mathbf{V} = \mathbf{I}$.

$$\bullet \quad \mathbf{X} = \mathbf{C}\mathbf{V}^T \implies \mathbf{X}\mathbf{V} = \mathbf{C}\mathbf{V}^T\mathbf{V} \implies \mathbf{X}\mathbf{V} = \mathbf{C}$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK FACTORIZATION

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as $\mathbf{X} = \mathbf{C}\mathbf{V}^T$.



Exercise: What is this coefficient matrix \mathbf{C} ? **Hint:** Use that $\mathbf{V}^T \mathbf{V} = \mathbf{I}$.

$$\bullet \quad \mathbf{X} = \mathbf{C}\mathbf{V}^T \implies \mathbf{X}\mathbf{V} = \mathbf{C}\mathbf{V}^T \mathbf{V} \implies \mathbf{X}\mathbf{V} = \mathbf{C}$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{C}\mathbf{V}^T.$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{XV}\mathbf{V}^T.$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{X}\mathbf{V}\mathbf{V}^T.$$

- $\mathbf{V}\mathbf{V}^T$ is a **projection matrix**, which projects the rows of \mathbf{X} (the data points $\vec{x}_1, \dots, \vec{x}_n$) onto the subspace \mathcal{V} .


$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

PROJECTION VIEW

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{X}\mathbf{V}\mathbf{V}^T.$$

- $\mathbf{V}\mathbf{V}^T$ is a **projection matrix**, which projects the rows of \mathbf{X} (the data points $\vec{x}_1, \dots, \vec{x}_n$) onto the subspace \mathcal{V} .




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

PROJECTION VIEW

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{X}\mathbf{V}\mathbf{V}^T.$$

- $\mathbf{V}\mathbf{V}^T$ is a **projection matrix**, which projects the rows of \mathbf{X} (the data points $\vec{x}_1, \dots, \vec{x}_n$) onto the subspace \mathcal{V} .




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

PROJECTION VIEW

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{X}\mathbf{V}\mathbf{V}^T.$$

- $\mathbf{V}\mathbf{V}^T$ is a **projection matrix**, which projects the rows of \mathbf{X} (the data points $\vec{x}_1, \dots, \vec{x}_n$) onto the subspace \mathcal{V} .




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROXIMATION

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie **close to** a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be **approximated as:**

$$\mathbf{X} \approx \mathbf{XV}\mathbf{V}^T$$




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROXIMATION

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie **close to** a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be **approximated as**:

$$\mathbf{X} \approx \mathbf{XV}\mathbf{V}^T$$




Note: $\mathbf{XV}\mathbf{V}^T$ has rank k . It is a **low-rank approximation** of \mathbf{X} .

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROXIMATION

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie **close to** a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be **approximated as**:

$$\mathbf{X} \approx \mathbf{XV}\mathbf{V}^T$$



Note: $\mathbf{XV}\mathbf{V}^T$ has rank k . It is a **low-rank approximation** of \mathbf{X} .

$$\mathbf{XV}\mathbf{V}^T = \arg \min_{\mathbf{B} \text{ with rows in } \mathcal{V}} \|\mathbf{X} - \mathbf{B}\|_F^2 = \arg \min_{\mathbf{B} \text{ with rows in } \mathcal{V}} \sum_{i,j} (\mathbf{X}_{i,j} - \mathbf{B}_{i,j})^2.$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthonormal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROXIMATION

So Far: If $\vec{x}_1, \dots, \vec{x}_n$ lie close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as:

$$\mathbf{X} \approx \mathbf{XV}\mathbf{V}^T.$$

This is the closest approximation to \mathbf{X} with rows in \mathcal{V} (i.e., in the column span of \mathbf{V}).

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROXIMATION

So Far: If $\vec{x}_1, \dots, \vec{x}_n$ lie close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as:

$$\mathbf{X} \approx \mathbf{XV}\mathbf{V}^T.$$

This is the closest approximation to \mathbf{X} with rows in \mathcal{V} (i.e., in the column span of \mathbf{V}).

- Letting $(\mathbf{XV}\mathbf{V}^T)_i, (\mathbf{XV}\mathbf{V}^T)_j$ be the i^{th} and j^{th} projected data points,
$$\|(\mathbf{XV}\mathbf{V}^T)_i - (\mathbf{XV}\mathbf{V}^T)_j\|_2 = \|[(\mathbf{XV})_i - (\mathbf{XV})_j]\mathbf{V}^T\|_2 = \|[(\mathbf{XV})_i - (\mathbf{XV})_j]\|_2.$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROXIMATION

So Far: If $\vec{x}_1, \dots, \vec{x}_n$ lie close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as:

$$\mathbf{X} \approx \mathbf{XV}\mathbf{V}^T.$$

This is the closest approximation to \mathbf{X} with rows in \mathcal{V} (i.e., in the column span of \mathbf{V}).

- Letting $(\mathbf{XV}\mathbf{V}^T)_i, (\mathbf{XV}\mathbf{V}^T)_j$ be the i^{th} and j^{th} projected data points,
$$\|(\mathbf{XV}\mathbf{V}^T)_i - (\mathbf{XV}\mathbf{V}^T)_j\|_2 = \|[(\mathbf{XV})_i - (\mathbf{XV})_j]\mathbf{V}^T\|_2 = \|[(\mathbf{XV})_i - (\mathbf{XV})_j]\|_2.$$
- Can use $\mathbf{XV} \in \mathbb{R}^{n \times k}$ as a compressed approximate data set.

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROXIMATION

So Far: If $\vec{x}_1, \dots, \vec{x}_n$ lie close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as:

$$\mathbf{X} \approx \mathbf{XV}\mathbf{V}^T.$$

This is the closest approximation to \mathbf{X} with rows in \mathcal{V} (i.e., in the column span of \mathbf{V}).

- Letting $(\mathbf{XV}\mathbf{V}^T)_i, (\mathbf{XV}\mathbf{V}^T)_j$ be the i^{th} and j^{th} projected data points,
$$\|(\mathbf{XV}\mathbf{V}^T)_i - (\mathbf{XV}\mathbf{V}^T)_j\|_2 = \|[(\mathbf{XV})_i - (\mathbf{XV})_j]\mathbf{V}^T\|_2 = \|[(\mathbf{XV})_i - (\mathbf{XV})_j]\|_2.$$
- Can use $\mathbf{XV} \in \mathbb{R}^{n \times k}$ as a compressed approximate data set.

Key question is how to find the subspace \mathcal{V} and correspondingly \mathbf{V} .

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.