


COMPSCI 514: ALGORITHMS FOR DATA SCIENCE

Andrew McGregor

Lecture 15

LAST CLASS: EMBEDDING WITH ASSUMPTIONS

Set Up: Assume that data points $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ lie in some k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns.

$$\|\mathbf{V}^T \vec{x}_i - \mathbf{V}^T \vec{x}_j\|_2^2 = \|\vec{x}_i - \vec{x}_j\|_2^2.$$


Letting $\tilde{x}_i = \mathbf{V}^T \vec{x}_i$, we have a perfect embedding from \mathcal{V} into \mathbb{R}^k .

LAST CLASS: PROJECTION VIEW

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{XV}\mathbf{V}^T = \mathbf{CV}^T$$

- $\mathbf{V}\mathbf{V}^T$ is a **projection matrix**, which projects the rows of \mathbf{X} (the data points $\vec{x}_1, \dots, \vec{x}_n$) onto the subspace \mathcal{V} .




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LAST CLASS: PROJECTION VIEW

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{XV}\mathbf{V}^T = \mathbf{CV}^T$$

- $\mathbf{V}\mathbf{V}^T$ is a **projection matrix**, which projects the rows of \mathbf{X} (the data points $\vec{x}_1, \dots, \vec{x}_n$) onto the subspace \mathcal{V} .




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LAST CLASS: PROJECTION VIEW

Claim: If $\vec{x}_1, \dots, \vec{x}_n$ lie in a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be written as

$$\mathbf{X} = \mathbf{XV}\mathbf{V}^T = \mathbf{CV}^T$$


- $\mathbf{V}\mathbf{V}^T$ is a **projection matrix**, which projects the rows of \mathbf{X} (the data points $\vec{x}_1, \dots, \vec{x}_n$) onto the subspace \mathcal{V} .



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.


EMBEDDING WITH ASSUMPTIONS

Assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie **close to** any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



EMBEDDING WITH ASSUMPTIONS


Assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie close to any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



Will show above in homework. Today's focus: How do we find \mathcal{V} and \mathbf{V} ?

EMBEDDING WITH ASSUMPTIONS

Assume that data points $\vec{x}_1, \dots, \vec{x}_n$ lie **close to** any k -dimensional subspace \mathcal{V} of \mathbb{R}^d .



Letting $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns, $\mathbf{V}^T \vec{x}_i \in \mathbb{R}^k$ is **still a good embedding for $x_i \in \mathbb{R}^d$** and \mathbf{XVV}^T is still a good approximation for \mathbf{X} :

$$\mathbf{XVV}^T = \arg \min_{\mathbf{B} \text{ with rows in } \mathcal{V}} \|\mathbf{X} - \mathbf{B}\|_F^2.$$

Will show above in homework. Today's focus: How do we find \mathcal{V} and \mathbf{V} ?

A STEP BACK: WHY LOW-RANK APPROXIMATION?

Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

A STEP BACK: WHY LOW-RANK APPROXIMATION?

Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

- The rows of \mathbf{X} can be approximately reconstructed from a basis of k vectors.

A STEP BACK: WHY LOW-RANK APPROXIMATION?


Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

- The rows of \mathbf{X} can be approximately reconstructed from a basis of k vectors.


784 dimensional vectors



projections onto 15 dimensional space



orthonormal basis v_1, \dots, v_{15}



⋮

DUAL VIEW OF LOW-RANK APPROXIMATION

Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

DUAL VIEW OF LOW-RANK APPROXIMATION

Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

- Equivalently, the columns of \mathbf{X} are approx. spanned by k vectors.

DUAL VIEW OF LOW-RANK APPROXIMATION

Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

- Equivalently, the columns of \mathbf{X} are approx. spanned by k vectors.

Linearly Dependent Variables:

	bedrooms	bathrooms	sq.ft.	floors	list price	sale price
home 1	2	2	1800	2	200,000	195,000
home 2	4	2.5	2700	1	300,000	310,000
.
.
.
home n	5	3.5	3600	3	450,000	450,000

DUAL VIEW OF LOW-RANK APPROXIMATION

Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

- Equivalently, the columns of \mathbf{X} are approx. spanned by k vectors.

Linearly Dependent Variables:

	bedrooms	bathrooms	sq.ft.	floors	list price	sale price
home 1	2	2	1800	2	200,000	195,000
home 2	4	2.5	2700	1	300,000	310,000
.
.
.
home n	5	3.5	3600	3	450,000	450,000

DUAL VIEW OF LOW-RANK APPROXIMATION

Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

- Equivalently, the columns of \mathbf{X} are approx. spanned by k vectors.

Linearly Dependent Variables:

	bedrooms	bathrooms	sq.ft.	floors	list price	sale price
home 1	2	2	1800	2	200,000	195,000
home 2	4	2.5	2700	1	300,000	310,000
.
.
.
home n	5	3.5	3600	3	450,000	450,000

DUAL VIEW OF LOW-RANK APPROXIMATION

Question: Why might we expect $\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$ to lie close to a k -dimensional subspace?

- Equivalently, the columns of \mathbf{X} are approx. spanned by k vectors.

Linearly Dependent Variables:

10000* bathrooms+ 10* (sq. ft.) \approx list price

	bedrooms	bathrooms	sq.ft.	floors	list price	sale price
home 1	2	2	1800	2	200,000	195,000
home 2	4	2.5	2700	1	300,000	310,000
.
.
.
home n	5	3.5	3600	3	450,000	450,000

Quick Exercise 1 : Show that $\mathbf{V}\mathbf{V}^T$ is **idempotent**. I.e.,
 $(\mathbf{V}\mathbf{V}^T)(\mathbf{V}\mathbf{V}^T)\vec{y} = (\mathbf{V}\mathbf{V}^T)\vec{y}$ for any $\vec{y} \in \mathbb{R}^d$.

Quick Exercise 1 : Show that $\mathbf{V}\mathbf{V}^T$ is **idempotent**. I.e., $(\mathbf{V}\mathbf{V}^T)(\mathbf{V}\mathbf{V}^T)\vec{y} = (\mathbf{V}\mathbf{V}^T)\vec{y}$ for any $\vec{y} \in \mathbb{R}^d$.

Quick Exercise 2: The projection is orthogonal to its complement: For any $\vec{y} \in \mathbb{R}^d$, $\langle \mathbf{V}\mathbf{V}^T \vec{y}, (\mathbf{I} - \mathbf{V}\mathbf{V}^T) \vec{y} \rangle = 0$

Quick Exercise 1 : Show that $\mathbf{V}\mathbf{V}^T$ is **idempotent**. I.e., $(\mathbf{V}\mathbf{V}^T)(\mathbf{V}\mathbf{V}^T)\vec{y} = (\mathbf{V}\mathbf{V}^T)\vec{y}$ for any $\vec{y} \in \mathbb{R}^d$.

Quick Exercise 2: The projection is orthogonal to its complement: For any $\vec{y} \in \mathbb{R}^d$, $\langle \mathbf{V}\mathbf{V}^T \vec{y}, (\mathbf{I} - \mathbf{V}\mathbf{V}^T) \vec{y} \rangle = 0$

Implies the Pythagorean Theorem: Show that for any $\vec{y} \in \mathbb{R}^d$,

$$\|\vec{y}\|_2^2 = \|(\mathbf{V}\mathbf{V}^T)\vec{y}\|_2^2 + \|\vec{y} - (\mathbf{V}\mathbf{V}^T)\vec{y}\|_2^2.$$

BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as \mathbf{XVV}^T and \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as \mathbf{XVV}^T and \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

$$\begin{aligned} \arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{XVV}^T\|_F^2 &= \sum_{i,j} (\mathbf{x}_{i,j} - (\mathbf{XVV}^T)_{i,j})^2 \\ &= \sum_{i=1}^n \|\vec{x}_i - \mathbf{V}\mathbf{V}^T \vec{x}_i\|_2^2 \\ &= \sum_{i=1}^n \|\vec{x}_i\|_2^2 - \|\mathbf{V}\mathbf{V}^T \vec{x}_i\|_2^2 \end{aligned}$$

BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as $\mathbf{X}\mathbf{V}\mathbf{V}^T$ and $\mathbf{X}\mathbf{V}$ gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

$$\begin{aligned} \arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 &= \sum_{i,j} (\mathbf{x}_{i,j} - (\mathbf{X}\mathbf{V}\mathbf{V}^T)_{i,j})^2 \\ &= \sum_{i=1}^n \|\vec{x}_i - \mathbf{V}\mathbf{V}^T \vec{x}_i\|_2^2 \\ &= \sum_{i=1}^n \|\vec{x}_i\|_2^2 - \|\mathbf{V}\mathbf{V}^T \vec{x}_i\|_2^2 \end{aligned}$$

So want to maximize $\sum_i \|\mathbf{V}\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_i \vec{x}_i^T \mathbf{V}\mathbf{V}^T \mathbf{V}\mathbf{V}^T \vec{x}_i = \sum_i \|\mathbf{V}^T \vec{x}_i\|_2^2$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SOLUTION VIA EIGENDECOMPOSITION

\mathbf{V} minimizing $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$ is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SOLUTION VIA EIGENDECOMPOSITION

\mathbf{V} minimizing $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$ is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2 = \sum_{j=1}^k \|\mathbf{X} \vec{v}_j\|_2^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SOLUTION VIA EIGENDECOMPOSITION

\mathbf{V} minimizing $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$ is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2 = \sum_{j=1}^k \|\mathbf{X} \vec{v}_j\|_2^2$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ greedily.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \|\mathbf{X} \vec{v}\|_2^2.$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SOLUTION VIA EIGENDECOMPOSITION

\mathbf{V} minimizing $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$ is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2 = \sum_{j=1}^k \|\mathbf{X} \vec{v}_j\|_2^2$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ greedily.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SOLUTION VIA EIGENDECOMPOSITION

\mathbf{V} minimizing $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$ is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2 = \sum_{j=1}^k \|\mathbf{X} \vec{v}_j\|_2^2$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ greedily.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SOLUTION VIA EIGENDECOMPOSITION

\mathbf{V} minimizing $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$ is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2 = \sum_{j=1}^k \|\mathbf{X} \vec{v}_j\|_2^2$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ greedily.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

...

$$\vec{v}_k = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle=0 \ \forall j < k} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SOLUTION VIA EIGENDECOMPOSITION

\mathbf{V} minimizing $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$ is given by:

$$\arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \sum_{i=1}^n \|\mathbf{V}^T \vec{x}_i\|_2^2 = \sum_{j=1}^k \sum_{i=1}^n \langle \vec{v}_j, \vec{x}_i \rangle^2 = \sum_{j=1}^k \|\mathbf{X} \vec{v}_j\|_2^2$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ greedily.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

...

$$\vec{v}_k = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle=0 \ \forall j < k} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

These are exactly the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$.

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.