

COMPSCI 514: ALGORITHMS FOR DATA SCIENCE

Andrew McGregor

Lecture 16

Last Class: Low-Rank Approximation

- When data lies in a k -dimensional subspace \mathcal{V} , we can perfectly embed into k dimensions using an orthonormal span $\mathbf{V} \in \mathbb{R}^{d \times k}$.
- When data lies **close** to \mathcal{V} , the optimal embedding in that space is given by projecting onto that space.

$$\mathbf{XV}\mathbf{V}^T = \arg \min_{\mathbf{B} \text{ with rows in } \mathcal{V}} \|\mathbf{X} - \mathbf{B}\|_F^2.$$

Last Class: Low-Rank Approximation

- When data lies in a k -dimensional subspace \mathcal{V} , we can perfectly embed into k dimensions using an orthonormal span $\mathbf{V} \in \mathbb{R}^{d \times k}$.
- When data lies **close** to \mathcal{V} , the optimal embedding in that space is given by projecting onto that space.


$$\mathbf{X}\mathbf{V}\mathbf{V}^T = \arg \min_{\mathbf{B} \text{ with rows in } \mathcal{V}} \|\mathbf{X} - \mathbf{B}\|_F^2.$$

This Class:

- The best subspace \mathcal{V} is the subspace spanned by the top k eigenvectors of $\mathbf{X}^T\mathbf{X}$. How good is this approximation?

RECAP: BASIC SET UP

Reminder of Set Up: Assume that $\vec{x}_1, \dots, \vec{x}_n$ lie **close to** any k -dimensional subspace \mathcal{V} of \mathbb{R}^d . Let $\mathbf{X} \in \mathbb{R}^{n \times d}$ be the data matrix.




Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns.

- $\mathbf{V}\mathbf{V}^T \in \mathbb{R}^{d \times d}$ is the **projection matrix** onto \mathcal{V} .
- $\mathbf{X} \approx \mathbf{X}(\mathbf{V}\mathbf{V}^T)$. Gives the closest approximation to \mathbf{X} with rows in \mathcal{V} .

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

RECAP: BASIC SET UP

Reminder of Set Up: Assume that $\vec{x}_1, \dots, \vec{x}_n$ lie **close to** any k -dimensional subspace \mathcal{V} of \mathbb{R}^d . Let $\mathbf{X} \in \mathbb{R}^{n \times d}$ be the data matrix.



Let $\vec{v}_1, \dots, \vec{v}_k$ be an orthonormal basis for \mathcal{V} and $\mathbf{V} \in \mathbb{R}^{d \times k}$ be the matrix with these vectors as its columns.

- $\mathbf{V}\mathbf{V}^T \in \mathbb{R}^{d \times d}$ is the **projection matrix** onto \mathcal{V} .
- $\mathbf{X} \approx \mathbf{X}(\mathbf{V}\mathbf{V}^T)$. Gives the closest approximation to \mathbf{X} with rows in \mathcal{V} .

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: orthogonal basis for subspace \mathcal{V} . $\mathbf{V} \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

RECAP: BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as $\mathbf{XV}\mathbf{V}^T$. \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

RECAP: BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as \mathbf{XVV}^T . \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

$$\arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{XVV}^T\|_F^2 = \arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{XV}\|_F^2.$$

RECAP: BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as \mathbf{XVV}^T . \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

$$\arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{XVV}^T\|_F^2 = \arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{XV}\|_F^2.$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ **greedily**.

RECAP: BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as \mathbf{XVV}^T . \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

$$\arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{XVV}^T\|_F^2 = \arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{XV}\|_F^2.$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ **greedily**.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

RECAP: BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as \mathbf{XVV}^T . \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

$$\arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{XVV}^T\|_F^2 = \arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{XV}\|_F^2.$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ **greedily**.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

RECAP: BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as \mathbf{XVV}^T . \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

$$\arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{XVV}^T\|_F^2 = \arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{XV}\|_F^2.$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ greedily.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

...

$$\vec{v}_k = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle=0 \ \forall j < k} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

RECAP: BEST FIT SUBSPACE

If $\vec{x}_1, \dots, \vec{x}_n$ are close to a k -dimensional subspace \mathcal{V} with orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times k}$, the data matrix can be approximated as \mathbf{XVV}^T . \mathbf{XV} gives optimal embedding of \mathbf{X} in \mathcal{V} . How do we find \mathcal{V} (equivalently \mathbf{V})?

$$\arg \min_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{X} - \mathbf{XVV}^T\|_F^2 = \arg \max_{\text{orthonormal } \mathbf{V} \in \mathbb{R}^{d \times k}} \|\mathbf{XV}\|_F^2.$$

Surprisingly, can find the columns of \mathbf{V} , $\vec{v}_1, \dots, \vec{v}_k$ greedily.

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

...

$$\vec{v}_k = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle=0 \ \forall j < k} \vec{v}^T \mathbf{X}^T \mathbf{X} \vec{v}.$$

These are exactly the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$.

Eigenvector: $\vec{x} \in \mathbb{R}^d$ is an eigenvector of a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$ if $\mathbf{A}\vec{x} = \lambda\vec{x}$ for some scalar λ (the eigenvalue corresponding to \vec{x}).

Eigenvector: $\vec{x} \in \mathbb{R}^d$ is an eigenvector of a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$ if $\mathbf{A}\vec{x} = \lambda\vec{x}$ for some scalar λ (the eigenvalue corresponding to \vec{x}).

- That is, \mathbf{A} just 'stretches' x .

REVIEW: EIGENVECTORS AND EIGENDECOMPOSITION

Eigenvector: $\vec{x} \in \mathbb{R}^d$ is an eigenvector of a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$ if $\mathbf{A}\vec{x} = \lambda\vec{x}$ for some scalar λ (the eigenvalue corresponding to \vec{x}).

- That is, \mathbf{A} just 'stretches' x .
- If \mathbf{A} is **symmetric**, can find d orthonormal eigenvectors $\vec{v}_1, \dots, \vec{v}_d$. Let $\mathbf{V} \in \mathbb{R}^{d \times d}$ have these vectors as columns and $\mathbf{\Lambda}$ be the diagonal matrix with the corresponding eigenvalues on the diagonal.

REVIEW: EIGENVECTORS AND EIGENDECOMPOSITION

Eigenvector: $\vec{x} \in \mathbb{R}^d$ is an eigenvector of a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$ if $\mathbf{A}\vec{x} = \lambda\vec{x}$ for some scalar λ (the eigenvalue corresponding to \vec{x}).

- That is, \mathbf{A} just ‘stretches’ x .
- If \mathbf{A} is **symmetric**, can find d orthonormal eigenvectors $\vec{v}_1, \dots, \vec{v}_d$. Let $\mathbf{V} \in \mathbb{R}^{d \times d}$ have these vectors as columns and $\mathbf{\Lambda}$ be the diagonal matrix with the corresponding eigenvalues on the diagonal.

$$\mathbf{AV} = \begin{bmatrix} | & | & | & | \\ \mathbf{A}\vec{v}_1 & \mathbf{A}\vec{v}_2 & \cdots & \mathbf{A}\vec{v}_d \\ | & | & | & | \end{bmatrix}$$

REVIEW: EIGENVECTORS AND EIGENDECOMPOSITION

Eigenvector: $\vec{x} \in \mathbb{R}^d$ is an eigenvector of a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$ if $\mathbf{A}\vec{x} = \lambda\vec{x}$ for some scalar λ (the eigenvalue corresponding to \vec{x}).

- That is, \mathbf{A} just ‘stretches’ \vec{x} .
- If \mathbf{A} is **symmetric**, can find d orthonormal eigenvectors $\vec{v}_1, \dots, \vec{v}_d$. Let $\mathbf{V} \in \mathbb{R}^{d \times d}$ have these vectors as columns and $\mathbf{\Lambda}$ be the diagonal matrix with the corresponding eigenvalues on the diagonal.

$$\mathbf{AV} = \begin{bmatrix} | & | & | & | \\ \mathbf{A}\vec{v}_1 & \mathbf{A}\vec{v}_2 & \cdots & \mathbf{A}\vec{v}_d \\ | & | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | & | \\ \lambda_1\vec{v}_1 & \lambda_2\vec{v}_2 & \cdots & \lambda\vec{v}_d \\ | & | & | & | \end{bmatrix}$$

REVIEW: EIGENVECTORS AND EIGENDECOMPOSITION

Eigenvector: $\vec{x} \in \mathbb{R}^d$ is an eigenvector of a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$ if $\mathbf{A}\vec{x} = \lambda\vec{x}$ for some scalar λ (the eigenvalue corresponding to \vec{x}).

- That is, \mathbf{A} just ‘stretches’ \vec{x} .
- If \mathbf{A} is **symmetric**, can find d orthonormal eigenvectors $\vec{v}_1, \dots, \vec{v}_d$. Let $\mathbf{V} \in \mathbb{R}^{d \times d}$ have these vectors as columns and $\mathbf{\Lambda}$ be the diagonal matrix with the corresponding eigenvalues on the diagonal.

$$\mathbf{AV} = \begin{bmatrix} | & | & | & | \\ \mathbf{A}\vec{v}_1 & \mathbf{A}\vec{v}_2 & \cdots & \mathbf{A}\vec{v}_d \\ | & | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | & | \\ \lambda_1\vec{v}_1 & \lambda_2\vec{v}_2 & \cdots & \lambda\vec{v}_d \\ | & | & | & | \end{bmatrix} = \mathbf{V}\mathbf{\Lambda}$$

REVIEW: EIGENVECTORS AND EIGENDECOMPOSITION


Eigenvector: $\vec{x} \in \mathbb{R}^d$ is an eigenvector of a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$ if $\mathbf{A}\vec{x} = \lambda\vec{x}$ for some scalar λ (the eigenvalue corresponding to \vec{x}).

- That is, \mathbf{A} just 'stretches' x .
- If \mathbf{A} is **symmetric**, can find d orthonormal eigenvectors $\vec{v}_1, \dots, \vec{v}_d$. Let $\mathbf{V} \in \mathbb{R}^{d \times d}$ have these vectors as columns and $\mathbf{\Lambda}$ be the diagonal matrix with the corresponding eigenvalues on the diagonal.

$$\mathbf{AV} = \begin{bmatrix} | & | & | & | \\ \mathbf{A}\vec{v}_1 & \mathbf{A}\vec{v}_2 & \cdots & \mathbf{A}\vec{v}_d \\ | & | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | & | \\ \lambda_1 \vec{v}_1 & \lambda_2 \vec{v}_2 & \cdots & \lambda \vec{v}_d \\ | & | & | & | \end{bmatrix} = \mathbf{V}\mathbf{\Lambda}$$

Yields eigendecomposition: $\mathbf{AVV}^T = \mathbf{A} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^T$.

REVIEW: EIGENVECTORS AND EIGENDECOMPOSITION



Typically order the eigenvectors in decreasing order:

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d$$

Courant-Fischer Principal: For symmetric \mathbf{A} , the eigenvectors are given via the greedy optimization:

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{A} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{A} \vec{v}.$$

...

$$\vec{v}_d = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle=0 \ \forall j < d} \vec{v}^T \mathbf{A} \vec{v}.$$

Courant-Fischer Principal: For symmetric \mathbf{A} , the eigenvectors are given via the greedy optimization:

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{A} \vec{v}.$$

$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{A} \vec{v}.$$

...

$$\vec{v}_d = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle=0 \ \forall j < d} \vec{v}^T \mathbf{A} \vec{v}.$$

- $\vec{v}_j^T \mathbf{A} \vec{v}_j = \lambda_j \cdot \vec{v}_j^T \vec{v}_j = \lambda_j$, the j^{th} largest eigenvalue.

Courant-Fischer Principal: For symmetric \mathbf{A} , the eigenvectors are given via the greedy optimization:

$$\vec{v}_1 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1} \vec{v}^T \mathbf{A} \vec{v}.$$


$$\vec{v}_2 = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_1 \rangle=0} \vec{v}^T \mathbf{A} \vec{v}.$$

...

$$\vec{v}_d = \arg \max_{\vec{v} \text{ with } \|\vec{v}\|_2=1, \langle \vec{v}, \vec{v}_j \rangle=0 \ \forall j < d} \vec{v}^T \mathbf{A} \vec{v}.$$

- $\vec{v}_j^T \mathbf{A} \vec{v}_j = \lambda_j \cdot \vec{v}_j^T \vec{v}_j = \lambda_j$, the j^{th} largest eigenvalue.
- The first k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (corresponding to the largest k eigenvalues) are exactly the directions of greatest variance in \mathbf{X} that we use for low-rank approximation.

LOW-RANK APPROX VIA EIGENDECOMPOSITION



LOW-RANK APPROX VIA EIGENDECOMPOSITION

Upshot: Letting \mathbf{V}_k have columns $\vec{v}_1, \dots, \vec{v}_k$ corresponding to the top k eigenvectors of the covariance matrix $\mathbf{X}^T \mathbf{X}$, \mathbf{V}_k is the orthogonal basis minimizing

$$\|\mathbf{X} - \mathbf{XV}_k\mathbf{V}_k^T\|_F^2,$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROX VIA EIGENDECOMPOSITION

Upshot: Letting \mathbf{V}_k have columns $\vec{v}_1, \dots, \vec{v}_k$ corresponding to the top k eigenvectors of the covariance matrix $\mathbf{X}^T \mathbf{X}$, \mathbf{V}_k is the orthogonal basis minimizing

$$\|\mathbf{X} - \mathbf{XV}_k\mathbf{V}_k^T\|_F^2,$$

This is principal component analysis (PCA).

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROX VIA EIGENDECOMPOSITION

Upshot: Letting \mathbf{V}_k have columns $\vec{v}_1, \dots, \vec{v}_k$ corresponding to the top k eigenvectors of the covariance matrix $\mathbf{X}^T \mathbf{X}$, \mathbf{V}_k is the orthogonal basis minimizing

$$\|\mathbf{X} - \mathbf{XV}_k\mathbf{V}_k^T\|_F^2,$$

This is principal component analysis (PCA). How accurate is this low-rank approximation?

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROX VIA EIGENDECOMPOSITION

Upshot: Letting \mathbf{V}_k have columns $\vec{v}_1, \dots, \vec{v}_k$ corresponding to the top k eigenvectors of the covariance matrix $\mathbf{X}^T \mathbf{X}$, \mathbf{V}_k is the orthogonal basis minimizing

$$\|\mathbf{X} - \mathbf{XV}_k\mathbf{V}_k^T\|_F^2,$$

This is principal component analysis (PCA). How accurate is this low-rank approximation? Can understand via eigenvalues of $\mathbf{X}^T \mathbf{X}$.

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

LOW-RANK APPROX VIA EIGENDECOMPOSITION

Upshot: Letting \mathbf{V}_k have columns $\vec{v}_1, \dots, \vec{v}_k$ corresponding to the top k eigenvectors of the covariance matrix $\mathbf{X}^T \mathbf{X}$, \mathbf{V}_k is the orthogonal basis minimizing

$$\|\mathbf{X} - \mathbf{XV}_k\mathbf{V}_k^T\|_F^2,$$

This is principal component analysis (PCA). **How accurate is this low-rank approximation?** Can understand via eigenvalues of $\mathbf{X}^T \mathbf{X}$.

- In Homework 3 we show matrix form of Pythagorus Theorem:

$$\|\mathbf{X}\|_F^2 = \|\mathbf{X} - \mathbf{XV}_k\mathbf{V}_k^T\|_F^2 + \|\mathbf{XV}_k\mathbf{V}_k^T\|_F^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Let $\vec{v}_1, \dots, \vec{v}_k$ be the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (the top k principal components). Approximation error is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Let $\vec{v}_1, \dots, \vec{v}_k$ be the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (the top k principal components). Approximation error is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \|\mathbf{X}\|_F^2 - \|\mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Let $\vec{v}_1, \dots, \vec{v}_k$ be the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (the top k principal components). Approximation error is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \|\mathbf{X}\|_F^2 - \|\mathbf{X}\mathbf{V}_k\|_F^2$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Let $\vec{v}_1, \dots, \vec{v}_k$ be the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (the top k principal components). Approximation error is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \|\mathbf{X}\|_F^2 - \|\mathbf{X}\mathbf{V}_k\|_F^2$$

- For any matrix \mathbf{A} , $\|\mathbf{A}\|_F^2 = \sum_{i=1}^d \|\vec{a}_i\|_2^2 = \text{tr}(\mathbf{A}^T \mathbf{A}) =$ sum of diagonal entries = sum eigenvalues.

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SPECTRUM ANALYSIS

Let $\vec{v}_1, \dots, \vec{v}_k$ be the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (the top k principal components). Approximation error is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \text{tr}(\mathbf{X}^T \mathbf{X}) - \text{tr}(\mathbf{V}_k^T \mathbf{X}^T \mathbf{X} \mathbf{V}_k)$$

- For any matrix \mathbf{A} , $\|\mathbf{A}\|_F^2 = \sum_{i=1}^d \|\vec{a}_i\|_2^2 = \text{tr}(\mathbf{A}^T \mathbf{A}) = \text{sum of diagonal entries} = \text{sum eigenvalues}$.

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Let $\vec{v}_1, \dots, \vec{v}_k$ be the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (the top k principal components). Approximation error is:

$$\begin{aligned}\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 &= \text{tr}(\mathbf{X}^T \mathbf{X}) - \text{tr}(\mathbf{V}_k^T \mathbf{X}^T \mathbf{X} \mathbf{V}_k) \\ &= \sum_{i=1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) - \sum_{i=1}^k \vec{v}_i^T \mathbf{X}^T \mathbf{X} \vec{v}_i\end{aligned}$$

- For any matrix \mathbf{A} , $\|\mathbf{A}\|_F^2 = \sum_{i=1}^d \|\vec{a}_i\|_2^2 = \text{tr}(\mathbf{A}^T \mathbf{A}) = \text{sum of diagonal entries} = \text{sum eigenvalues}.$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Let $\vec{v}_1, \dots, \vec{v}_k$ be the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (the top k principal components). Approximation error is:

$$\begin{aligned} \|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 &= \text{tr}(\mathbf{X}^T \mathbf{X}) - \text{tr}(\mathbf{V}_k^T \mathbf{X}^T \mathbf{X} \mathbf{V}_k) \\ &= \sum_{i=1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) - \sum_{i=1}^k \vec{v}_i^T \mathbf{X}^T \mathbf{X} \vec{v}_i \\ &= \sum_{i=1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) - \sum_{i=1}^k \lambda_i(\mathbf{X}^T \mathbf{X}) \end{aligned}$$

- For any matrix \mathbf{A} , $\|\mathbf{A}\|_F^2 = \sum_{i=1}^d \|\vec{a}_i\|_2^2 = \text{tr}(\mathbf{A}^T \mathbf{A}) = \text{sum of diagonal entries} = \text{sum eigenvalues}$.

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Let $\vec{v}_1, \dots, \vec{v}_k$ be the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$ (the top k principal components). Approximation error is:

$$\begin{aligned} \|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 &= \text{tr}(\mathbf{X}^T \mathbf{X}) - \text{tr}(\mathbf{V}_k^T \mathbf{X}^T \mathbf{X} \mathbf{V}_k) \\ &= \sum_{i=1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) - \sum_{i=1}^k \vec{v}_i^T \mathbf{X}^T \mathbf{X} \vec{v}_i \\ &= \sum_{i=1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) - \sum_{i=1}^k \lambda_i(\mathbf{X}^T \mathbf{X}) = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T \mathbf{X}) \end{aligned}$$

- For any matrix \mathbf{A} , $\|\mathbf{A}\|_F^2 = \sum_{i=1}^d \|\vec{a}_i\|_2^2 = \text{tr}(\mathbf{A}^T \mathbf{A}) = \text{sum of diagonal entries} = \text{sum eigenvalues}.$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

Claim: The error in approximating \mathbf{X} with the best rank k approximation (projecting onto the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$) is:


$$\|\mathbf{X} - \mathbf{X} \mathbf{V}_k \mathbf{V}_k^T\|_F^2 = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T \mathbf{X})$$

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SPECTRUM ANALYSIS

Claim: The error in approximating \mathbf{X} with the best rank k approximation (projecting onto the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$) is:

$$\|\mathbf{X} - \mathbf{X} \mathbf{V}_k \mathbf{V}_k^T\|_F^2 = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T \mathbf{X})$$



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SPECTRUM ANALYSIS


Claim: The error in approximating \mathbf{X} with the best rank k approximation (projecting onto the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$) is:

$$\|\mathbf{X} - \mathbf{X} \mathbf{V}_k \mathbf{V}_k^T\|_F^2 = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T \mathbf{X})$$

784 dimensional vectors



eigendecomposition




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SPECTRUM ANALYSIS


Claim: The error in approximating \mathbf{X} with the best rank k approximation (projecting onto the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$) is:

$$\|\mathbf{X} - \mathbf{X}\mathbf{V}_k\mathbf{V}_k^T\|_F^2 = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T \mathbf{X})$$

784 dimensional vectors



eigendecomposition




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SPECTRUM ANALYSIS


Claim: The error in approximating \mathbf{X} with the best rank k approximation (projecting onto the top k eigenvectors of $\mathbf{X}^T \mathbf{X}$) is:

$$\|\mathbf{X} - \mathbf{X} \mathbf{V}_k \mathbf{V}_k^T\|_F^2 = \sum_{i=k+1}^d \lambda_i(\mathbf{X}^T \mathbf{X})$$

784 dimensional vectors



eigendecomposition



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SPECTRUM ANALYSIS

Plotting the **spectrum** of the covariance matrix $\mathbf{X}^T \mathbf{X}$ (its eigenvalues) shows how compressible \mathbf{X} is using low-rank approximation (i.e., how close $\vec{x}_1, \dots, \vec{x}_n$ are to a low-dimensional subspace).

$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.


SPECTRUM ANALYSIS

Plotting the **spectrum** of the covariance matrix $\mathbf{X}^T \mathbf{X}$ (its eigenvalues) shows how compressible \mathbf{X} is using low-rank approximation (i.e., how close $\vec{x}_1, \dots, \vec{x}_n$ are to a low-dimensional subspace).

784 dimensional vectors



eigendecomposition




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.


SPECTRUM ANALYSIS

Plotting the **spectrum** of the covariance matrix $\mathbf{X}^T \mathbf{X}$ (its eigenvalues) shows how compressible \mathbf{X} is using low-rank approximation (i.e., how close $\vec{x}_1, \dots, \vec{x}_n$ are to a low-dimensional subspace).

784 dimensional vectors




eigendecomposition



$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SPECTRUM ANALYSIS

Plotting the **spectrum** of the covariance matrix $\mathbf{X}^T \mathbf{X}$ (its eigenvalues) shows how compressible \mathbf{X} is using low-rank approximation (i.e., how close $\vec{x}_1, \dots, \vec{x}_n$ are to a low-dimensional subspace).




$\vec{x}_1, \dots, \vec{x}_n \in \mathbb{R}^d$: data points, $\mathbf{X} \in \mathbb{R}^{n \times d}$: data matrix, $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^d$: top eigenvectors of $\mathbf{X}^T \mathbf{X}$, $\mathbf{V}_k \in \mathbb{R}^{d \times k}$: matrix with columns $\vec{v}_1, \dots, \vec{v}_k$.

SPECTRUM ANALYSIS

784 dimensional vectors



eigendecomposition



Exercise: Show that the eigenvalues of $\mathbf{X}^T \mathbf{X}$ are always positive.

Hint: Use that $\lambda_j = \vec{v}_j^T \mathbf{X}^T \mathbf{X} \vec{v}_j$.

- Many (most) datasets can be approximated via projection onto a low-dimensional subspace.
- Find this subspace via a maximization problem:

$$\max_{\text{orthonormal } \mathbf{V}} \|\mathbf{XV}\|_F^2.$$

- Greedy solution via eigendecomposition of $\mathbf{X}^T\mathbf{X}$.
- Columns of \mathbf{V} are the top eigenvectors of $\mathbf{X}^T\mathbf{X}$.
- Error of best low-rank approximation is determined by the tail of $\mathbf{X}^T\mathbf{X}$'s eigenvalue spectrum.

- Many (most) datasets can be approximated via projection onto a low-dimensional subspace.
- Find this subspace via a maximization problem:

$$\max_{\text{orthonormal } \mathbf{V}} \|\mathbf{XV}\|_F^2.$$

- Greedy solution via eigendecomposition of $\mathbf{X}^T \mathbf{X}$.
- Columns of \mathbf{V} are the top eigenvectors of $\mathbf{X}^T \mathbf{X}$.
- Error of best low-rank approximation is determined by the tail of $\mathbf{X}^T \mathbf{X}$'s eigenvalue spectrum.
- We'll return to the problem how to quickly compute the top eigenvectors of $\mathbf{X}^T \mathbf{X}$.

