




On Nešetřil's Pentagon Problem



Hossein Hajiabolhassan

Department of Mathematical Sciences

Shahid Beheshti University, G.C.


Tehran, Iran

Workshop on Graph Homomorphisms and Related Topics
Nova Louka, Czech Republic

November 23-27, 2009


GRAPHS HOMOMORPHISM

- A **homomorphism** $f : G \longrightarrow H$ from a graph G to a graph H is a map $f : V(G) \longrightarrow V(H)$ such that if $uv \in E(G)$ then $f(u)f(v) \in E(H)$.




GRAPHS HOMOMORPHISM

- A **homomorphism** $f : G \longrightarrow H$ from a graph G to a graph H is a map $f : V(G) \longrightarrow V(H)$ such that if $uv \in E(G)$ then $f(u)f(v) \in E(H)$.



NEŠETŘIL'S PENTAGON PROBLEM

- Let G be a **cubic graph** of **sufficiently large girth**, is it true that $\text{Hom}(G, C_5) \neq \emptyset$?





NEŠETŘIL'S PENTAGON PROBLEM

- ▶ If in the **Nešetřil's Pentagon problem** C_5 is replaced by C_3 , then the answer is **affirmative**; and in fact it is a quick consequence of **Brooks' theorem**.
- ▶ (A. Kostochka, J. Nešetřil, and P. Smolikova, 2001) If in the problem C_5 is replaced by C_{11} , then the answer is **negative**.
- ▶ (I.M. Wanless and N.C. Wormald, 2001) If in the problem C_5 is replaced by C_9 , then the answer is **negative**.
- ▶ (H. Hatami, 2005) If in the problem C_5 is replaced by C_7 , then the answer is **negative**.



NEŠETŘIL'S PENTAGON PROBLEM

- ▶ If in the **Nešetřil's Pentagon problem** C_5 is replaced by C_3 , then the answer is **affirmative**; and in fact it is a quick consequence of **Brooks' theorem**.
- ▶ (**A. Kostochka, J. Nešetřil, and P. Smolikova, 2001**) If in the problem C_5 is replaced by C_{11} , then the answer is **negative**.
- ▶ (**I.M. Wanless and N.C. Wormald, 2001**) If in the problem C_5 is replaced by C_9 , then the answer is **negative**.
- ▶ (**H. Hatami, 2005**) If in the problem C_5 is replaced by C_7 , then the answer is **negative**.



NEŠETŘIL'S PENTAGON PROBLEM

- ▶ If in the **Nešetřil's Pentagon problem** C_5 is replaced by C_3 , then the answer is **affirmative**; and in fact it is a quick consequence of **Brooks' theorem**.
- ▶ (**A. Kostochka, J. Nešetřil, and P. Smolikova, 2001**) If in the problem C_5 is replaced by C_{11} , then the answer is **negative**.
- ▶ (**I.M. Wanless and N.C. Wormald, 2001**) If in the problem C_5 is replaced by C_9 , then the answer is **negative**.
- ▶ (**H. Hatami, 2005**) If in the problem C_5 is replaced by C_7 , then the answer is **negative**.




NEŠETŘIL'S PENTAGON PROBLEM

- ▶ If in the **Nešetřil's Pentagon problem** C_5 is replaced by C_3 , then the answer is **affirmative**; and in fact it is a quick consequence of **Brooks' theorem**.
- ▶ (**A. Kostochka, J. Nešetřil, and P. Smolikova, 2001**) If in the problem C_5 is replaced by C_{11} , then the answer is **negative**.
- ▶ (**I.M. Wanless and N.C. Wormald, 2001**) If in the problem C_5 is replaced by C_9 , then the answer is **negative**.
- ▶ (**H. Hatami, 2005**) If in the problem C_5 is replaced by C_7 , then the answer is **negative**.



GRAPH POWERS


For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .





GRAPH POWERS


For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .





GRAPH POWERS


For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .



GRAPH POWERS


For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .

Complete Graph
 K_5




GRAPH POWERS

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .




GRAPH POWERS

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .




GRAPH POWERS

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .



GRAPH POWERS

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .






GRAPH POWERS


- **Lemma.** Let G and H be two simple graphs such that $\text{Hom}(G, H) \neq \emptyset$. Then, for any positive integer k , $\text{Hom}(G^k, H^k) \neq \emptyset$.

GRAPH POWERS

- **Lemma.** Let G and H be two simple graphs such that $\text{Hom}(G, H) \neq \emptyset$. Then, for any positive integer k , $\text{Hom}(G^k, H^k) \neq \emptyset$.




?



GRAPH POWERS

- **Lemma.** Let G and H be two simple graphs such that $\text{Hom}(G, H) \neq \emptyset$. Then, for any positive integer k , $\text{Hom}(G^k, H^k) \neq \emptyset$.





NEŠETŘIL'S PENTAGON PROBLEM

- ▶ If G is a cubic graph and $\text{Hom}(G, C_5) \neq \emptyset$. Then,
 $\text{Hom}(G^3, K_5) \neq \emptyset$.
- ▶ Question. (A. Daneshgar and H. Hajiabolhassan) Is it true that for any natural number g_0 , there exists a cubic graph G whose girth is larger than g_0 and $\chi(G^3) \geq 6$?
- ▶ What is the exact value of $\max_{g(G) \geq g} \chi(G^3)$? (maximum is taken over all cubic graphs with girth at least g)



NEŠETŘIL'S PENTAGON PROBLEM


- ▶ If G is a cubic graph and $\text{Hom}(G, C_5) \neq \emptyset$. Then,
 $\text{Hom}(G^3, K_5) \neq \emptyset$.
- ▶ **Question.** (A. Daneshgar and H. Hajiabolhassan) Is it true that for any natural number g_0 , there exists a cubic graph G whose girth is larger than g_0 and $\chi(G^3) \geq 6$?
- ▶ What is the exact value of $\max_{g(G) \geq g} \chi(G^3)$? (maximum is taken over all cubic graphs with girth at least g)




NEŠETŘIL'S PENTAGON PROBLEM

- ▶ If G is a cubic graph and $\text{Hom}(G, C_5) \neq \emptyset$. Then,
 $\text{Hom}(G^3, K_5) \neq \emptyset$.
- ▶ **Question.** (A. Daneshgar and H. Hajiabolhassan) Is it true that for any natural number g_0 , there exists a cubic graph G whose girth is larger than g_0 and $\chi(G^3) \geq 6$?
- ▶ What is the exact value of $\max_{g(G) \geq g} \chi(G^3)$? (maximum is taken over all cubic graphs with girth at least g)


FRACTIONAL POWERS



FRACTIONAL POWERS



FRACTIONAL POWERS





FRACTIONAL POWERS

- ▶ The graph $G^{\frac{1}{t}}$ is said to be the t -subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ▶ Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.
- ▶ For any graph G , set $G^{\frac{t}{s}} := (G^{\frac{1}{s}})^t$.
- ▶ For instance, if $n \geq 3$ is a positive integer, then $K_n^{\frac{7}{3}} = K_{n^2}$.
Moreover, $K_n^{\frac{6r+1}{2r+1}} = K_{rn^2-rn+n}$.
- ▶ A rational number r is called positive odd rational number if numerator and denominator are both positive odd integers.



FRACTIONAL POWERS

- ▶ The graph $G^{\frac{1}{t}}$ is said to be the t -subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ▶ Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.
- ▶ For any graph G , set $G^{\frac{t}{s}} := (G^{\frac{1}{s}})^t$.
- ▶ For instance, if $n \geq 3$ is a positive integer, then $K_n^{\frac{7}{3}} = K_{n^2}$.
Moreover, $K_n^{\frac{6r+1}{2r+1}} = K_{rn^2-rn+n}$.
- ▶ A rational number r is called positive odd rational number if numerator and denominator are both positive odd integers.



FRACTIONAL POWERS

- ▶ The graph $G^{\frac{1}{t}}$ is said to be the t -subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ▶ Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.
- ▶ For any graph G , set $G^{\frac{t}{s}} := (G^{\frac{1}{s}})^t$.
- ▶ For instance, if $n \geq 3$ is a positive integer, then $K_n^{\frac{7}{3}} = K_{n^2}$.
Moreover, $K_n^{\frac{6r+1}{2r+1}} = K_{rn^2-rn+n}$.
- ▶ A rational number r is called positive odd rational number if numerator and denominator are both positive odd integers.



FRACTIONAL POWERS

- ▶ The graph $G^{\frac{1}{t}}$ is said to be the t -subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ▶ Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.
- ▶ For any graph G , set $G^{\frac{t}{s}} := (G^{\frac{1}{s}})^t$.
- ▶ For instance, if $n \geq 3$ is a positive integer, then $K_n^{\frac{7}{3}} = K_{n^2}$.
Moreover, $K_n^{\frac{6r+1}{2r+1}} = K_{rn^2-rn+n}$.
- ▶ A rational number r is called positive odd rational number if numerator and denominator are both positive odd integers.



FRACTIONAL POWERS

- ▶ The graph $G^{\frac{1}{t}}$ is said to be the t -subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ▶ Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.
- ▶ For any graph G , set $G^{\frac{t}{s}} := (G^{\frac{1}{s}})^t$.
- ▶ For instance, if $n \geq 3$ is a positive integer, then $K_n^{\frac{7}{3}} = K_{n^2}$.
 Moreover, $K_n^{\frac{6r+1}{2r+1}} = K_{rn^2-rn+n}$.
- ▶ A rational number r is called positive odd rational number if numerator and denominator are both positive odd integers.



BLOW UP A GRAPH

- ▶ $\omega(G^r)$ stands for the clique number of G^r where r is a positive odd rational number and $r < og(G)$. One can see that if r approaches to $og(G)$, then $\omega(G^r)$ approaches infinity.

BLOW UP A GRAPH

- ▶ $\omega(G^r)$ stands for the **clique number of G^r** where r is a positive odd rational number and $r < og(G)$. One can see that if r **approaches to $og(G)$** , then $\omega(G^r)$ **approaches infinity**.
- ▶ Let G and H be **non-bipartite graphs**. Then there exists a positive odd rational number $r \geq 1$ such that $H \longrightarrow G^r$.






BLOW OUT A GRAPH

- For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .

BLOW OUT A GRAPH

- ▶ For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .
- ▶ Let G and H be non-bipartite graphs. Then there exists a positive odd rational number $r \leq 1$ such that $G^r \longrightarrow H$.





PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani

- ▶ If a, b and s are non-negative integer, then $G^{\frac{(2s+1)a}{(2s+1)b}} \longleftrightarrow G^{\frac{a}{b}}$.
- ▶ Let r be a positive odd rational number and a is a positive odd integer. If $ra < og(G)$, then $(G^r)^a \longleftrightarrow G^{ra}$.
- ▶ Assume that r and s are positive odd rational numbers. If $rs < og(G)$, then $(G^r)^s \longrightarrow G^{rs}$. It does not hold, in general, that $G^{rs} \longrightarrow (G^r)^s$.
- ▶ Let G and H be two graphs where $2s+1 < og(H)$. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.
- ▶ $G < H$ means $G \longrightarrow H$ and $H \not\rightarrow G$.
- ▶ Let r and s be positive odd rational numbers and G be a non-bipartite graph. If $r < s < og(G)$, then $G^r < G^s$.



PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani

- ▶ If a, b and s are non-negative integer, then $G^{\frac{(2s+1)a}{(2s+1)b}} \longleftrightarrow G^{\frac{a}{b}}$.
- ▶ Let r be a positive odd rational number and a is a positive odd integer. If $ra < og(G)$, then $(G^r)^a \longleftrightarrow G^{ra}$.
- ▶ Assume that r and s are positive odd rational numbers. If $rs < og(G)$, then $(G^r)^s \longrightarrow G^{rs}$. It does not hold, in general, that $G^{rs} \longrightarrow (G^r)^s$.
- ▶ Let G and H be two graphs where $2s+1 < og(H)$. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.
- ▶ $G < H$ means $G \longrightarrow H$ and $H \not\rightarrow G$.
- ▶ Let r and s be positive odd rational numbers and G be a non-bipartite graph. If $r < s < og(G)$, then $G^r < G^s$.



PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani

- ▶ If a, b and s are non-negative integer, then $G^{\frac{(2s+1)a}{(2s+1)b}} \longleftrightarrow G^{\frac{a}{b}}$.
- ▶ Let r be a positive odd rational number and a is a positive odd integer. If $ra < og(G)$, then $(G^r)^a \longleftrightarrow G^{ra}$.
- ▶ Assume that r and s are positive odd rational numbers. If $rs < og(G)$, then $(G^r)^s \longrightarrow G^{rs}$. It does not hold, in general, that $G^{rs} \longrightarrow (G^r)^s$.
- ▶ Let G and H be two graphs where $2s+1 < og(H)$. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.
- ▶ $G < H$ means $G \longrightarrow H$ and $H \not\longrightarrow G$.
- ▶ Let r and s be positive odd rational numbers and G be a non-bipartite graph. If $r < s < og(G)$, then $G^r < G^s$.



PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani

- ▶ If a, b and s are non-negative integer, then $G^{\frac{(2s+1)a}{(2s+1)b}} \longleftrightarrow G^{\frac{a}{b}}$.
- ▶ Let r be a positive odd rational number and a is a positive odd integer. If $ra < og(G)$, then $(G^r)^a \longleftrightarrow G^{ra}$.
- ▶ Assume that r and s are positive odd rational numbers. If $rs < og(G)$, then $(G^r)^s \longrightarrow G^{rs}$. It does not hold, in general, that $G^{rs} \longrightarrow (G^r)^s$.
- ▶ Let G and H be two graphs where $2s+1 < og(H)$. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.
- ▶ $G < H$ means $G \longrightarrow H$ and $H \not\longrightarrow G$.
- ▶ Let r and s be positive odd rational numbers and G be a non-bipartite graph. If $r < s < og(G)$, then $G^r < G^s$.



PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani

- ▶ If a, b and s are non-negative integer, then $G^{\frac{(2s+1)a}{(2s+1)b}} \longleftrightarrow G^{\frac{a}{b}}$.
- ▶ Let r be a positive odd rational number and a is a positive odd integer. If $ra < og(G)$, then $(G^r)^a \longleftrightarrow G^{ra}$.
- ▶ Assume that r and s are positive odd rational numbers. If $rs < og(G)$, then $(G^r)^s \longrightarrow G^{rs}$. It does not hold, in general, that $G^{rs} \longrightarrow (G^r)^s$.
- ▶ Let G and H be two graphs where $2s+1 < og(H)$. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.
- ▶ $G < H$ means $G \longrightarrow H$ and $H \not\longrightarrow G$.
- ▶ Let r and s be positive odd rational numbers and G be a non-bipartite graph. If $r < s < og(G)$, then $G^r < G^s$.



PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani

- ▶ If a, b and s are non-negative integer, then $G^{\frac{(2s+1)a}{(2s+1)b}} \longleftrightarrow G^{\frac{a}{b}}$.
- ▶ Let r be a positive odd rational number and a is a positive odd integer. If $ra < og(G)$, then $(G^r)^a \longleftrightarrow G^{ra}$.
- ▶ Assume that r and s are positive odd rational numbers. If $rs < og(G)$, then $(G^r)^s \longrightarrow G^{rs}$. It does not hold, in general, that $G^{rs} \longrightarrow (G^r)^s$.
- ▶ Let G and H be two graphs where $2s+1 < og(H)$. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.
- ▶ $G < H$ means $G \longrightarrow H$ and $H \not\rightarrow G$.
- ▶ Let r and s be positive odd rational numbers and G be a non-bipartite graph. If $r < s < og(G)$, then $G^r < G^s$.



PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani

- ▶ If a, b and s are non-negative integer, then $G^{\frac{(2s+1)a}{(2s+1)b}} \longleftrightarrow G^{\frac{a}{b}}$.
- ▶ Let r be a positive odd rational number and a is a positive odd integer. If $ra < og(G)$, then $(G^r)^a \longleftrightarrow G^{ra}$.
- ▶ Assume that r and s are positive odd rational numbers. If $rs < og(G)$, then $(G^r)^s \longrightarrow G^{rs}$. It does not hold, in general, that $G^{rs} \longrightarrow (G^r)^s$.
- ▶ Let G and H be two graphs where $2s + 1 < og(H)$. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.
- ▶ $G < H$ means $G \longrightarrow H$ and $H \not\longrightarrow G$.
- ▶ Let r and s be positive odd rational numbers and G be a non-bipartite graph. If $r < s < og(G)$, then $G^r < G^s$.



POWER THICKNESS

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < \text{og}(G)\}?$$

- ▶ Let K_n be complete graph with $n \geq 3$ vertices. Then, $\theta(K_n) = 1$.
- ▶ Let n be a positive integer. Then, $\theta(C_{2n+1}) = \frac{2n+1}{3}$.
- ▶ Let G and H be two non-bipartite graphs with $\chi(G) = \chi(H)$. If $G \longrightarrow H$, then $\theta(G) \geq \theta(H)$.



POWER THICKNESS

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < \text{og}(G)\}?$$

- ▶ Let K_n be complete graph with $n \geq 3$ vertices. Then, $\theta(K_n) = 1$.
- ▶ Let n be a positive integer. Then, $\theta(C_{2n+1}) = \frac{2n+1}{3}$.
- ▶ Let G and H be two non-bipartite graphs with $\chi(G) = \chi(H)$. If $G \rightarrow H$, then $\theta(G) \geq \theta(H)$.



POWER THICKNESS

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k+1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < \text{og}(G)\}?$$

- ▶ Let K_n be complete graph with $n \geq 3$ vertices. Then, $\theta(K_n) = 1$.
- ▶ Let n be a positive integer. Then, $\theta(C_{2n+1}) = \frac{2n+1}{3}$.
- ▶ Let G and H be two non-bipartite graphs with $\chi(G) = \chi(H)$. If $G \longrightarrow H$, then $\theta(G) \geq \theta(H)$.



POWER THICKNESS

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < \text{og}(G)\}?$$

- ▶ Let K_n be complete graph with $n \geq 3$ vertices. Then, $\theta(K_n) = 1$.
- ▶ Let n be a positive integer. Then, $\theta(C_{2n+1}) = \frac{2n+1}{3}$.
- ▶ Let G and H be two non-bipartite graphs with $\chi(G) = \chi(H)$. If $G \rightarrow H$, then $\theta(G) \geq \theta(H)$.



POWER THICKNESS

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < \text{og}(G)\}?$$

- ▶ Let K_n be complete graph with $n \geq 3$ vertices. Then, $\theta(K_n) = 1$.
- ▶ Let n be a positive integer. Then, $\theta(C_{2n+1}) = \frac{2n+1}{3}$.
- ▶ Let G and H be two non-bipartite graphs with $\chi(G) = \chi(H)$. If $G \longrightarrow H$, then $\theta(G) \geq \theta(H)$.



ODD CYCLES

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ (Nešetřil's Pentagon Problem) Let G be a cubic graph of sufficiently large girth, is it true that $\Theta(G) \leq \frac{5}{3}$?
- ▶ (Jaeger's Conjecture) Every planar graph with girth at least $4k$ has a homomorphism to C_{2k+1} .
- ▶ (Jaeger's Conjecture) Let P be a planar graph with girth at least $4k$. Then we have $\chi(P^{\frac{2k+1}{3}}) \leq 3$.



ODD CYCLES

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ (Nešetřil's Pentagon Problem) Let G be a cubic graph of sufficiently large girth, is it true that $\Theta(G) \leq \frac{5}{3}$?
- ▶ (Jaeger's Conjecture) Every planar graph with girth at least $4k$ has a homomorphism to C_{2k+1} .
- ▶ (Jaeger's Conjecture) Let P be a planar graph with girth at least $4k$. Then we have $\chi(P^{\frac{2k+1}{3}}) \leq 3$.



ODD CYCLES

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ (Nešetřil's Pentagon Problem) Let G be a cubic graph of sufficiently large girth, is it true that $\Theta(G) \leq \frac{5}{3}$?
- ▶ (Jaeger's Conjecture) Every planar graph with girth at least $4k$ has a homomorphism to C_{2k+1} .
- ▶ (Jaeger's Conjecture) Let P be a planar graph with girth at least $4k$. Then we have $\chi(P^{\frac{2k+1}{3}}) \leq 3$.




ODD CYCLES

- ▶ (H. Hajiabolhassan) Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .
- ▶ (Nešetřil's Pentagon Problem) Let G be a cubic graph of sufficiently large girth, is it true that $\Theta(G) \leq \frac{5}{3}$?
- ▶ (Jaeger's Conjecture) Every planar graph with girth at least $4k$ has a homomorphism to C_{2k+1} .
- ▶ (Jaeger's Conjecture) Let P be a planar graph with girth at least $4k$. Then we have $\chi(P^{\frac{2k+1}{3}}) \leq 3$.

CIRCULAR CHROMATIC NUMBER


- Let C be a circle of (Euclidean) length r . An r -circular coloring of a graph G is a mapping c which assigns to each vertex v of G an open unit length arc $c(v)$ of C , such that for every edge $uv \in E(G)$ we have $c(u) \cap c(v) = \emptyset$. The circular chromatic number of a graph, denoted by $\chi_c(G)$, is defined as, $\chi_c(G) = \inf\{r : G \text{ admits an } r\text{-circular coloring}\}$.





CIRCULAR COMPLETE GRAPHS

- ▶ The circular complete graph $K_{\frac{p}{q}}$ has the vertex set $\{0, 1, \dots, p-1\}$ and the edge set $\{ij : q \leq |i-j| \leq p-q\}$.




- ▶ Circular chromatic number: $\chi_c(G) = \inf\{\frac{p}{q} \mid G \rightarrow K_{\frac{p}{q}}\}$.



CIRCULAR COMPLETE GRAPHS

- ▶ The circular complete graph $K_{\frac{p}{q}}$ has the vertex set $\{0, 1, \dots, p-1\}$ and the edge set $\{ij : q \leq |i-j| \leq p-q\}$.



- ▶ Circular chromatic number: $\chi_c(G) = \inf\{\frac{p}{q} \mid G \rightarrow K_{\frac{p}{q}}\}$.



CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani

- ▶ Let G be a **non-bipartite graph** with chromatic number $\chi(G)$.
 - ▶ If r is a positive odd rational number and $r \leq \frac{\chi(G)}{3(\chi(G)-2)}$, then $\chi(G^r) = 3$.
 - ▶ There exists a positive odd rational number $r > \frac{\chi(G)}{3(\chi(G)-2)}$ for which $\chi(G^r) = 3$ **if and only if** $\chi(G) \neq \chi_c(G)$.
- ▶ $\chi_c(G) = \inf \left\{ \frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0 \right\}$.
- ▶ For any rational number $\frac{p}{q} > 2$, where $q \nmid p$, we have $\theta(K_{\frac{p}{q}}) > 1$.



CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani

- ▶ Let G be a **non-bipartite graph** with chromatic number $\chi(G)$.
 - ▶ If r is a positive odd rational number and $r \leq \frac{\chi(G)}{3(\chi(G)-2)}$, then $\chi(G^r) = 3$.
 - ▶ There exists a positive odd rational number $r > \frac{\chi(G)}{3(\chi(G)-2)}$ for which $\chi(G^r) = 3$ if and only if $\chi(G) \neq \chi_c(G)$.
- ▶ $\chi_c(G) = \inf \left\{ \frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0 \right\}$.
- ▶ For any rational number $\frac{p}{q} > 2$, where $q \nmid p$, we have $\theta(K_{\frac{p}{q}}) > 1$.



CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani

- ▶ Let G be a **non-bipartite graph** with chromatic number $\chi(G)$.
 - ▶ If r is a positive odd rational number and $r \leq \frac{\chi(G)}{3(\chi(G)-2)}$, then $\chi(G^r) = 3$.
 - ▶ There exists a positive odd rational number $r > \frac{\chi(G)}{3(\chi(G)-2)}$ for which $\chi(G^r) = 3$ **if and only if** $\chi(G) \neq \chi_c(G)$.
- ▶ $\chi_c(G) = \inf \left\{ \frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0 \right\}$.
- ▶ For any rational number $\frac{p}{q} > 2$, where $q \nmid p$, we have $\theta(K_{\frac{p}{q}}) > 1$.



CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani

- ▶ Let G be a **non-bipartite graph** with chromatic number $\chi(G)$.
 - ▶ If r is a positive odd rational number and $r \leq \frac{\chi(G)}{3(\chi(G)-2)}$, then $\chi(G^r) = 3$.
 - ▶ There exists a positive odd rational number $r > \frac{\chi(G)}{3(\chi(G)-2)}$ for which $\chi(G^r) = 3$ **if and only if** $\chi(G) \neq \chi_c(G)$.
- ▶ $\chi_c(G) = \inf\{\frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0\}$.
- ▶ For any rational number $\frac{p}{q} > 2$, where $q \nmid p$, we have $\theta(K_{\frac{p}{q}}) > 1$.



CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani

- ▶ Let G be a **non-bipartite graph** with chromatic number $\chi(G)$.
 - ▶ If r is a positive odd rational number and $r \leq \frac{\chi(G)}{3(\chi(G)-2)}$, then $\chi(G^r) = 3$.
 - ▶ There exists a positive odd rational number $r > \frac{\chi(G)}{3(\chi(G)-2)}$ for which $\chi(G^r) = 3$ **if and only if** $\chi(G) \neq \chi_c(G)$.
- ▶ $\chi_c(G) = \inf\{\frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0\}$.
- ▶ For any rational number $\frac{p}{q} > 2$, **where** $q \nmid p$, we have $\theta(K_{\frac{p}{q}}) > 1$.



SUBDIVISION AND CIRCULAR COLORINGS

- ▶ For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .
- ▶ Let G be a non-bipartite graph.

$$\triangleright f(G, 2t+1) := \max\{2n+1 \mid G^{\frac{1}{2n+1}} \rightarrow C_{2n+1}\}.$$

$$\triangleright (\text{H. H. and A. Taherkhani}) \quad f(G, 2t+1) = 2 \left\lfloor \frac{1+t\chi_0(G)}{\chi_0(G)-2} \right\rfloor + 1.$$

- ▶ (A. Daneshgar and H. H.) If $\sigma \in \text{Hom}(G, C_{2n+1})$, then,

$$\lambda_m^G \geq \frac{|E(G)|}{|V(G)|} \lambda_{2n+1}^{C_{2n+1}},$$

where λ_m^G and $\lambda_{2n+1}^{C_{2n+1}}$ stand for the largest eigenvalues of Laplacian matrices of G and C_{2n+1} , respectively.



SUBDIVISION AND CIRCULAR COLORINGS

- ▶ For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .
- ▶ Let G be a **non-bipartite graph**.

$$\text{▶ } f(G, 2t+1) := \max\{2n+1 \mid G^{\frac{1}{2t+1}} \longrightarrow C_{2n+1}\}.$$

$$\text{▶ (H. H. and A. Taherkhani) } f(G, 2t+1) = 2 \lfloor \frac{1+t\chi_c(G)}{\chi_c(G)-2} \rfloor + 1.$$

- ▶ (A. Daneshgar and H. H.) If $\sigma \in \text{Hom}(G, C_{2n+1})$, then,

$$\lambda_m^G \geq \frac{|E(G)|}{|V(G)|} \lambda_{2n+1}^{C_{2n+1}},$$

where λ_m^G and $\lambda_{2n+1}^{C_{2n+1}}$ stand for the largest eigenvalues of Laplacian matrices of G and C_{2n+1} , respectively.



SUBDIVISION AND CIRCULAR COLORINGS

- ▶ For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .
- ▶ Let G be a **non-bipartite graph**.
 - ▶ $f(G, 2t+1) := \max\{2n+1 \mid G^{\frac{1}{2t+1}} \longrightarrow C_{2n+1}\}.$
 - ▶ (H. H. and A. Taherkhani) $f(G, 2t+1) = 2 \lfloor \frac{1+t\chi_c(G)}{\chi_c(G)-2} \rfloor + 1.$
 - ▶ (A. Daneshgar and H. H.) If $\sigma \in \text{Hom}(G, C_{2n+1})$, then,

$$\lambda_m^G \geq \frac{|E(G)|}{|V(G)|} \lambda_{2n+1}^{C_{2n+1}},$$

where λ_m^G and $\lambda_{2n+1}^{C_{2n+1}}$ stand for the largest eigenvalues of Laplacian matrices of G and C_{2n+1} , respectively.



SUBDIVISION AND CIRCULAR COLORINGS

- ▶ For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .
- ▶ Let G be a **non-bipartite graph**.
 - ▶ $f(G, 2t+1) := \max\{2n+1 \mid G^{\frac{1}{2t+1}} \longrightarrow C_{2n+1}\}$.
 - ▶ (H. H. and A. Taherkhani) $f(G, 2t+1) = 2 \lfloor \frac{1+t\chi_c(G)}{\chi_c(G)-2} \rfloor + 1$.
- ▶ (A. Daneshgar and H. H.) If $\sigma \in \text{Hom}(G, C_{2n+1})$, then,

$$\lambda_m^G \geq \frac{|E(G)|}{|V(G)|} \lambda_{2n+1}^{C_{2n+1}},$$

where λ_m^G and $\lambda_{2n+1}^{C_{2n+1}}$ stand for the largest eigenvalues of Laplacian matrices of G and C_{2n+1} , respectively.



SUBDIVISION AND CIRCULAR COLORINGS





- ▶ For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .
- ▶ Let G be a **non-bipartite graph**.
 - ▶ $f(G, 2t+1) := \max\{2n+1 \mid G^{\frac{1}{2t+1}} \longrightarrow C_{2n+1}\}.$
 - ▶ (H. H. and A. Taherkhani) $f(G, 2t+1) = 2 \lfloor \frac{1+t\chi_c(G)}{\chi_c(G)-2} \rfloor + 1.$
- ▶ (A. Daneshgar and H. H.) If $\sigma \in \text{Hom}(G, C_{2n+1})$, then,

$$\lambda_m^G \geq \frac{|E(G)|}{|V(G)|} \lambda_{2n+1}^{C_{2n+1}},$$

where λ_m^G and $\lambda_{2n+1}^{C_{2n+1}}$ stand for **the largest eigenvalues** of **Laplacian matrices** of G and C_{2n+1} , respectively.



REFERENCES

-  A. Daneshgar and H. Hajiabolhassan, Graph homomorphisms through random walks.
Journal of Graph Theory, 44 (2003), 15–38.
-  A. Daneshgar and H. Hajiabolhassan, Density and power graphs in graph homomorphism problem.
Discrete Mathematics, 308 (2008), 4027–4030.
-  H. Hajiabolhassan, On colorings of graph powers.
Discrete Mathematics, 309 (2009), 4299–4305.
-  H. Hajiabolhassan and A. Taherkhani, Graph powers and graph homomorphisms.
Manuscript 2008, arXiv: 0808.0362v3.

Thank You!

