



# Graph Colorings via The Borsuk-Ulam Theorem

Hossein Hajabolhassan

Department of Mathematical Sciences

Shahid Beheshti University  
Tehran, Iran

*Workshop On Topological Combinatorics  
Shahid Beheshti University, G.C.  
Tehran, Iran*

Wednesday, October 21, 2009



# BASIC IDEA



*Combinatorial  
object  $G$*



# BASIC IDEA



*Combinatorial  
object  $G$*        $\longrightarrow$     *Topological space  $T(G)$*



# BASIC IDEA



*Combinatorial  
object  $G$*        $\longrightarrow$     *Topological space  $T(G)$*



*Topological  
properties of  $T(G)$*



# BASIC IDEA



*Combinatorial  
object  $G$*        $\longrightarrow$     *Topological space  $T(G)$*



*Combinatorial  
properties of  $G$*        $\longleftarrow$     *Topological  
properties of  $T(G)$*



# SIMPLICIAL MAP

- ▶ Let  $K$  and  $L$  be simplicial complexes, and let  $f$  be a map from vertices of  $K$  to  $L$  such that whenever the vertices  $v_0, \dots, v_t$  span a simplex of  $K$ , the points  $f(v_0), \dots, f(v_t)$ , are vertices of a simplex of  $L$ . We call  $f$  a **simplicial map** and it is easy to see that  $f$  can be extended to a **continuous map**  $\|f\| : \|K\| \longrightarrow \|L\|$ .



## NEIGHBORHOOD COMPLEX

- **Neighborhood complex** of a graph  $G$  is denoted by  $\mathcal{N}(G)$  and has  $V(G)$  as vertices.



# NEIGHBORHOOD COMPLEX

- **Neighborhood complex** of a graph  $G$  is denoted by  $\mathcal{N}(G)$  and has  $V(G)$  as vertices.
- $CN(A) = \{v \in V(G) : av \in E(G) \text{ for all } a \in A\}.$





## NEIGHBORHOOD COMPLEX

- **Neighborhood complex** of a graph  $G$  is denoted by  $\mathcal{N}(G)$  and has  $V(G)$  as vertices.
- $CN(A) = \{v \in V(G) : av \in E(G) \text{ for all } a \in A\}$ .
- $\mathcal{N}(G) = \{A : A \subseteq V(G) \text{ & } CN(A) \neq \emptyset\}$ .

# NEIGHBORHOOD COMPLEX



- **Neighborhood complex** of a graph  $G$  is denoted by  $\mathcal{N}(G)$  and has  $V(G)$  as vertices.
- $CN(A) = \{v \in V(G) : av \in E(G) \text{ for all } a \in A\}$ .
- $\mathcal{N}(G) = \{A : A \subseteq V(G) \text{ & } CN(A) \neq \emptyset\}$ .







# NEIGHBORHOOD COMPLEX

- Neighborhood complex of a graph  $G$  is denoted by  $\mathcal{N}(G)$  and has  $V(G)$  as vertices.
- $CN(A) = \{v \in V(G) : av \in E(G) \text{ for all } a \in A\}$ .
- $\mathcal{N}(G) = \{A : A \subseteq V(G) \text{ & } CN(A) \neq \emptyset\}$ .





# NEIGHBORHOOD COMPLEX

- Neighborhood complex of a graph  $G$  is denoted by  $\mathcal{N}(G)$  and has  $V(G)$  as vertices.
- $CN(A) = \{v \in V(G) : av \in E(G) \text{ for all } a \in A\}$ .
- $\mathcal{N}(G) = \{A : A \subseteq V(G) \text{ & } CN(A) \neq \emptyset\}$ .




## NEIGHBORHOOD COMPLEX


 $G$  $N(G)$ 

# NEIGHBORHOOD COMPLEX



$G$



$N(G)$



## NEIGHBORHOOD COMPLEX

 $G$  $N(G)$ 



# LOVÁSZ COMPLEX

- ▶ Set  $\mathcal{N}_1(G) := sd(\mathcal{N}(G))$ , i.e., the barycentric subdivision of  $\mathcal{N}(G)$ .




# LOVÁSZ COMPLEX

- ▶ Set  $\mathcal{N}_1(G) := sd(\mathcal{N}(G))$ , i.e., the barycentric subdivision of  $\mathcal{N}(G)$ .
- ▶ Consider  $CN : V(\mathcal{N}_1(G)) \rightarrow V(\mathcal{N}_1(G))$ . Note that  $CN$  is a simplicial map, since  $A \subseteq B$  implies  $CN(A) \supseteq CN(B)$ . Is it true that  $CN^2 = id$ ?



## LOVÁSZ COMPLEX


- ▶ Set  $\mathcal{N}_1(G) := sd(\mathcal{N}(G))$ , i.e., the barycentric subdivision of  $\mathcal{N}(G)$ .
- ▶ Consider  $CN : V(\mathcal{N}_1(G)) \rightarrow V(\mathcal{N}_1(G))$ . Note that  $CN$  is a simplicial map, since  $A \subseteq B$  implies  $CN(A) \supseteq CN(B)$ . Is it true that  $CN^2 = id$ ?
- ▶  $A \neq CN^2(A)!$





## LOVÁSZ COMPLEX

- ▶ Set  $\mathcal{N}_1(G) := sd(\mathcal{N}(G))$ , i.e., the barycentric subdivision of  $\mathcal{N}(G)$ .
- ▶ Consider  $CN : V(\mathcal{N}_1(G)) \rightarrow V(\mathcal{N}_1(G))$ . Note that  $CN$  is a simplicial map, since  $A \subseteq B$  implies  $CN(A) \supseteq CN(B)$ . Is it true that  $CN^2 = id$ ?
- ▶  $A \neq CN^2(A)!$




- ▶ One can check that  $CN^3 = CN$ .



# LOVÁSZ COMPLEX

- ▶ Set  $\mathcal{N}_1(G) := sd(\mathcal{N}(G))$ , i.e., the barycentric subdivision of  $\mathcal{N}(G)$ .
- ▶ Consider  $CN : V(\mathcal{N}_1(G)) \rightarrow V(\mathcal{N}_1(G))$ . Note that  $CN$  is a simplicial map, since  $A \subseteq B$  implies  $CN(A) \supseteq CN(B)$ . Is it true that  $CN^2 = id$ ?
- ▶  $A \neq CN^2(A)!$




- ▶ One can check that  $CN^3 = CN$ .
- ▶ Let  $L(G) \subseteq \mathcal{N}_1(G)$  where  $V(L(G)) := \{CN(A) \neq \emptyset : A \subseteq V(G)\}$ .



# LOVÁSZ COMPLEX

- ▶ Set  $\mathcal{N}_1(G) := sd(\mathcal{N}(G))$ , i.e., the barycentric subdivision of  $\mathcal{N}(G)$ .
- ▶ Consider  $CN : V(\mathcal{N}_1(G)) \rightarrow V(\mathcal{N}_1(G))$ . Note that  $CN$  is a simplicial map, since  $A \subseteq B$  implies  $CN(A) \supseteq CN(B)$ . Is it true that  $CN^2 = id$ ?
- ▶  $A \neq CN^2(A)!$



- ▶ One can check that  $CN^3 = CN$ .
- ▶ Let  $L(G) \subseteq \mathcal{N}_1(G)$  where  $V(L(G)) := \{CN(A) \neq \emptyset : A \subseteq V(G)\}$ .
- ▶  $(L(G), CN)$  is a free  $\mathbb{Z}_2$ -space.





## Box COMPLEX

- **Box complex** of a graph  $G$  is denoted by  $B(G)$  and has  $V(G) \times \{1, 2\}$  as vertices.

# BOX COMPLEX

- **Box complex** of a graph  $G$  is denoted by  $B(G)$  and has  $V(G) \times \{1, 2\}$  as vertices.
- 


$$B(G) = \left\{ \begin{array}{l} A \uplus B : \quad A, B \subseteq V(G), \quad A \cap B = \emptyset \\ G[A, B] \quad \text{is complete, } CN(A) \neq \emptyset \text{ & } CN(B) \neq \emptyset \end{array} \right\}$$




## Box COMPLEX

- **Box complex** of a graph  $G$  is denoted by  $B(G)$  and has  $V(G) \times \{1, 2\}$  as vertices.
- 

$$B(G) = \left\{ \begin{array}{l} A \uplus B : \quad A, B \subseteq V(G), \quad A \cap B = \emptyset \\ G[A, B] \quad \text{is complete, } CN(A) \neq \emptyset \text{ & } CN(B) \neq \emptyset \end{array} \right\}$$



$C_5$



$B(C_5)$




## DEFINITION

- A **homomorphism**  $f : G \longrightarrow H$  from a graph  $G$  to a graph  $H$  is a map  $f : V(G) \longrightarrow V(H)$  such that if  $uv \in E(G)$  then  $f(u)f(v) \in E(H)$ .


## DEFINITION

- ▶ A **homomorphism**  $f : G \longrightarrow H$  from a graph  $G$  to a graph  $H$  is a map  $f : V(G) \longrightarrow V(H)$  such that if  $uv \in E(G)$  then  $f(u)f(v) \in E(H)$ .



## DEFINITION

- ▶ A **homomorphism**  $f : G \longrightarrow H$  from a graph  $G$  to a graph  $H$  is a map  $f : V(G) \longrightarrow V(H)$  such that if  $uv \in E(G)$  then  $f(u)f(v) \in E(H)$ .





# GRAPH COLORINGS

- ▶ Chromatic number:

$\chi(G) = \min\{n | \text{there exists a homomorphism } \sigma : G \rightarrow K_n\}.$



# GRAPH COLORINGS

- ▶ Chromatic number:

$\chi(G) = \min\{n | \text{there exists a homomorphism } \sigma : G \rightarrow K_n\}.$

- ▶ Circular chromatic number:

$\chi_c(G) = \inf\{\frac{n}{d} | \text{there exists a homomorphism } \sigma : G \rightarrow K_{\frac{n}{d}}\}.$



# GRAPH COLORINGS

- ▶ Chromatic number:

$\chi(G) = \min\{n | \text{there exists a homomorphism } \sigma : G \rightarrow K_n\}.$

- ▶ Circular chromatic number:

$\chi_c(G) = \inf\{\frac{n}{d} | \text{there exists a homomorphism } \sigma : G \rightarrow K_{\frac{n}{d}}\}.$

- ▶ Fractional chromatic number:

$\chi_f(G) = \inf\{\frac{n}{k} | \text{there exists a homomorphism } \sigma : G \rightarrow KG(n, k)\}.$



# GRAPH COLORINGS

- ▶ Chromatic number:

$\chi(G) = \min\{n | \text{there exists a homomorphism } \sigma : G \rightarrow K_n\}.$

- ▶ Circular chromatic number:

$\chi_c(G) = \inf\{\frac{n}{d} | \text{there exists a homomorphism } \sigma : G \rightarrow K_{\frac{n}{d}}\}.$

- ▶ Fractional chromatic number:

$\chi_f(G) = \inf\{\frac{n}{k} | \text{there exists a homomorphism } \sigma : G \rightarrow KG(n, k)\}.$

- ▶ Local chromatic number:

$\psi(G) = \min\{r | \exists n, \text{ there exists a homomorphism } \sigma : G \rightarrow U(n, r)\}.$



# SIMPLICIAL MAP

- ▶ Let  $K$  and  $L$  be simplicial complexes, and let  $f$  be a map from vertices of  $K$  to  $L$  such that whenever the vertices  $v_0, \dots, v_t$  span a simplex of  $K$ , the points  $f(v_0), \dots, f(v_t)$ , are vertices of a simplex of  $L$ . We call  $f$  a **simplicial map** and it is easy to see that  $f$  can be extended to a **continuous map**  $\|f\| : \|K\| \longrightarrow \|L\|$ .



# SIMPLICIAL MAP

- ▶ Let  $K$  and  $L$  be simplicial complexes, and let  $f$  be a map from vertices of  $K$  to  $L$  such that whenever the vertices  $v_0, \dots, v_t$  span a simplex of  $K$ , the points  $f(v_0), \dots, f(v_t)$ , are vertices of a simplex of  $L$ . We call  $f$  a simplicial map and it is easy to see that  $f$  can be extended to a continuous map  $\|f\| : \|K\| \longrightarrow \|L\|$ .
- ▶ If there exists a homomorphism  $f : G \longrightarrow H$  then there are simplicial maps  $N(f) : \mathcal{N}(G) \longrightarrow \mathcal{N}(H)$  and  $B(f) : B(G) \longrightarrow B(H)$ .



# INDEX AND COINDEX

- $\mathbb{Z}_2$ -index:  $ind(X) = \min\{n \geq 0 : X \xrightarrow{\mathbb{Z}_2} S^n\}$



# INDEX AND COINDEX

- ▶  $\mathbb{Z}_2$ -index:  $ind(X) = \min\{n \geq 0 : X \xrightarrow{\mathbb{Z}_2} S^n\}$
- ▶  $\mathbb{Z}_2$ -coindex:  $coind(X) = \max\{n \geq 0 : S^n \xrightarrow{\mathbb{Z}_2} X\}$



# INDEX AND COINDEX

- ▶  $Z_2$ -index:  $ind(X) = \min\{n \geq 0 : X \xrightarrow{Z_2} S^n\}$
- ▶  $Z_2$ -coindex:  $coind(X) = \max\{n \geq 0 : S^n \xrightarrow{Z_2} X\}$
- ▶  $ind(S^n) = coind(S^n) = n.$



# INDEX AND COINDEX

- ▶  $Z_2$ -index:  $ind(X) = \min\{n \geq 0 : X \xrightarrow{Z_2} S^n\}$
- ▶  $Z_2$ -coindex:  $coind(X) = \max\{n \geq 0 : S^n \xrightarrow{Z_2} X\}$
- ▶  $ind(S^n) = coind(S^n) = n.$
- ▶ If  $G$  is a graph, then  $B(G)$  is a  $Z_2$ -space:  $\nu(A \uplus B) := B \uplus A.$



# INDEX AND COINDEX

- ▶  $Z_2$ -index:  $ind(X) = \min\{n \geq 0 : X \xrightarrow{Z_2} S^n\}$
- ▶  $Z_2$ -coindex:  $coind(X) = \max\{n \geq 0 : S^n \xrightarrow{Z_2} X\}$
- ▶  $ind(S^n) = coind(S^n) = n.$
- ▶ If  $G$  is a graph, then  $B(G)$  is a  $Z_2$ -space:  $\nu(A \uplus B) := B \uplus A.$
- ▶ If  $f : G \longrightarrow H$  is a graph homomorphism, then there is a  $Z_2$ -map  $B(f) : B(G) \longrightarrow B(H).$  In fact,

$$B(f)(A \uplus B) := f(A) \uplus f(B).$$

# INDEX AND COINDEX

- ▶  $Z_2$ -index:  $ind(X) = \min\{n \geq 0 : X \xrightarrow{Z_2} S^n\}$
- ▶  $Z_2$ -coindex:  $coind(X) = \max\{n \geq 0 : S^n \xrightarrow{Z_2} X\}$
- ▶  $ind(S^n) = coind(S^n) = n.$
- ▶ If  $G$  is a graph, then  $B(G)$  is a  $Z_2$ -space:  $\nu(A \uplus B) := B \uplus A.$
- ▶ If  $f : G \longrightarrow H$  is a graph homomorphism, then there is a  $Z_2$ -map  $B(f) : B(G) \longrightarrow B(H).$  In fact,

$$B(f)(A \uplus B) := f(A) \uplus f(B).$$

- ▶

$$\begin{array}{ccc} A \uplus B & \rightarrow & f(A) \uplus f(B) \\ \downarrow & & \downarrow \\ B \uplus A & \rightarrow & f(B) \uplus f(A) \end{array}$$



# CHROMATIC NUMBER

- ▶ Let  $G$  and  $H$  be two graphs &  $\text{ind}(B(G)) > \text{ind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .



# CHROMATIC NUMBER

- ▶ Let  $G$  and  $H$  be two graphs &  $\text{ind}(B(G)) > \text{ind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .
- ▶ Let  $G$  and  $H$  be two graphs &  $\text{coind}(B(G)) > \text{coind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .



# CHROMATIC NUMBER

- ▶ Let  $G$  and  $H$  be two graphs &  $\text{ind}(B(G)) > \text{ind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .
- ▶ Let  $G$  and  $H$  be two graphs &  $\text{coind}(B(G)) > \text{coind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .
- ▶  $B(K_n) \xrightarrow{Z_2} S^{n-2}$  &  $\text{ind}(B(K_n)) = n - 2$ .



# CHROMATIC NUMBER

- ▶ Let  $G$  and  $H$  be two graphs &  $\text{ind}(B(G)) > \text{ind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .
- ▶ Let  $G$  and  $H$  be two graphs &  $\text{coind}(B(G)) > \text{coind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .
- ▶  $B(K_n) \xrightarrow{Z_2} S^{n-2}$  &  $\text{ind}(B(K_n)) = n - 2$ .
- ▶ (Lovasz, 1978) For any graph  $G$ ,  $\chi(G) \geq \text{ind}(B(G)) + 2$ .



# CHROMATIC NUMBER

- ▶ Let  $G$  and  $H$  be two graphs &  $\text{ind}(B(G)) > \text{ind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .
- ▶ Let  $G$  and  $H$  be two graphs &  $\text{coind}(B(G)) > \text{coind}(B(H))$  then there is **no homomorphism** from  $G$  to  $H$ .
- ▶  $B(K_n) \xrightarrow{Z_2} S^{n-2}$  &  $\text{ind}(B(K_n)) = n - 2$ .
- ▶ (Lovasz, 1978) For any graph  $G$ ,  $\chi(G) \geq \text{ind}(B(G)) + 2$ .
- ▶ (Sarkaria) For any graph  $G$ ,  $\chi(G) \geq \text{ind}(\text{susp}(B(G))) + 1$ .



# COLORFUL $K_{l,m}$ THEOREM

- ▶ Let  $G$  be a finite graph with  $t = \text{coind}(\text{susp}(B(G))) + 1$  and let  $c$  be an arbitrary proper coloring of  $G$  by colors 1, 2, etc and let  $A, B \subseteq [t]$  form a **bipartition** of the color set, i.e.,  $A \cup B = [t]$  and  $A \cap B = \emptyset$ . Then there exists a **complete bipartite subgraph**  $K_{l,m}$  of  $G$  with sides  $L, M$  such that  $|L| = l = |A|$ ,  $|M| = m = |B|$ , and  $\{c(v) : v \in L\} = A$ , and  $\{c(v) : v \in M\} = B$ . In particular, this  $K_{l,m}$  is **completely multicolored** at  $c$ .



## REFERENCES

-  L. Lovász, Kneser's conjecture, chromatic number, and homotopy.  
*J. Combin. Theory Ser. A*, 25(3):319–324, 1978.
-  Gábor Simonyi and Gábor Tardos, Local chromatic number, Ky Fan's theorem and circular colorings.  
*Combinatorica*, 26(5):587–626, 2006.
-  Gábor Simonyi and Gábor Tardos, Colorful subgraphs in Kneser-like graphs.  
*European J. Combin.*, 28(8):2188–2200, 2007.
-  Jiří Matoušek, *Using the Borsuk-Ulam theorem*, Universitext. Springer-Verlag, Berlin, 2003.  
Lectures on topological methods in combinatorics and geometry, Written in cooperation with Anders Björner and Günter M. Ziegler.



Thank You!

