

On the Altermatic Number of Graphs

Hossein Hajiabolhassan

Joint Work With Meysam Alishahi

Department of Mathematical Sciences
Shahid Beheshti University
Tehran, Iran

Frontiers in Mathematical Sciences
Institute for Research in Fundamental Sciences


December 30, 2014



CHROMATIC NUMBER

Definition (Chromatic number)


The **chromatic number** $\chi(G)$ of a graph G is the **smallest number** of colors needed to color the vertices of G so that **no two adjacent** vertices share the same color.



CHROMATIC NUMBER

Definition (Chromatic number)


The **chromatic number** $\chi(G)$ of a graph G is the **smallest number** of colors needed to color the vertices of G so that **no two adjacent** vertices share the same color.



CHROMATIC NUMBER

Definition (Chromatic number)

The **chromatic number** $\chi(G)$ of a graph G is the **smallest number** of colors needed to color the vertices of G so that **no two adjacent** vertices share the same color.




Chromatic number

It is **NP-hard** to compute the chromatic number of a graph!

GRAPHS HOMOMORPHISM

Definition (Graph Homomorphism)


A **homomorphism** $f : G \rightarrow H$ from a graph G to a graph H is a map $f : V(G) \rightarrow V(H)$ such that if $uv \in E(G)$ then $f(u)f(v) \in E(H)$. Also, the existence of a homomorphism is indicated by the symbols $G \rightarrow H$. Also, $G \leftrightarrow H$ means that $G \rightarrow H$ and $H \rightarrow G$.



GRAPHS HOMOMORPHISM

Definition (Graph Homomorphism)


A **homomorphism** $f : G \rightarrow H$ from a graph G to a graph H is a map $f : V(G) \rightarrow V(H)$ such that if $uv \in E(G)$ then $f(u)f(v) \in E(H)$. Also, the existence of a homomorphism is indicated by the symbols $G \rightarrow H$. Also, $G \leftrightarrow H$ means that $G \rightarrow H$ and $H \rightarrow G$.



GRAPHS HOMOMORPHISM

Definition (Graph Homomorphism)

A **homomorphism** $f : G \rightarrow H$ from a graph G to a graph H is a map $f : V(G) \rightarrow V(H)$ such that if $uv \in E(G)$ then $f(u)f(v) \in E(H)$. Also, the existence of a homomorphism is indicated by the symbols $G \rightarrow H$. Also, $G \leftrightarrow H$ means that $G \rightarrow H$ and $H \rightarrow G$.




Observation

For any graph G , we have $\chi(G) = \min\{n : G \rightarrow K_n\}$.

KNESER REPRESENTATION

$$V(H) = \text{A ground set} = \{1, 2, \dots, n\}$$

$$E(H) = \{e, f, g, \dots\} \subseteq 2^{V(H)}$$



KNESER REPRESENTATION


For a **hypergraph** H , consider the graph $KG(H)$ whose vertex set is $E(H)$ and whose edge set consists of all **disjoint pairs**. For instance, if

$$V(H) = \{1, 2, 3, 4, 5\},$$

$$E(H) = \binom{[5]}{2},$$

then

$$KG(H) = KG(5, 2) = \text{Petersen Graph}.$$



KNESER REPRESENTATION


For a **hypergraph** H , consider the graph $KG(H)$ whose vertex set is $E(H)$ and whose edge set consists of all **disjoint pairs**. For instance, if

$$V(H) = \{1, 2, 3, 4, 5\},$$


$$E(H) = \{\{1, 2\}, \{3, 4\}, \{1, 5\}, \{2, 3\}, \{4, 5\}\},$$

then


$$KG(H) = C_5.$$




KNESER REPRESENTATIONS OF GRAPHS



KNESER REPRESENTATIONS OF GRAPHS



KNESER REPRESENTATIONS OF GRAPHS



ALTERMATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1 2 3 4 5



ALTERNATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	+	0	—	+

A subsequence of **nonzero elements** is termed an **alternating subsequence** if any **two consecutive terms** in this subsequence are **different**.



ALTERMATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	+	0	—	+
+	0	+	—	+

A subsequence of **nonzero elements** is termed an **alternating subsequence** if any **two consecutive terms** in this subsequence are **different**.



ALTERNATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	0	—	+	0

A subsequence of **nonzero elements** is termed an **alternating subsequence** if any **two consecutive terms** in this subsequence are **different**.



ALTERNATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	0	—	+	0

$\{1, 4\}$

$Alt(H)$ = The number of nonzero elements of a longest alternating subsequence which does not contain a positive OR negative hyperedge of H



ALTERNATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	—	0	0	+

$Alt(H)$ = The number of nonzero elements of a longest alternating subsequence which does not contain a positive OR negative hyperedge of H



ALTERNATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	—	+	—	+
0	+	—	+	—
	{1, 3}		{3, 5}	

$Alt(H)$ = The number of nonzero elements of a longest alternating subsequence which does not contain a positive OR negative hyperedge of H



ALTERNATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	—	0	0	+

$Alt(H)$ = The number of nonzero elements of a longest alternating subsequence which does not contain a positive OR negative hyperedge of H

$$|V(H)| - Alt(H) = 2$$



$$V(H) = [m] = \{1, 2, \dots, m\} \quad \& \quad E(H) = \binom{[m]}{n}$$

$$\text{KG}(H) = \text{KG}(m, n)$$

Theorem (L. Lovász 1978)

For any $m \geq 2n$, we have $\chi(\text{KG}(m, n)) = m - 2n + 2$.



KNESER GRAPH

$$V(H) = [m] = \{1, 2, \dots, m\} \quad \& \quad E(H) = \binom{[m]}{n}$$

$$\text{KG}(H) = \text{KG}(m, n)$$

Theorem (L. Lovász 1978)

For any $m \geq 2n$, we have $\chi(\text{KG}(m, n)) = m - 2n + 2$.

1	2	3	4	...	$2n-3$	$2n-2$	$2n-1$	$2n$...	$m-1$	m
+	-	+	-	...	+	-	0	0	...	0	0



KNESER GRAPH

$$V(H) = [m] = \{1, 2, \dots, m\} \quad \& \quad E(H) = \binom{[m]}{n}$$

$$KG(H) = KG(m, n)$$

Theorem (L. Lovász 1978)

For any $m \geq 2n$, we have $\chi(KG(m, n)) = m - 2n + 2$.

1	2	3	4	...	$2n-3$	$2n-2$	$2n-1$	$2n$...	$m-1$	m
+	-	+	-	...	+	-	0	0	...	0	0

$$Alt(H) = 2n - 2$$



KNESER GRAPH

$$V(H) = [m] = \{1, 2, \dots, m\} \quad \& \quad E(H) = \binom{[m]}{n}$$

$$KG(H) = KG(m, n)$$

Theorem (L. Lovász 1978)

For any $m \geq 2n$, we have $\chi(KG(m, n)) = m - 2n + 2$.

1	2	3	4	...	$2n-3$	$2n-2$	$2n-1$	$2n$...	$m-1$	m
+	-	+	-	...	+	-	0	0	...	0	0

$$|V(H)| - Alt(H) \geq m - 2n + 2$$



STRONG ALTERMATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	−	0	+	−
	{1, 4}		{2, 5}	

$SAlt(H)$ = The number of nonzero elements of a longest alternating subsequence which does not contain a positive AND negative hyperedge of H

$$|V(H)| - SAlt(H) + 1$$



STRONG ALTERMATIC NUMBER

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
+	-	0	+	0

$\{1, 4\}$

$SAlt(H)$ = The number of nonzero elements of a longest alternating subsequence which does not contain a positive AND negative hyperedge of H

$$|V(H)| - SAlt(H) + 1 = 3$$



Definition (Altermatic Number and Strong Altermatic Number)

The **altermatic number** $\zeta(G)$ and the **strong altermatic number** $\zeta_s(G)$ of a graph G are defined, respectively, as follows:

$$\zeta(G) = \max_H \{|V(H)| - \text{Alt}(H) : \text{KG}(H) \longleftrightarrow G\}.$$

$$\zeta_s(G) = \max_H \{|V(H)| - \text{SAlt}(H) + 1 : \text{KG}(H) \longleftrightarrow G\}.$$



ALTERMATIC NUMBER

Definition (Altermatic Number and Strong Altermatic Number)

The **altermatic number** $\zeta(G)$ and the **strong altermatic number** $\zeta_s(G)$ of a graph G are defined, respectively, as follows:

$$\zeta(G) = \max_H \{ |V(H)| - \text{Alt}(H) : \text{KG}(H) \longleftrightarrow G \}.$$

$$\zeta_s(G) = \max_H \{ |V(H)| - \text{SAlt}(H) + 1 : \text{KG}(H) \longleftrightarrow G \}.$$


Theorem (M. Alishahi and H.H., 2013)

For any graph G , we have

$$\chi(G) \geq \zeta(G),$$

$$\chi(G) \geq \zeta_s(G).$$

THE ALTERMATIC NUMBER OF FIVE CYCLE




$$V(H) = \{1, 2, 3, 4, 5\}$$

$$E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

$$|V(H)| - Alt(H) = 2$$



THE ALTERMATIC NUMBER OF FIVE CYCLE




$$V(H) = \{1, 2, 3, 4, 5, a, b, c, d\}$$

$$E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$



THE ALTERMATIC NUMBER OF FIVE CYCLE




$$V(H) = \{1, 2, 3, 4, 5, a, b, c, d\}$$

$$E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1 *a* 3 *b* 5 *c* 2 *d* 4



THE ALTERMATIC NUMBER OF FIVE CYCLE




$$V(H) = \{1, 2, 3, 4, 5, a, b, c, d\}$$

$$E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	<i>a</i>	3	<i>b</i>	5	<i>c</i>	2	<i>d</i>	4
+	-	+	0	0	0	-	+	-
			$\{1, 3\}$	$\{2, 4\}$				



THE ALTERNATING NUMBER OF FIVE CYCLE




$$V(H) = \{1, 2, 3, 4, 5, a, b, c, d\}$$

$$E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1 *a* 3 *b* 5 *c* 2 *d* 4
 0 0



THE ALTERNATING NUMBER OF FIVE CYCLE




$$V(H) = \{1, 2, 3, 4, 5, a, b, c, d\}$$

$$E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

$$\begin{array}{ccccccccc} 1 & a & 3 & b & 5 & c & 2 & d & 4 \\ + & - & 0 & + & - & + & 0 & - & + \\ & & & & \{1, 4\} & & & & \end{array}$$



THE ALTERMATIC NUMBER OF FIVE CYCLE



$$V(H) = \{1, 2, 3, 4, 5, a, b, c, d\}$$


$$E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	a	3	b	5	c	2	d	4
+	-	0	+	-	+	0	-	0

$$|V(H)| - Alt(H) = 3$$



KNESER REPRESENTATION



$$V(H) = \{1, 2, 3, 4, 5, a, b, c, d\}$$

$$E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	<i>a</i>	3		2	<i>d</i>	4
+	-	0	+	-	-	0

$$|V(H)| - \text{Alt}(H) = 3$$


What is the best representation for a graph?
(in sense of alternatic number)



KNESER REPRESENTATION

$$V(H) = \text{A ground set} = \{1, 2, \dots, n\}$$

$$E(H) = \{e, f, g, \dots\} \subseteq 2^{V(H)}$$





KNESER REPRESENTATION

$V(H)$ = The edge set of the graph G

$E(H)$ = Every subgraph of G isomorphic to F

$KG(H) = ?$





KNESER REPRESENTATION

$V(H)$ = The edge set of the graph G

$E(H)$ = Every subgraph of G isomorphic to F


$KG(H)$ = Petersen Graph = $KG(G, F)$



GENERAL KNESER GRAPHS

Definition (General Kneser Graphs)

Let G be a graph and \mathcal{F} be a family of graphs. By $KG(G, \mathcal{F})$, we denote the general Kneser graph whose vertex set is the set of all subgraphs of G isomorphic to some member of \mathcal{F} and in which two vertices are adjacent if the corresponding subgraphs are edge-disjoint.




$$KG(C_5, 2K_2)$$



GENERAL KNESER GRAPHS

Definition (General Kneser Graphs)

Let G be a graph and \mathcal{F} be a family of graphs. By $KG(G, \mathcal{F})$, we denote the general Kneser graph whose vertex set is the set of all subgraphs of G isomorphic to some member of \mathcal{F} and in which two vertices are adjacent if the corresponding subgraphs are edge-disjoint.



$$KG(C_5, 2K_2) = C_5$$



A REPRESENTATION FOR SOME GRAPHS

Kneser Graphs

- 1 **Kneser Graphs:** $KG(nK_2, rK_2)$, where nK_2 is a matching of size n .
- 2 $\chi(KG(nK_2, rK_2)) = n - 2r + 2$ (L. Lovász, 1978)



A REPRESENTATION FOR SOME GRAPHS

Kneser Graphs

- ① **Kneser Graphs:** $KG(nK_2, rK_2)$, where nK_2 is a matching of size n .
- ② $\chi(KG(nK_2, rK_2)) = n - 2r + 2$ (L. Lovász, 1978)

Schrijver Graphs

- ① **Schrijver Graphs:** $KG(C_n, rK_2)$, where C_n is a cycle of size n .
- ② $\chi(KG(C_n, rK_2)) = n - 2r + 2$ (A. Schrijver, 1978)



A REPRESENTATION FOR SOME GRAPHS

Kneser Graphs

- ① **Kneser Graphs:** $KG(nK_2, rK_2)$, where nK_2 is a matching of size n .
- ② $\chi(KG(nK_2, rK_2)) = n - 2r + 2$ (L. Lovász, 1978)

Schrijver Graphs

- ① **Schrijver Graphs:** $KG(C_n, rK_2)$, where C_n is a cycle of size n .
- ② $\chi(KG(C_n, rK_2)) = n - 2r + 2$ (A. Schrijver, 1978)


Circular Complete Graphs and Permutation Graphs

- ① **Circular Complete Graphs:** $KG(C_n, P_d)$; P_d is a path of length d .
- ② **Permutation Graphs:** $KG(K_{m,n}, rK_2)$, where $K_{m,n}$ is a complete bipartite graph.

TURÁN NUMBER

Definition (Generalized Turán Number)

We define the **generalized Turán number** $\text{ex}(G, \mathcal{F})$ as the **largest number** m such that there exists a **spanning subgraph** K of G with m edges which contains **no subgraph** isomorphic to a member of \mathcal{F} .




$$\text{ex}(K_6, K_3) = 9$$



TURÁN NUMBER

Definition (Generalized Turán Number)



We define the **generalized Turán number** $\text{ex}(G, \mathcal{F})$ as the **largest number** m such that there exists a **spanning subgraph** K of G with m edges which contains **no subgraph** isomorphic to a member of \mathcal{F} .



TURÁN NUMBER

Definition (Generalized Turán Number)


We define the **generalized Turán number** $\text{ex}(G, \mathcal{F})$ as the **largest number** m such that there exists a **spanning subgraph** K of G with m edges which contains **no subgraph** isomorphic to a member of \mathcal{F} .



UPPER BOUND FOR CHROMATIC NUMBER

Observation

Let G be a graph and \mathcal{F} be a family of graphs. For the general Kneser graph $KG(G, \mathcal{F})$, we have $\chi(KG(G, \mathcal{F})) \leq |E(G)| - \text{ex}(G, \mathcal{F})$.



Proof

Let K has no \mathcal{F} -subgraph and $|E(K)| = \text{ex}(G, \mathcal{F})$. Consider an ordering for $E(G) \setminus E(K)$. Define $c : V(KG(G, \mathcal{F})) \rightarrow \{\text{Colors}\} = E(G) \setminus E(K)$ as follows. Set $c(F)$ to be the smallest edge of F in $E(G) \setminus E(K)$.

LOWER BOUND FOR CHROMATIC NUMBER

Question

Let G be a graph and \mathcal{F} be a family of graphs. What is the best lower bound for the chromatic number of the general Kneser graph $KG(G, \mathcal{F})$?



LOWER BOUND FOR CHROMATIC NUMBER

Question

Let G be a graph and \mathcal{F} be a family of graphs. What is the best lower bound for the chromatic number of the general Kneser graph $\text{KG}(G, \mathcal{F})$?

Observation

$$|E(G)| - 2\text{ex}(G, \mathcal{F}) \leq \chi(\text{KG}(G, \mathcal{F})) \leq |E(G)| - \text{ex}(G, \mathcal{F}).$$

Proof

Set \mathcal{F}' to be all subgraphs of G with exactly $n = \text{ex}(G, \mathcal{F}) + 1$ edges. Consider a graph homomorphism $g : \text{KG}(G, \mathcal{F}') \rightarrow \text{KG}(G, \mathcal{F})$. Let $m = |E(G)|$. One can check that

$$\chi(\text{KG}(G, \mathcal{F}')) = \chi(\text{KG}(m, n)) = |E(G)| - 2\text{ex}(G, \mathcal{F}) \leq \chi(\text{KG}(G, \mathcal{F})).$$


CHROMATIC NUMBER VIA TURÁN NUMBER

Theorem (L. Lovász, 1978)

If $n \geq 2k$, for the Kneser graph $KG(nK_2, kK_2)$, we have

$$\chi(KG(nK_2, kK_2)) = |E(nK_2)| - 2\text{ex}(nK_2, kK_2) = n - 2k + 2.$$



CHROMATIC NUMBER VIA TURÁN NUMBER

Theorem (L. Lovász, 1978)

If $n \geq 2k$, for the Kneser graph $KG(nK_2, kK_2)$, we have

$$\chi(KG(nK_2, kK_2)) = |E(nK_2)| - 2\text{ex}(nK_2, kK_2) = n - 2k + 2.$$

Theorem (A. Schrijver, 1978)

If $n \geq 2k$, for the Schrijver graph $KG(C_n, kK_2)$, we have

$$\chi(KG(C_n, kK_2)) = |E(C_n)| - \text{ex}(C_n, kK_2) = n - 2k + 2.$$



CHROMATIC NUMBER VIA TURÁN NUMBER

Theorem (L. Lovász, 1978)

If $n \geq 2k$, for the Kneser graph $KG(nK_2, kK_2)$, we have

$$\chi(KG(nK_2, kK_2)) = |E(nK_2)| - 2\text{ex}(nK_2, kK_2) = n - 2k + 2.$$

Theorem (A. Schrijver, 1978)

If $n \geq 2k$, for the Schrijver graph $KG(C_n, kK_2)$, we have

$$\chi(KG(C_n, kK_2)) = |E(C_n)| - \text{ex}(C_n, kK_2) = n - 2k + 2.$$

Theorem (P. Frankl, 1985)

For the generalized Kneser graph $KG(K_n, K_k)$, we have

$$\chi(KG(K_n, K_k)) = |E(K_n)| - \text{ex}(K_n, K_k) = (k-1)\binom{s}{2} + rs,$$

where $n = (k-1)s + r$, $0 \leq r < k-1$, and n is sufficiently large.

Problem (G.O.H. Katona and Z. Tuza, 2013)

If q is a prime power and $n = q^2 + q + 1$, does the following equality hold?

$$\chi(\text{KG}(K_n, C_4)) = |E(K_n)| - \text{ex}(K_n, C_4) = \binom{q^2+q+1}{2} - \frac{1}{2}q(q+1)^2$$



CONJECTURES AND PROBLEMS

Problem (G.O.H. Katona and Z. Tuza, 2013)

If q is a prime power and $n = q^2 + q + 1$, does the following equality hold?

$$\chi(\text{KG}(K_n, C_4)) = |E(K_n)| - \text{ex}(K_n, C_4) = \binom{q^2+q+1}{2} - \frac{1}{2}q(q+1)^2$$

Conjecture (G.O.H. Katona and Z. Tuza, 2013)

If k is an odd integer and n is sufficiently large, then

$$\chi(\text{KG}(K_n, C_k)) = |E(K_n)| - \text{ex}(K_n, C_k) = \left\lfloor \frac{(n-1)^2}{4} \right\rfloor.$$



CONJECTURES AND PROBLEMS

Problem (G.O.H. Katona and Z. Tuza, 2013)

If q is a prime power and $n = q^2 + q + 1$, does the following equality hold?

$$\chi(\text{KG}(K_n, C_4)) = |E(K_n)| - \text{ex}(K_n, C_4) = \binom{q^2+q+1}{2} - \frac{1}{2}q(q+1)^2$$

Conjecture (G.O.H. Katona and Z. Tuza, 2013)

If k is an odd integer and n is sufficiently large, then

$$\chi(\text{KG}(K_n, C_k)) = |E(K_n)| - \text{ex}(K_n, C_k) = \lfloor \frac{(n-1)^2}{4} \rfloor.$$

Conjecture (P. Frankl, 1985)


If $k > s \geq 2$, $n \geq 2k - s + 1$, and n is sufficiently large, then

$$\chi(\text{KG}(K_{n,s}, K_{k,s})) = |E(K_{n,s})| - \text{ex}(K_{n,s}, K_{k,s}),$$

where the complete hypergraph $K_{n,s}$ contains all of s -subsets of $[n]$.

ALTERNATING TURÁN NUMBER

Assume that $\sigma = (e_1, e_2, \dots, e_t)$ is an **ordering** of the **edges** of G , where $t = |E(G)|$.




ALTERNATING TURÁN NUMBER

Assume that $\sigma = (e_1, e_2, \dots, e_t)$ is an **ordering** of the **edges of G** , where $t = |E(G)|$.

Definition (Alternating Coloring)

A 2-coloring of a **subset $T \subseteq E(G)$** (with two colors **red** and **blue**) is called an **alternating coloring** (with respect to σ) for T , if we assign two colors **alternatively** to **all members of T** with respect to the ordering σ .



ALTERNATING TURÁN NUMBER

Assume that \mathcal{F} is a family of graphs and G is a graph G .

Definition (Alternating Turán Number)

The **maximum number** of edges of a **spanning subgraph of G** such that there exists an **alternating coloring** for the edges of this subgraph with respect to the ordering σ and also the red subgraph **AND** the blue subgraph has **no subgraph** isomorphic to a member of \mathcal{F} is denoted by $\text{ex}_{alt}(G, \mathcal{F}, \sigma)$. Set

$$\text{ex}_{alt}(G, \mathcal{F}) = \min\{\text{ex}_{alt}(G, \mathcal{F}, \sigma) : \sigma \text{ is an ordering of } E(G)\}.$$





ALTERNATING TURÁN NUMBER

Assume that \mathcal{F} is a family of graphs and G is a graph G .

Definition (Alternating Turán Number)

The **maximum number** of edges of a **spanning subgraph of G** such that there exists an **alternating coloring** for the edges of this subgraph with respect to the ordering σ and also the red subgraph **AND** the blue subgraph has **no subgraph** isomorphic to a member of \mathcal{F} is denoted by $\text{ex}_{\text{alt}}(G, \mathcal{F}, \sigma)$. Set

$$\text{ex}_{\text{alt}}(G, \mathcal{F}) = \min\{\text{ex}_{\text{alt}}(G, \mathcal{F}, \sigma) : \sigma \text{ is an ordering of } E(G)\}.$$



ALTERNATING TURÁN NUMBER

Assume that \mathcal{F} is a family of graphs and G is a graph G .

Definition (Alternating Turán Number)

The **maximum number** of edges of a **spanning subgraph of G** such that there exists an **alternating coloring** for the edges of this subgraph with respect to the ordering σ and also the red subgraph **AND** the blue subgraph has **no subgraph** isomorphic to a member of \mathcal{F} is denoted by $\text{ex}_{alt}(G, \mathcal{F}, \sigma)$. Set

$$\text{ex}_{alt}(G, \mathcal{F}) = \min\{\text{ex}_{alt}(G, \mathcal{F}, \sigma) : \sigma \text{ is an ordering of } E(G)\}.$$

Observation

$$\text{ex}(G, \mathcal{F}) \leq \text{ex}_{alt}(G, \mathcal{F}) \leq 2\text{ex}(G, \mathcal{F})$$



ALTERNATING TURÁN NUMBER

Assume that \mathcal{F} is a family of graphs and G is a graph G .

Definition (Alternating Turán Number)

The **maximum number** of edges of a **spanning subgraph of G** such that there exists an **alternating coloring** for the edges of this subgraph with respect to the ordering σ and also the red subgraph **AND** the blue subgraph has **no subgraph** isomorphic to a member of \mathcal{F} is denoted by $\text{ex}_{alt}(G, \mathcal{F}, \sigma)$. Set

$$\text{ex}_{alt}(G, \mathcal{F}) = \min\{\text{ex}_{alt}(G, \mathcal{F}, \sigma) : \sigma \text{ is an ordering of } E(G)\}.$$

Lemma (M. Alishahi and H.H., 2013)

$$|E(G)| - \text{ex}_{alt}(G, \mathcal{F}) \leq \chi(\text{KG}(G, \mathcal{F})) \leq |E(G)| - \text{ex}(G, \mathcal{F}).$$



ALTERNATING TURÁN NUMBER

Assume that \mathcal{F} is a family of graphs and G is a graph G .

Definition (Alternating Turán Number)

The **maximum number** of edges of a **spanning subgraph of G** such that there exists an **alternating coloring** for the edges of this subgraph with respect to the ordering σ and also the red subgraph **AND** the blue subgraph has **no subgraph** isomorphic to a member of \mathcal{F} is denoted by $\text{ex}_{alt}(G, \mathcal{F}, \sigma)$. Set

$$\text{ex}_{alt}(G, \mathcal{F}) = \min\{\text{ex}_{alt}(G, \mathcal{F}, \sigma) : \sigma \text{ is an ordering of } E(G)\}.$$

Lemma (M. Alishahi and H.H., 2013)

$$|E(G)| - \text{ex}_{alt}(G, \mathcal{F}) \leq \chi(\text{KG}(G, \mathcal{F})) \leq |E(G)| - \text{ex}(G, \mathcal{F}).$$

Corollary (M. Alishahi and H.H., 2013)

If $\text{ex}_{alt}(G, \mathcal{F}) = \text{ex}(G, \mathcal{F})$, then $\chi(\text{KG}(G, \mathcal{F})) = |E(G)| - \text{ex}(G, \mathcal{F})$.

STRONG ALTERNATING TURÁN NUMBER

Assume that \mathcal{F} is a family of graphs and G is a graph G .

Definition (Strong Alternating Turán Number)

The **maximum number** of edges of a **spanning subgraph of G** such that there exists an **alternating coloring** for the edges of this subgraph with respect to the ordering σ and also the red subgraph **OR** the blue subgraph has **no subgraph** isomorphic to a member of \mathcal{F} is denoted by $\text{ex}_{\text{salt}}(G, \mathcal{F}, \sigma)$. Set

$$\text{ex}_{\text{salt}}(G, \mathcal{F}) = \min\{\text{ex}_{\text{salt}}(G, \mathcal{F}, \sigma) : \sigma \text{ is an ordering of } E(G)\}.$$



STRONG ALTERNATING TURÁN NUMBER

Assume that \mathcal{F} is a family of graphs and G is a graph G .

Definition (Strong Alternating Turán Number)

The **maximum number** of edges of a **spanning subgraph of G** such that there exists an **alternating coloring** for the edges of this subgraph with respect to the ordering σ and also the red subgraph **OR** the blue subgraph has **no subgraph** isomorphic to a member of \mathcal{F} is denoted by $\text{ex}_{\text{salt}}(G, \mathcal{F}, \sigma)$. Set

$$\text{ex}_{\text{salt}}(G, \mathcal{F}) = \min\{\text{ex}_{\text{salt}}(G, \mathcal{F}, \sigma) : \sigma \text{ is an ordering of } E(G)\}.$$

Lemma (M. Alishahi and H.H., 2013)

$$|E(G)| - \text{ex}_{\text{salt}}(G, \mathcal{F}) + 1 \leq \chi(\text{KG}(G, \mathcal{F})) \leq |E(G)| - \text{ex}(G, \mathcal{F}).$$

Corollary (M. Alishahi and H.H., 2013)

If $\text{ex}_{\text{salt}}(G, \mathcal{F}) - 1 = \text{ex}(G, \mathcal{F})$, then $\chi(\text{KG}(G, \mathcal{F})) = |E(G)| - \text{ex}(G, \mathcal{F})$.

Observation

$$\chi(KG(nK_2, rK_2)) = n - 2r + 2 = |E(nK_2)| - 2\text{ex}(nK_2, rK_2).$$



Observation

$$\chi(KG(nK_2, rK_2)) = n - 2r + 2 = |E(nK_2)| - 2\text{ex}(nK_2, rK_2).$$

Theorem (M. Alishahi and H.H., 2013-2014)

If G is a sufficiently large **dense graph** or a **sparse connected graph** (with some conditions), then $\chi(KG(G, rK_2)) = |E(G)| - \text{ex}(G, rK_2)$.



MATCHING GRAPHS

Observation

$$\chi(\text{KG}(nK_2, rK_2)) = n - 2r + 2 = |E(nK_2)| - 2\text{ex}(nK_2, rK_2).$$

Theorem (M. Alishahi and H.H., 2013-2014)

If G is a sufficiently large **dense graph** or a **sparse connected graph** (with some conditions), then $\chi(\text{KG}(G, rK_2)) = |E(G)| - \text{ex}(G, rK_2)$.

Proof!

- 1 Present an appropriate **ordering** for $E(G)$.
- 2 In view of **TutteBerge formula**, we show that $\text{ex}_{alt}(G, F) = \text{ex}(G, F)$ or $\text{ex}_{salt}(G, F) - 1 = \text{ex}(G, F)$!





GENERAL KNESER GRAPHS

Theorem (M. Alishahi and H.H., 2013)

If G is a **multigraph** such that the **multiplicity** of each edge is at least 2 and F is a **simple graph**, then $\chi(KG(G, F)) = |E(G)| - \text{ex}(G, F)$.

Proof!

- 1 Present an appropriate **ordering** for $E(G)$.
- 2 $\text{ex}_{alt}(G, F) = \text{ex}(G, F)$




SPANNING TREE GRAPHS

Theorem (M. Alishahi and H.H., 2014)

If G is a **sufficiently large dense graph** and \mathcal{T}_n is the family of the **spanning trees** of G , then $\chi(\text{KG}(G, \mathcal{T}_n)) = |\text{MinimumCUT}(G)|$.

Proof!

- 1 Present an appropriate **ordering** for $E(G)$.
- 2 $\text{ex}_{alt}(G, \mathcal{T}_n) = \text{ex}(G, \mathcal{T}_n)$



$$|\text{MinimumCUT}(G)| = 1$$



THE ALTERMATIC NUMBER OF SPARSE GRAPHS

Theorem (M. Alishahi and H.H., 2013)

For any graph G , we have $\zeta(G) \leq \max\{n : K_{\lceil \frac{n}{2} \rceil, \lfloor \frac{n}{2} \rfloor} \text{ is a subgraph of } G\}$.



THE ALTERMATIC NUMBER OF SPARSE GRAPHS

Theorem (M. Alishahi and H.H., 2013)

For any graph G , we have $\zeta(G) \leq \max\{n : K_{\lceil \frac{n}{2} \rceil, \lfloor \frac{n}{2} \rfloor} \text{ is a subgraph of } G\}$.

Question

Is it true that for any graph G , we have $\zeta(G \vee K_n) \leq \zeta(G) + n$?



THE ALTERMATIC NUMBER OF SPARSE GRAPHS

Theorem (M. Alishahi and H.H., 2013)


For any graph G , we have $\zeta(G) \leq \max\{n : K_{\lceil \frac{n}{2} \rceil, \lfloor \frac{n}{2} \rfloor} \text{ is a subgraph of } G\}$.

Question

Is it true that for any graph G , we have $\zeta(G \vee K_n) \leq \zeta(G) + n$?

Theorem (M. Alishahi and H.H., 2014)

For any graph G , we have $\zeta(M(G)) \geq \zeta(G) + 1$.



TUCKER'S LEMMA

Definition

Let $X = (x_1, \dots, x_n), Y = (y_1, \dots, y_n) \in \{-1, 0, +1\}^n$. Set $X^+ = \{i \in [n] : x_i = +1\}$ and $X^- = \{i \in [n] : x_i = -1\}$. By $X \preceq Y$, we mean $X^+ \subseteq Y^+$ and $X^- \subseteq Y^-$.



TUCKER'S LEMMA

Definition

Let $X = (x_1, \dots, x_n)$, $Y = (y_1, \dots, y_n) \in \{-1, 0, +1\}^n$. Set $X^+ = \{i \in [n] : x_i = +1\}$ and $X^- = \{i \in [n] : x_i = -1\}$. By $X \preceq Y$, we mean $X^+ \subseteq Y^+$ and $X^- \subseteq Y^-$.

Tucker's Lemma, 1946

Let $\lambda : \{-1, 0, +1\}^n \setminus \{(0, \dots, 0)\} \rightarrow \{\pm 1, \pm 2, \dots, \pm(n-1)\}$. Also, assume that for any $X \in \{-1, 0, +1\}^n \setminus \{(0, \dots, 0)\}$, we have $\lambda(-X) = -\lambda(X)$. Then there exist two vectors $X, Y \in \{-1, 0, +1\}^n \setminus \{(0, \dots, 0)\}$ such that $X \preceq Y$ and also $\lambda(X) = -\lambda(Y)$.



TUCKER'S LEMMA

Definition

Let $X = (x_1, \dots, x_n)$, $Y = (y_1, \dots, y_n) \in \{-1, 0, +1\}^n$. Set $X^+ = \{i \in [n] : x_i = +1\}$ and $X^- = \{i \in [n] : x_i = -1\}$. By $X \preceq Y$, we mean $X^+ \subseteq Y^+$ and $X^- \subseteq Y^-$.

Tucker's Lemma, 1946

Let $\lambda : \{-1, 0, +1\}^n \setminus \{(0, \dots, 0)\} \rightarrow \{\pm 1, \pm 2, \dots, \pm(n-1)\}$. Also, assume that for any $X \in \{-1, 0, +1\}^n \setminus \{(0, \dots, 0)\}$, we have $\lambda(-X) = -\lambda(X)$. Then there exist two vectors $X, Y \in \{-1, 0, +1\}^n \setminus \{(0, \dots, 0)\}$ such that $X \preceq Y$ and also $\lambda(X) = -\lambda(Y)$.

Z_p -Tucker Lemma (G.M. Ziegler, 2002)





Definition (The Chromatic Number of Hypergraphs)

Let $H = (V(H), E(H))$ be a hypergraph. The hypergraph H is called r -colorable if there exists a map $c : V(H) \rightarrow \{1, 2, \dots, r\}$ such that no hyperedge is monochromatic. The chromatic number $\chi(H)$ of H is the minimum r such that H is r -colorable.




HYPERGRAPH COLORING

Definition (The Chromatic Number of Hypergraphs)

Let $H = (V(H), E(H))$ be a hypergraph. The hypergraph H is called r -colorable if there exists a map $c : V(H) \rightarrow \{1, 2, \dots, r\}$ such that no hyperedge is monochromatic. The chromatic number $\chi(H)$ of H is the minimum r such that H is r -colorable.

$$V(H) = \{A, B, C, D, a, b, c\}$$



$$E(H) = \{\{A, B, c\}, \{A, D, a\}, \{A, C, b\}, \\ \{B, D, b\}, \{B, C, a\}, \{C, D, c\}, \\ \{a, b, c\}\}$$



HYPERGRAPH COLORING

Definition (The Chromatic Number of Hypergraphs)

Let $H = (V(H), E(H))$ be a hypergraph. The hypergraph H is called r -colorable if there exists a map $c : V(H) \rightarrow \{1, 2, \dots, r\}$ such that no hyperedge is monochromatic. The chromatic number $\chi(H)$ of H is the minimum r such that H is r -colorable.



For a **hypergraph** H and positive integer $r \geq 2$, the Kneser hypergraph $KG_r(H)$ is an r -uniform hypergraph whose vertex set is $E(H)$ and whose hyperedge set consists of all **r -tuples of pairwise disjoint hyperedges of H** .



KNESER HYPERGRAPH

For a **hypergraph** H and positive integer $r \geq 2$, the Kneser hypergraph $KG^r(H)$ is an r -uniform hypergraph whose vertex set is $E(H)$ and whose hyperedge set consists of all r -tuples of pairwise disjoint hyperedges of H . For instance, if $r = 2$,

$$V(H) = \{1, 2, 3, 4, 5\},$$

$$E(H) = \binom{[5]}{2},$$

then

$$KG^2(H) = KG(H) = KG(5, 2) = \text{Petersen Graph}.$$



KNESER HYPERGRAPH

For a **hypergraph** H and positive integer $r \geq 2$, the Kneser hypergraph $KG^r(H)$ is an r -uniform hypergraph whose vertex set is $E(H)$ and whose hyperedge set consists of all **r -tuples of pairwise disjoint hyperedges of H** . For instance, if $r = 3$,

$$V(H) = \{1, 2, 3, 4, 5, 6\},$$

$$E(H) = \binom{[6]}{2},$$

then

$$E(KG^3(H)) = \{\{\{1, 2\}, \{3, 4\}, \{5, 6\}\}, \{\{1, 3\}, \{2, 4\}, \{5, 6\}\}, \dots\}.$$



KNESER HYPERGRAPH

For a **hypergraph** H and positive integer $r \geq 2$, the Kneser hypergraph $\text{KG}^r(H)$ is an r -uniform hypergraph whose vertex set is $E(H)$ and whose hyperedge set consists of all r -tuples of pairwise disjoint hyperedges of H .
For

$$V(H) = [m],$$

$$E(H) = \binom{[m]}{n},$$

then denote the **usual Kneser hypergraph** $\text{KG}^r(H)$ by

$$\text{KG}^r(m, n).$$



Observation (P. Erdős, 1976)

For $m \geq rn$ and $r \geq 2$, we have $\chi(KG^r(m, n)) \leq \left\lceil \frac{m-r(n-1)}{r-1} \right\rceil$.

Conjecture (P. Erdős, 1976)

If $m \geq rn$ and $r \geq 2$, then $\chi(KG^r(m, n)) = \left\lceil \frac{m-r(n-1)}{r-1} \right\rceil$.



Observation (P. Erdős, 1976)

For $m \geq rn$ and $r \geq 2$, we have $\chi(KG^r(m, n)) \leq \left\lceil \frac{m-r(n-1)}{r-1} \right\rceil$.

Conjecture (P. Erdős, 1976)

If $m \geq rn$ and $r \geq 2$, then $\chi(KG^r(m, n)) = \left\lceil \frac{m-r(n-1)}{r-1} \right\rceil$.

Theorem (N. Alon, P. Frankl, and L. Lovász, 1986)

For $m \geq rn$ and $r \geq 2$, we have $\chi(KG^r(m, n)) = \left\lceil \frac{m-r(n-1)}{r-1} \right\rceil$.



THE r -ALTERNATION NUMBER OF A HYPERGRAPH

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1 2 3 4 5



THE r -ALTERNATION NUMBER OF A HYPERGRAPH

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
ω	ω	0	ω^2	ω^3

A subsequence of **nonzero elements** is termed an **alternating subsequence** if any **two consecutive terms** in this subsequence are **different**.



THE r -ALTERNATION NUMBER OF A HYPERGRAPH

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
ω	ω^3	0	ω^2	ω

A subsequence of **nonzero elements** is termed an **alternating subsequence** if any **two consecutive terms** in this subsequence are **different**.



THE r -ALTERNATION NUMBER OF A HYPERGRAPH

$$V(H) = \{1, 2, 3, 4, 5\} \quad \& \quad E(H) = \{\{1, 3\}, \{1, 4\}, \{2, 4\}, \{2, 5\}, \{3, 5\}\}$$

1	2	3	4	5
ω	0	ω^2	ω	ω^3
		$\{1, 4\}$		

$Alt_r(H)$ = The number of nonzero elements of a longest alternating alternating subsequence with at most r signs which does not contain a positive OR negative hyperedge of H



THE ALTERMATIC NUMBER OF HYPERGRAPHS

Theorem (M. Alishahi and H.H., 2013)

For any hypergraph H and positive integer $r \geq 2$, we have

$$\chi(KG^r(H)) \geq \left\lceil \frac{|V(H)| - alt_r(H)}{r-1} \right\rceil.$$



THE ALTERMATIC NUMBER OF HYPERGRAPHS

Theorem (M. Alishahi and H.H., 2013)

For any hypergraph H and positive integer $r \geq 2$, we have

$$\chi(KG^r(H)) \geq \left\lceil \frac{|V(H)| - alt_r(H)}{r-1} \right\rceil.$$

Observation

For any positive integer $r \geq 2$ and the hypergraph $H = (V(H), E(H))$, where $V(H) = [m]$, and $E(H) = \binom{[m]}{n}$, we have $alt_r(H) \leq r(n-1)$.



THE ALTERMATIC NUMBER OF HYPERGRAPHS

Theorem (M. Alishahi and H.H., 2013)

For any hypergraph H and positive integer $r \geq 2$, we have

$$\chi(\text{KG}^r(H)) \geq \left\lceil \frac{|V(H)| - \text{alt}_r(H)}{r-1} \right\rceil.$$

Observation

For any positive integer $r \geq 2$ and the hypergraph $H = (V(H), E(H))$, where $V(H) = [m]$, and $E(H) = \binom{[m]}{n}$, we have $\text{alt}_r(H) \leq r(n-1)$.

Consequently,

Theorem (N. Alon, P. Frankl, and L. Lovász, 1986)

For $m \geq rn$ and $r \geq 2$, we have $\chi(\text{KG}^r(m, n)) = \left\lceil \frac{m - r(n-1)}{r-1} \right\rceil$.

Definition (r -Colorability Defect)

Let the r -colorability defect, denoted by $cd_r(H)$, be the minimum size of a subset $X \subseteq V(H)$ such that the hyperedges of H that contain no points of X is r -colorable. Precisely,
$$cd_r(H) = \min\{|X| : (V(H) \setminus X, \{F \in E(H) : F \cap X = \emptyset\}) \text{ is } r\text{-colorable}\}.$$




DOL'NIKOV'S THEOREM


Definition (r -Colorability Defect)

Let the r -colorability defect, denoted by $\text{cd}_r(H)$, be the minimum size of a subset $X \subseteq V(H)$ such that the hyperedges of H that contain no points of X is r -colorable. Precisely,

$\text{cd}_r(H) = \min\{|X| : (V(H) \setminus X, \{F \in E(H) : F \cap X = \emptyset\}) \text{ is } r\text{-colorable}\}.$



$V(H) - X$



X



DOL'NIKOV'S THEOREM

Theorem (V.L. Dol'nikov for $r = 2$ 1988, I. Kříž 1992)

For any hypergraph H and positive integer $r \geq 2$, we have

$$\chi(KG^r(H)) \geq \left\lceil \frac{\text{cd}_r(H)}{r-1} \right\rceil.$$



DOL'NIKOV'S THEOREM

Theorem (V.L. Dol'nikov for $r = 2$ 1988, I. Kříž 1992)

For any hypergraph H and positive integer $r \geq 2$, we have

$$\chi(KG^r(H)) \geq \left\lceil \frac{\text{cd}_r(H)}{r-1} \right\rceil.$$

Observation

For any hypergraph $H = (V(H), E(H))$ and positive integer $r \geq 2$, we have $\text{alt}_r(H) \leq |V(H)| - \text{cd}_r(H)$.



DOL'NIKOV'S THEOREM

Theorem (V.L. Dol'nikov for $r = 2$ 1988, I. Kříž 1992)

For any hypergraph H and positive integer $r \geq 2$, we have

$$\chi(KG^r(H)) \geq \left\lceil \frac{\text{cd}_r(H)}{r-1} \right\rceil.$$





Observation

For any hypergraph $H = (V(H), E(H))$ and positive integer $r \geq 2$, we have $\text{alt}_r(H) \leq |V(H)| - \text{cd}_r(H)$.

Consequently,

Observation

For any hypergraph $H = (V(H), E(H))$ and positive integer $r \geq 2$, we have $\chi(KG^r(H)) \geq \left\lceil \frac{|V(H)| - \text{alt}_r(H)}{r-1} \right\rceil \geq \left\lceil \frac{\text{cd}_r(H)}{r-1} \right\rceil$.

-  M. Alishahi and H. Hajiabolhassan, On the chromatic number of general Kneser hypergraphs. ArXiv e-prints, February 2013.
-  M. Alishahi and H. Hajiabolhassan, Chromatic number via Turán number. ArXiv e-prints, December 2013.
-  M. Alishahi and H. Hajiabolhassan, Hedetniemi's conjecture via alternating chromatic number. ArXiv e-prints, March 2014.
-  M. Alishahi and H. Hajiabolhassan, On chromatic number and minimum cut. ArXiv e-prints, July 2014.



Thank You!

I would like to express my heartfelt gratitude to **Professor Carsten Thomassen** who generously shared his time with me.



QUESTIONS?!

