

Graph Homomorphisms Through Graph Powers

Hossein Hajiabolhassan



Department of Mathematical Sciences
Shahid Beheshti University, G.C.
Tehran, Iran

ICRTGC-2010, Cochin, India.

August 12-15, 2010

Outline


- ① What is the best choice for definition of power of graphs?



3

Outline


- ① What is the best choice for definition of power of graphs?



- ② For any graph G , we introduce graph powers $G^{\frac{2a+1}{2b+1}}$ and $G^{\frac{2a+1}{2b+1}}$.

Outline


- ① What is the best choice for definition of power of graphs?



- ② For any graph G , we introduce graph powers $G^{\frac{2a+1}{2b+1}}$ and $G^{\frac{2a+1}{2b+1}}$.
③ These powers inherit some properties from power in numbers.

Outline

- ① What is the best choice for definition of power of graphs?




- ② For any graph G , we introduce graph powers $G^{\frac{2a+1}{2b+1}}$ and $G^{\frac{2b+1}{2a+1}}$.
- ③ These powers inherit some properties from power in numbers.
- ④ These powers can be considered as **dual** of each other

$$G^{\frac{2a+1}{2b+1}} \longrightarrow H \iff G \longrightarrow H^{\frac{2b+1}{2a+1}}.$$

Outline

- ① What is the best choice for definition of power of graphs?



- ② For any graph G , we introduce graph powers $G^{\frac{2a+1}{2b+1}}$ and $G^{\frac{2b+1}{2a+1}}$.
- ③ These powers inherit some properties from power in numbers.
- ④ These powers can be considered as **dual** of each other


$$G^{\frac{2a+1}{2b+1}} \longrightarrow H \iff G \longrightarrow H^{\frac{2b+1}{2a+1}}.$$

- ⑤ Applications: Graph homomorphisms, Circular coloring, and etc.

GRAPHS HOMOMORPHISM

Graph Homomorphism


A homomorphism $f : G \rightarrow H$ from a graph G to a graph H is a map $f : V(G) \rightarrow V(H)$ such that if $uv \in E(G)$ then $f(u)f(v) \in E(H)$.



GRAPHS COLORINGS

Chromatic number

How many colors does one need to color the vertices of a given graph G , so that if two vertices are connected by an edge, then they get different colors? The chromatic number of a graph G is the smallest number of colors needed to color the vertices of G .



Chromatic Number of Five Cycle is three!


$$\chi(G) = \min\{n \mid \text{There exists a homomorphism } \sigma : G \rightarrow K_n\}$$

NEŠETŘIL'S PENTAGON PROBLEM, 1999

Let G be a cubic graph of sufficiently large girth, is it true that $\text{Hom}(G, C_5) \neq \emptyset$?



Jarik Nešetřil



NEŠETŘIL'S PENTAGON PROBLEM, 1999

R.L. Brooks, 1941

If in the Nešetřil's Pentagon problem C_5 is replaced by C_3 , then the answer is affirmative; and in fact it is a quick consequence of Brooks' theorem.

A. Kostochka, J. Nešetřil, and P. Smolikova, 2001

If in the problem C_5 is replaced by C_{11} , then the answer is negative.

I.M. Wanless and N.C. Wormald, 2001

If in the problem C_5 is replaced by C_9 , then the answer is negative.


H. Hatami, 2005

If in the problem C_5 is replaced by C_7 , then the answer is negative.

GRAPH POWERS

Graph Powers


For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .



GRAPH POWERS

Graph Powers


For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .



GRAPH POWERS

Graph Powers


For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .



GRAPH POWERS

Graph Powers

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .




GRAPH POWERS

Graph Powers

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .

The Petersen graph




GRAPH POWERS

Graph Powers

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .

The Petersen graph




GRAPH POWERS

Graph Powers

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .

The Petersen graph




GRAPH POWERS

Graph Powers

For a graph G , let G^k be the k th power of G , which is obtained on the vertex set $V(G)$, by connecting any two vertices u and v for which there exists a walk of length k between u and v in G .

The Complete
graph K_{-10}



GRAPH POWERS



Observation.

Let G and H be two simple graphs such that $\text{Hom}(G, H) \neq \emptyset$. Then, for any positive integer k , $\text{Hom}(G^k, H^k) \neq \emptyset$.

GRAPH POWERS

Observation.


Let G and H be two simple graphs such that $\text{Hom}(G, H) \neq \emptyset$. Then, for any positive integer k , $\text{Hom}(G^k, H^k) \neq \emptyset$.



GRAPH POWERS

Observation.

Let G and H be two simple graphs such that $\text{Hom}(G, H) \neq \emptyset$. Then, for any positive integer k , $\text{Hom}(G^k, H^k) \neq \emptyset$.



NEŠETŘIL'S PENTAGON PROBLEM, 1999

Observation.

If G is a cubic graph and $\text{Hom}(G, C_5) \neq \emptyset$. Then, $\text{Hom}(G^3, K_5) \neq \emptyset$.

NEŠETŘIL'S PENTAGON PROBLEM, 1999

Observation.

If G is a cubic graph and $\text{Hom}(G, C_5) \neq \emptyset$. Then, $\text{Hom}(G^3, K_5) \neq \emptyset$.

Question. (A. Daneshgar and H. Hajabolhassan, 2008)

Is it true that for any natural number g_0 , there exists a cubic graph G whose girth is larger than g_0 and $\chi(G^3) \geq 6$?

NEŠETŘIL'S PENTAGON PROBLEM, 1999

Observation.

If G is a cubic graph and $\text{Hom}(G, C_5) \neq \emptyset$. Then, $\text{Hom}(G^3, K_5) \neq \emptyset$.


Question. (A. Daneshgar and H. Hajabolhassan, 2008)

Is it true that for any natural number g_0 , there exists a cubic graph G whose girth is larger than g_0 and $\chi(G^3) \geq 6$?

Conjecture. (Naserasl, 2007)


Let \mathcal{P}_{2k+1} be the class of planar graphs of odd-girth at least $2k + 1$. For every $G \in \mathcal{P}_{2k+1}$ we have $\chi(G^{2k-1}) \leq 2^{2k}$.

FRACTIONAL POWERS




FRACTIONAL POWERS

3



FRACTIONAL POWERS

The third
subdivision
of the Petersen
graph



FRACTIONAL POWERS

- ① The graph $G^{\frac{1}{t}}$ is said to be the *t*-subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.

FRACTIONAL POWERS

- ① The graph $G^{\frac{1}{t}}$ is said to be the *t*-subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ② Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.

FRACTIONAL POWERS

- ① The graph $G^{\frac{1}{t}}$ is said to be the *t*-subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ② Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.

Lemma.

Let G and H be two graphs. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.

FRACTIONAL POWERS

- ① The graph $G^{\frac{1}{t}}$ is said to be the t -subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ② Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.

Lemma.

Let G and H be two graphs. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.

Definition.

For any graph G , set $G^{\frac{t}{s}} := (G^{\frac{1}{s}})^t$.

FRACTIONAL POWERS

- ① The graph $G^{\frac{1}{t}}$ is said to be the t -subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ② Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.

Lemma.

Let G and H be two graphs. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.

Definition.

For any graph G , set $G^{\frac{t}{s}} := (G^{\frac{1}{s}})^t$.

- ③ For instance, if $n \geq 3$ is a positive integer, then $K_n^{\frac{7}{3}} = K_{n^2}$.

FRACTIONAL POWERS

- ① The graph $G^{\frac{1}{t}}$ is said to be the t -subdivision of a graph G if G is obtained from G by replacing each edge by a path with exactly $t - 1$ inner vertices.
- ② Note that $G^{\frac{1}{1}}$ is isomorphic to G . Also, $(G^{\frac{1}{t}})^{\frac{1}{s}}$ is isomorphic to $G^{\frac{1}{st}}$.

Lemma.

Let G and H be two graphs. Then, $G^{\frac{1}{2s+1}} \longrightarrow H$ if and only if $G \longrightarrow H^{2s+1}$.

Definition.

For any graph G , set $G^{\frac{t}{s}} := (G^{\frac{1}{s}})^t$.

- ③ For instance, if $n \geq 3$ is a positive integer, then $K_n^{\frac{7}{3}} = K_{n^2}$.
- ④ A rational number r is called positive odd rational number if numerator and denominator are both positive odd integers.

PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani, 2010

Question.

Let a, b and s be positive odd integers. Is it true that $G^{\frac{sa}{sb}} \longleftrightarrow G^{\frac{a}{b}}$?

PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani, 2010

Question.

Let a, b and s be positive odd integers. Is it true that $G^{\frac{s/a}{s/b}} \longleftrightarrow G^{\frac{a}{b}}$?

PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani, 2010

Lemma.

If a, b and s are positive odd integers, then $G^{\frac{sa}{sb}} \longleftrightarrow G^{\frac{a}{b}}$.

PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani, 2010

Lemma.

If a, b and s are positive odd integers, then $G^{\frac{sa}{sb}} \longleftrightarrow G^{\frac{a}{b}}$.

Observation.

Assume that r and s are positive odd rational number. It **does not hold**, in general, that $G^{rs} \longleftrightarrow (G^r)^s$.

PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani, 2010

Lemma.

If a, b and s are positive odd integers, then $G^{\frac{sa}{sb}} \longleftrightarrow G^{\frac{a}{b}}$.

Lemma.

Assume that r and s are positive odd rational number. If $rs < \text{og}(G)$, then $(G^r)^s \longrightarrow G^{rs}$.

PROPERTIES OF FRACTIONAL POWERS

H. Hajiabolhassan and A. Taherkhani, 2010

Lemma.

If a, b and s are positive odd integers, then $G^{\frac{sa}{sb}} \longleftrightarrow G^{\frac{a}{b}}$.

Lemma.

Assume that r and s are positive odd rational number. If $rs < og(G)$, then $(G^r)^s \longrightarrow G^{rs}$.

Theorem.

$G < H$ means $G \longrightarrow H$ and $H \not\rightarrow G$. Let r and s be positive odd rational numbers and G be a non-bipartite graph. If $r < s < og(G)$, then $G^r < G^s$.

BLOW UP A GRAPH

Observation.

$\omega(G^r)$ stands for the clique number of G^r where r is a positive odd rational number and $r < og(G)$. One can see that if r approaches to $og(G)$, then $\omega(G^r)$ approaches infinity.

BLOW UP A GRAPH

Observation.

$\omega(G^r)$ stands for the clique number of G^r where r is a positive odd rational number and $r < og(G)$. One can see that if r approaches to $og(G)$, then $\omega(G^r)$ approaches infinity.

Observation.

Let G and H be non-bipartite graphs. Then there exists a positive odd rational number $r \geq 1$ such that $H \longrightarrow G^r$.



BLOW OUT A GRAPH

Observation.

For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .


BLOW OUT A GRAPH

Observation.

For any positive integer $a \geq 3$. There exists a graph homomorphism from $G^{\frac{1}{a}}$ to C_a .

Observation.

Let G and H be non-bipartite graphs. Then there exists a positive odd rational number $r \leq 1$ such that $G^r \rightarrow H$.



POWER THICKNESS

H. Hajiabolhassan, 2009

Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .

POWER THICKNESS

H. Hajiabolhassan, 2009

Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .

Definition.

Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < og(G)\}?$$

POWER THICKNESS

H. Hajiabolhassan, 2009

Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .

Definition.

Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < og(G)\}?$$

- ① Let K_n be complete graph with $n \geq 3$ vertices. Then, $\theta(K_n) = 1$.

POWER THICKNESS

H. Hajiabolhassan, 2009

Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .

Definition.

Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < og(G)\}?$$

- ① Let K_n be complete graph with $n \geq 3$ vertices. Then, $\theta(K_n) = 1$.
- ② Let n be a positive integer. Then, $\theta(C_{2n+1}) = \frac{2n+1}{3}$.

POWER THICKNESS

H. Hajabolhassan, 2009

Let G be a graph with odd girth at least $2k + 1$. Then, $\chi(G^{\frac{2k+1}{3}}) \leq 3$ if and only if there exists a graph homomorphism from G to C_{2k+1} .

Definition.

Let G be a non-bipartite graph. What is the value of

$$\Theta(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = \chi(G), r < og(G)\}?$$

- ① Let K_n be complete graph with $n \geq 3$ vertices. Then, $\theta(K_n) = 1$.
- ② Let n be a positive integer. Then, $\theta(C_{2n+1}) = \frac{2n+1}{3}$.
- ③ For any rational number $r \geq 1$, there exists a graph G with $\theta(G) = r$.

ODD CYCLES

Nešetřil's Pentagon Problem, 1999

Let G be a cubic graph of sufficiently large girth, is it true that $\chi(G^{\frac{5}{3}}) \leq 3$?

Jaeger's Conjecture, 1981

Every planar graph with girth at least $4k$ has a homomorphism to C_{2k+1} .

Jaeger's Conjecture, 1981

Let P be a planar graph with girth at least $4k$. Then we have $\chi(P^{\frac{2k+1}{3}}) \leq 3$.

Colorful Graphs

H. Hajiabolhassan and A. Taherkhani, 2010

Definition.

Let G be a graph and $\chi(G) = k$. G is called a **colorful graph** if for any proper k -coloring of G , there exists an induced subgraph H of G such that for any $v \in V(H)$, all colors appear in the closed neighborhood of v .

Colorful Graphs

H. Hajiabolhassan and A. Taherkhani, 2010

Definition.

Let G be a graph and $\chi(G) = k$. G is called a **colorful graph** if for any proper k -coloring of G , there exists an induced subgraph H of G such that for any $v \in V(H)$, all colors appear in the closed neighborhood of v .

Theorem.

For any non-bipartite **colorful graph** G , we have $\theta(G) = 1$.

Colorful Graphs

H. Hajiabolhassan and A. Taherkhani, 2010

Definition.

Let G be a graph and $\chi(G) = k$. G is called a **colorful graph** if for any proper k -coloring of G , there exists an induced subgraph H of G such that for any $v \in V(H)$, all colors appear in the closed neighborhood of v .

Theorem.

For any non-bipartite **colorful graph** G , we have $\theta(G) = 1$.

Question.

Is it true that any graph G with $\theta(G) = 1$ is a colorful graph?

Colorful Graphs

H. Hajiabolhassan and A. Taherkhani, 2010

Definition.

Let G be a graph and $\chi(G) = k$. G is called a **colorful graph** if for any proper k -coloring of G , there exists an induced subgraph H of G such that for any $v \in V(H)$, all colors appear in the closed neighborhood of v .

Theorem.

For any non-bipartite **colorful graph** G , we have $\theta(G) = 1$.

Question.

Is it true that any graph G with $\theta(G) = 1$ is a colorful graph?


Question.

Let m and n be positive integers where $m \geq 2n$. Is the Kneser graph $\text{KG}(m, n)$ a **colorful graph**? Is it true that $\theta(\text{KG}(m, n)) = 1$?

CHROMATIC NUMBER

Chromatic number


Let I be a line of length r . An r -coloring of a graph G is a mapping c which assigns to each vertex v of G an open unit length interval $c(v)$ of I , such that for every edge $uv \in E(G)$ we have $c(u) \cap c(v) = \emptyset$. The chromatic number of a graph, denoted by $\chi(G)$, is defined as, $\chi(G) = \inf\{r : G \text{ admits an } r - \text{coloring}\}$.



CIRCULAR CHROMATIC NUMBER

Circular Chromatic number

Let C be a circle of (Euclidean) length r . An r -circular coloring of a graph G is a mapping c which assigns to each vertex v of G an open unit length arc $c(v)$ of C , such that for every edge $uv \in E(G)$ we have $c(u) \cap c(v) = \emptyset$. The circular chromatic number of a graph, denoted by $\chi_c(G)$, is defined as, $\chi_c(G) = \inf\{r : G \text{ admits an } r - \text{circular coloring}\}$.



Andrew Vince




Xuding Zhu

CIRCULAR CHROMATIC NUMBER

Circular Complete Graphs

The circular complete graph $K_{\frac{p}{q}}$ has the vertex set $\{0, 1, \dots, p - 1\}$ and the edge set $\{ij : q \leq |i - j| \leq p - q\}$.



Circular chromatic number: $\chi_c(G) = \inf\left\{\frac{p}{q} \mid G \rightarrow K_{\frac{p}{q}}\right\}$

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Theorem.

Let G be a **non-bipartite graph** with chromatic number $\chi(G)$.

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Theorem.

Let G be a **non-bipartite graph** with chromatic number $\chi(G)$.

① $\chi_c(G) = \inf\left\{\frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0\right\}.$

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Theorem.

Let G be a non-bipartite graph with chromatic number $\chi(G)$.

- ① $\chi_c(G) = \inf\left\{\frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0\right\}$.
- ② If r is a positive odd rational number and $r \leq \frac{\chi(G)}{3(\chi(G)-2)}$, then $\chi(G^r) = 3$.

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Theorem.

Let G be a **non-bipartite graph** with chromatic number $\chi(G)$.

- ① $\chi_c(G) = \inf\left\{\frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0\right\}$.
- ② If r is a positive odd rational number and $r \leq \frac{\chi(G)}{3(\chi(G)-2)}$, then $\chi(G^r) = 3$.
- ③ There exists a positive odd rational number $r > \frac{\chi(G)}{3(\chi(G)-2)}$ for which $\chi(G^r) = 3$ if and only if $\chi(G) \neq \chi_c(G)$.

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Theorem.

Let G be a non-bipartite graph with chromatic number $\chi(G)$.

- ① $\chi_c(G) = \inf\left\{\frac{2n+1}{n-t} \mid \chi(G^{\frac{2n+1}{3(2t+1)}}) = 3, n > t > 0\right\}$.
- ② If r is a positive odd rational number and $r \leq \frac{\chi(G)}{3(\chi(G)-2)}$, then $\chi(G^r) = 3$.
- ③ There exists a positive odd rational number $r > \frac{\chi(G)}{3(\chi(G)-2)}$ for which $\chi(G^r) = 3$ if and only if $\chi(G) \neq \chi_c(G)$.

Theorem.

For any rational number $\frac{p}{q} > 2$, where $q \nmid p$, we have $\theta(K_{\frac{p}{q}}) > 1$.

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Definition.

Let G be a **non-bipartite graph**. Set

$$\mu(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = 3, r < og(G)\}.$$

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Definition.

Let G be a **non-bipartite graph**. Set

$$\mu(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = 3, r < \text{og}(G)\}.$$

Theorem.

Let G be a non-bipartite graph. Then $\mu(G) = \frac{\chi_c(G)}{3(\chi_c(G)-2)}$.

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Definition.

Let G be a **non-bipartite graph**. Set

$$\mu(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = 3, r < \text{og}(G)\}.$$

Theorem.

Let G be a non-bipartite graph. Then $\mu(G) = \frac{\chi_c(G)}{3(\chi_c(G)-2)}$.

Conjecture. (X. Zhu, 1992)

Let G and H be two graphs. Then $\chi_c(G \times H) = \min\{\chi_c(G), \chi_c(H)\}$.

CIRCULAR COLORINGS

H. Hajiabolhassan and A. Taherkhani, 2010

Definition.

Let G be a **non-bipartite graph**. Set

$$\mu(G) := \sup\{r \mid r = \frac{2t+1}{2s+1}, \chi(G^r) = 3, r < \text{og}(G)\}.$$

Theorem.

Let G be a non-bipartite graph. Then $\mu(G) = \frac{\chi_c(G)}{3(\chi_c(G)-2)}$.

Conjecture. (X. Zhu, 1992)

Let G and H be two graphs. Then $\chi_c(G \times H) = \min\{\chi_c(G), \chi_c(H)\}$.

Conjecture.

Let G and H be two graphs. Then $\mu(G \times H) = \max\{\mu(G), \mu(H)\}$.

DUAL POWER

Definition.

For any non-negative integer s , define the graph $G^{\frac{1}{2s+1}}$ as follows.

DUAL POWER

Definition.

For any non-negative integer s , define the graph $G^{\frac{1}{2s+1}}$ as follows.

- ① $V(G^{\frac{1}{2s+1}}) := \{(A_1, \dots, A_{s+1}) \mid A_i \subseteq V(G), |A_1| = 1, \emptyset \neq A_i \subseteq N_{i-1}(A_1), i \leq s+1\}$, where
 $N_i(v) := \{u \mid \text{there is a walk of length } i \text{ joining } u \text{ and } v\}.$

DUAL POWER

Definition.

For any non-negative integer s , define the graph $G^{\frac{1}{2s+1}}$ as follows.

- ① $V(G^{\frac{1}{2s+1}}) := \{(A_1, \dots, A_{s+1}) \mid A_i \subseteq V(G), |A_1| = 1, \emptyset \neq A_i \subseteq N_{i-1}(A_1), i \leq s+1\}$, where
 $N_i(v) := \{u \mid \text{there is a walk of length } i \text{ joining } u \text{ and } v\}.$
- ② $(A_1, \dots, A_{s+1})(B_1, \dots, B_{s+1}) \in E(G^{\frac{1}{2s+1}})$ if for any $1 \leq i \leq s$ and $1 \leq j \leq s+1$, $A_i \subseteq B_{i+1}$, $B_i \subseteq A_{i+1}$, and every vertex of A_j is joined to every vertex of B_j .

DUAL POWER

Definition.

For any non-negative integer s , define the graph $G^{\frac{1}{2s+1}}$ as follows.

- ① $V(G^{\frac{1}{2s+1}}) := \{(A_1, \dots, A_{s+1}) \mid A_i \subseteq V(G), |A_1| = 1, \emptyset \neq A_i \subseteq N_{i-1}(A_1), i \leq s+1\}$, where
 $N_i(v) := \{u \mid \text{there is a walk of length } i \text{ joining } u \text{ and } v\}.$
- ② $(A_1, \dots, A_{s+1})(B_1, \dots, B_{s+1}) \in E(G^{\frac{1}{2s+1}})$ if for any $1 \leq i \leq s$ and $1 \leq j \leq s+1$, $A_i \subseteq B_{i+1}$, $B_i \subseteq A_{i+1}$, and every vertex of A_j is joined to every vertex of B_j .

Definition.

For any non-negative integers r and s , define $G^{\frac{2r+1}{2s+1}} := (G^{\frac{1}{2s+1}})^{2r+1}$.

DUAL POWER

For any positive integer n we have

$$\text{KG}(2n+1, n)^{\frac{1}{2s+1}} \longleftrightarrow \text{KG}(2n+1, n)^{\frac{1}{2s+1}}.$$

DUAL POWER

For any positive integer n we have

$$\text{KG}(2n+1, n)^{\frac{1}{2s+1}} \longleftrightarrow \text{KG}(2n+1, n)^{\frac{1}{2s+1}}.$$

For any $s \geq 0$, we have

$$K_3^{\frac{1}{2s+1}} \simeq C_{6s+3}.$$

DUAL POWER

For any positive integer n we have

$$\text{KG}(2n+1, n)^{\frac{1}{2s+1}} \longleftrightarrow \text{KG}(2n+1, n)^{\frac{1}{2s+1}}.$$

For any $s \geq 0$, we have

$$K_3^{\frac{1}{2s+1}} \simeq C_{6s+3}.$$

For any $r, s, n \geq 0$, we have

$$C_{2n+1}^{\frac{2r+1}{2s+1}} \longleftrightarrow K_{\frac{4ns+2n+2s+1}{2ns+n+s-r}}.$$

HISTORY OF DUAL POWER

In 2004, A. Gyärfás, T. Jensen, and M. Stiebitz defined the graph $K_n^{\frac{1}{3}}$ to show that there is a *n*-chromatic graph with an *n*-coloring where the neighbors of each color-class form an independent set.



A. Gyärfás, T. Jensen, and M. Stiebitz

HISTORY OF DUAL POWER

In 2004, A. Gyärfás, T. Jensen, and M. Stiebitz defined the graph $K_n^{\frac{1}{3}}$ to show that there is a n -chromatic graph with an n -coloring where the neighbors of each color-class form an independent set.



A. Gyärfás, T. Jensen, and M. Stiebitz

In 2005, C. Tardif defined the graph $G^{\frac{1}{3}}$ to show that the circular complete graph $K_{\frac{p}{q}}$, for $\frac{p}{q} < 4$, is a multiplicative graph, i.e., $G \times H \rightarrow K_{\frac{p}{q}} \implies G \rightarrow K_{\frac{p}{q}}$ or $H \rightarrow K_{\frac{p}{q}}$.



Claude Tardif

HISTORY OF DUAL POWER

In 2004, A. Gyärfás, T. Jensen, and M. Stiebitz defined the graph $K_n^{\frac{1}{3}}$ to show that there is a n -chromatic graph with an n -coloring where the neighbors of each color-class form an independent set.



A. Gyärfás, T. Jensen, and M. Stiebitz

In 2005, C. Tardif defined the graph $G^{\frac{1}{3}}$ to show that the circular complete graph $K_{\frac{p}{q}}$, for $\frac{p}{q} < 4$, is a multiplicative graph, i.e., $G \times H \rightarrow K_{\frac{p}{q}} \implies G \rightarrow K_{\frac{p}{q}}$ or $H \rightarrow K_{\frac{p}{q}}$.



Claude Tardif

In 2006, G. Simonyi and G. Tardos showed that $K_n^{\frac{1}{2k+1}}$ is an n -critical graph.



Gabor Simonyi and Gabor Tardos

PROPERTIES OF DUAL POWERS

H. Hajiabolhassan and A. Taherkhani, 2010

Lemma.

- ① If a, b and s are positive odd integers, then $G^{\frac{sa}{sb}} \longleftrightarrow G^{\frac{a}{b}}$.

Lemma.

- ② Assume that a, b, c and d are positive odd integers. If $ac < og(G^{\frac{1}{bd}})$, then $G^{\frac{ac}{bd}} \rightarrow (G^{\frac{a}{b}})^{\frac{c}{d}}$.

Theorem.

- ③ Let a, b, c and d be positive odd integers and G be a non-bipartite graph. If $\frac{a}{b} < \frac{c}{d}$ and $c < og(G^{\frac{1}{d}})$, then $G^{\frac{a}{b}} < G^{\frac{c}{d}}$.

DUAL POWER

Theorem. (H. Hajiabolhassan and A. Taherkhani, 2010)

Let G and H be two graphs. Also, assume that $\frac{2a+1}{2b+1} < og(G)$ and $2b + 1 < og(H^{\frac{1}{2a+1}})$. We have $G^{\frac{2a+1}{2b+1}} \rightarrow H$ if and only if $G \rightarrow H^{\frac{2b+1}{2a+1}}$.

DUAL POWER

Theorem. (H. Hajiabolhassan and A. Taherkhani, 2010)

Let G and H be two graphs. Also, assume that $\frac{2a+1}{2b+1} < og(G)$ and $2b + 1 < og(H^{\frac{1}{2a+1}})$. We have $G^{\frac{2a+1}{2b+1}} \rightarrow H$ if and only if $G \rightarrow H^{\frac{2b+1}{2a+1}}$.

Theorem. (H. Hajiabolhassan, 2009)

Let m, n , and k be positive integers with $m \geq 2n$. Then

$\chi(KG(m, n)^{\frac{1}{2k-1}}) = m - 2n + 2$. Moreover, if m is an even positive integer, then $\chi_c(KG(m, n)^{\frac{1}{2k-1}}) = m - 2n + 2$.

DUAL POWER

Theorem. (H. Hajiabolhassan and A. Taherkhani, 2010)

Let G and H be two graphs. Also, assume that $\frac{2a+1}{2b+1} < og(G)$ and $2b+1 < og(H^{\frac{1}{2a+1}})$. We have $G^{\frac{2a+1}{2b+1}} \rightarrow H$ if and only if $G \rightarrow H^{\frac{2b+1}{2a+1}}$.

Theorem. (H. Hajiabolhassan, 2009)

Let m , n , and k be positive integers with $m \geq 2n$. Then

$\chi(KG(m, n)^{\frac{1}{2k-1}}) = m - 2n + 2$. Moreover, if m is an even positive integer, then $\chi_c(KG(m, n)^{\frac{1}{2k-1}}) = m - 2n + 2$.

Theorem. (H. Hajiabolhassan and A. Taherkhani, 2010)

Let k , l , and m be positive integers where $m \geq 3$ and $\frac{2l-1}{2k-1} \leq 1$. Then
 $\theta(K_m^{\frac{2l-1}{2k-1}}) = \frac{2k-1}{2l-1}$.

REFERENCES

-  A. Daneshgar and H. Hajabolhassan, Density and power graphs in graph homomorphism problem.
Discrete Mathematics, 308 (2008), 4027-4030.
-  H. Hajabolhassan, On colorings of graph powers.
Discrete Mathematics, 309 (2009), 4299-4305.
-  H. Hajabolhassan and A. Taherkhani, Graph powers and graph homomorphisms.
Electronic Journal of Combinatorics, 17 (2010), no. 1, Research Paper 17, 16 pp.

Thank You!

I would like to acknowledge Dr Ali Taherkhani for his useful comments.

