

# Cantor-Bernstein for Theories

Princeton Logic Group

December 14, 2020

The purpose of this note is to ask: under what conditions could a pair of theories  $(T_1, T_2)$  fail to have the Cantor-Bernstein or co-Cantor-Bernstein property?

**Definition.** We say that the pair  $(T_1, T_2)$  has the Cantor-Bernstein property just in case: if  $T_1$  and  $T_2$  are mutually faithfully interpretable, then  $T_1$  and  $T_2$  are bi-interpretable. In other words: if  $(T_1, T_2)$  does not have the Cantor-Bernstein property then (a)  $T_1$  and  $T_2$  are mutually faithfully interpretable, and (b)  $T_1$  and  $T_2$  are not bi-interpretable.

Here “mutually faithfully interpretable” means that there are conservative (strong, equality preserving) translations  $F : T_1 \rightarrow T_2$  and  $G : T_2 \rightarrow T_1$ .

**Definition.** We say that the pair  $(T_1, T_2)$  has the co-Cantor-Bernstein property just in case: if there are essentially surjective translations  $F : T_1 \rightarrow T_2$  and  $G : T_2 \rightarrow T_1$ , then  $T_1$  and  $T_2$  are bi-interpretable.

## 1 Examples of theory pairs that are not CB

1.  $T_1$  is the empty theory on a countably infinite propositional signature.  $T_2$  is the “fan theory” with axioms  $p_0 \vdash p_i$ , for  $i \geq 0$ . These theories are counterexamples both to the CB and the co-CB properties.

But are these theories “pathological” in some sense? The fact that these theories are propositional should not (I think) be seen as a pathology. However, these theories are incomplete, and the second of them is not finitely axiomatizable.

2. The many examples of pairs of set theories described in (Freire and Hamkins 2020, p 8).
3. The pair of theories described in (Andréka, Madarász, and Németi 2005).

## 2 Results

**Proposition.** *If  $T_1$  or  $T_2$  is complete, then  $(T_1, T_2)$  has the co-CB property.*

*Proof.* If  $T_1$  is complete, then every translation  $F : T_1 \rightarrow T_2$  is conservative. So if  $F : T_1 \rightarrow T_2$  is eso, then  $F$  is a strong equivalence, i.e.  $T_1$  and  $T_2$  are bi-interpretable.  $\square$

Recall that  $F : T_1 \rightarrow T_2$  is conservative iff  $F^* : M(T_2) \rightarrow M(T_1)$  is a full functor (Barrett 2020). Recall also that if  $F$  is a strong (equality-preserving) translation, then  $F^*$  preserves cardinality of models. In particular, if  $T_1$  and  $T_2$  are  $\aleph_0$ -categorical theories, then a translation  $F : T_1 \rightarrow T_2$  induces a group homomorphism  $F^* : \text{Aut}(M_2) \rightarrow \text{Aut}(M_1)$ , and  $F$  is conservative iff  $F^*$  is surjective. (In fact,  $\text{Aut}(M_i)$  is naturally a topological group, and I conjecture that  $F^*$  is a continuous group homomorphism.)

Recall that if  $T$  has countable signature, and if  $T$  is  $\aleph_0$ -categorical, then  $T$  is complete. The following result would be interesting because an  $\aleph_0$ -categorical theory is essentially characterized by a topological group, viz. the group of automorphisms of its unique (up to isomorphism) countable model.

**Conjecture.** *There are  $\aleph_0$ -categorical theories  $T_1$  and  $T_2$  that do not have the CB property.*

**Definition.** Let  $I$  be the set of axioms  $\{\exists_{>1}, \exists_{>2}, \dots\}$ . We say that a theory  $T$  is essentially finitely axiomatizable just in case there is a finite set  $E$  of axioms such that  $Cn(T) = Cn(E \cup I)$ .

**Conjecture.** *There are essentially finitely axiomatizable theories  $T_1$  and  $T_2$  that do not have the CB property.*

### 3 Conjecture: co-CB fails only for theories with many models

*Remark.* For a theory  $T$  and a cardinal number  $\kappa$ , let  $I(T, \kappa)$  be the number of non-isomorphic countable models of  $T$ . Interestingly,  $I(T, \aleph_0)$  can be countably infinite, or any finite cardinal besides 2. See (Marker 2006, p 155ff).

**Proposition.** *If  $F : T_1 \rightarrow T_2$  is essentially surjective, then for any fixed cardinal number  $\kappa$ ,  $I(T_2, \kappa) \leq I(T_1, \kappa)$ .*

*Proof.* Recall that the dual functor  $F^*$  is always faithful. If  $F : T_1 \rightarrow T_2$  is eso, then  $F^*$  is also full (Halvorsen 2019, Prop 6.6.13). In particular, for any models  $M, N$  of  $T_2$ , if  $F^*(M)$  is isomorphic to  $F^*(N)$ , then  $M$  is isomorphic to  $N$ . Now fix a cardinal number  $\kappa$ , and let  $[M(T_i)]_\kappa$  be the set of isomorphism classes of models of  $T_i$  of cardinality  $\kappa$ . Then  $F^*$  induces a one-to-one mapping from  $[M(T_2)]_\kappa$  into  $[M(T_1)]_\kappa$ .  $\square$

**Proposition.** *Suppose that  $T_2$  has finitely many non-isomorphic models of each cardinality. If  $F : T_1 \rightarrow T_2$  and  $G : T_2 \rightarrow T_1$  are essentially surjective, then  $(F^*, G^*)$  is an equivalence of categories. If  $T_1$  and  $T_2$  are proper theories then  $F^*$  is part of a homotopy equivalence.*

*Proof.* For the first part it will suffice to show that  $F^*$  is essentially surjective. By the previous proof,  $G^*$  induces an injection of  $[M(T_1)]_\kappa$  into  $[M(T_2)]_\kappa$ . Since the latter is finite, so is the former. Since  $F^*$  is an injection of one finite set into a not-larger finite set, it follows that  $F^*$  is bijection. Therefore  $F^*$  is essentially surjective.

The second part follows from Theorem 7.1 of (D'Arienzo, Pagano, and Johnson 2020).  $\square$

**Corollary.** *Let  $T_1$  and  $T_2$  be proper theories. If  $(T_1, T_2)$  violate the co-CB property then there is a cardinal number  $\kappa$  such that  $T_1$  and  $T_2$  have infinitely many non-isomorphic models of size  $\kappa$ .*

TO DO: I would like to come up with a similar necessary condition for  $T_1$  and  $T_2$  violating the CB property. However, we do not yet have any interesting result of the form: “if  $T_1$  or  $T_2$  is ... and  $F : T_1 \rightarrow T_2$  is conservative then  $F^* : M(T_2) \rightarrow M(T_1)$  is ...”

## 4 Groups that are not CB

**Definition.** Let  $P(\mathbb{N})$  be the permutation group of the natural numbers, equipped with the topology of pointwise convergence. (TO DO: explain the sense in which this topology on  $P(\mathbb{N})$  is definable from the theory of infinite sets. Explain more generally the sense in which for a  $\Sigma$ -structure  $M$ ,  $\text{Aut}(M)$  is naturally a topological group.)

**Fact.** *Let  $G$  be a subgroup of  $P(\mathbb{N})$ . Then  $G$  is the automorphism group of an  $\aleph_0$ -categorical theory iff  $G$  is a closed subset of  $P(\mathbb{N})$ .*

**Conjecture.** *There are closed subgroups  $G$  and  $H$  of  $P(\mathbb{N})$  such that  $G$  is isomorphic to a closed subgroup of  $H$  and vice versa, but  $G$  and  $H$  are not isomorphic.*

It is not difficult at all to find groups that violate the Cantor-Bernstein condition — but I do not immediately know if any of these groups are of the form  $\text{Aut}(M)$  for an  $\aleph_0$ -categorical structure  $M$ .

1. The group  $S_\infty$  of finite permutations of  $\mathbb{N}$  and the alternating group  $A_\infty$ . See <https://math.stackexchange.com/questions/1259081/if-there-are-injecti>
2. Infinite direct sums of  $\mathbb{Z}_{2^i}$ .
3. The free group on 2 generators and the free group on 3 generators.

**Proposition** (Ahlbrandt and Ziegler 1986). *Two countable  $\aleph_0$ -categorical structures are bi-interpretable iff their automorphism groups are isomorphic as topological groups.*

Conjecture: the previous result can be lifted to  $\aleph_0$ -categorical theories (with countable signature). But we need to be careful about terminology. First of all, Ahlbrandt and Ziegler are working with a notion of “interpretation” between structures of one language and structures of another language: given a  $\Sigma_1$ -structure  $\mathcal{M}_1$  and a  $\Sigma_2$ -structure  $\mathcal{M}_2$ , and interpretation  $f : \mathcal{M}_1 \rightarrow \mathcal{M}_2$  consists of a surjection  $f : U \rightarrow M_2$  where  $U$  is a definable subset of the domain of  $\mathcal{M}_1$  and  $M_2$  is the domain of  $\mathcal{M}_2$ , etc.

**Conjecture.** *An interpretation from  $\mathcal{M}_1$  to  $\mathcal{M}_2$  is a translation in our sense from  $\text{Th}(\mathcal{M}_1)$  to  $\text{Th}(\mathcal{M}_2)$ .*

The following proposition follows immediately from the fact that  $\aleph_0$ -categorical theories are complete.

**Proposition.** *If  $T$  is  $\aleph_0$ -categorical with unique model  $\mathcal{M}$ , then  $T$  is logically equivalent to  $Th(\mathcal{M})$ .*

This result would be interesting because it would show that the structure of an  $\aleph_0$ -categorical theory is captured by its countable model. i.e., there is no need to look at the models of higher cardinality and the arrows between them.

**Proposition** (Evans and Hewitt 1990). *There are closed subgroups  $G$  and  $H$  of  $P(\mathbb{N})$  that are isomorphic qua groups but not as topological groups. Hence, the corresponding structures are not bi-interpretable.*

The former result is intriguing: there is a group isomorphism  $\varphi : G \xrightarrow{\sim} H$  that does not correspond to a homotopy equivalence between the corresponding theories. In short: the symmetries of the model are not enough to capture the structure of the theories.

## 5 $\aleph_0$ -categorical theories

For this discussion, we restrict to theories with countable signatures. In this case, the downward Löwenheim-Skolem theorem shows that an  $\aleph_0$ -categorical theory is complete.

Question: How can we characterize the syntactic categories of  $\aleph_0$ -categorical theories? (Hint: Look at the Ryll-Nardzewski theorem [https://en.wikipedia.org/wiki/Omega-categorical\\_theory](https://en.wikipedia.org/wiki/Omega-categorical_theory) and (Cameron 1990, p 30). I suspect that the characterization will have something to do with finiteness of subobject lattices.)

TO DO: Establish a correspondence between the 2-category of  $\aleph_0$ -categorical theories and some subcategory of the 2-category of topological groups. Note 1: every group is a category, and group homomorphisms are functors. Note 2: all translations between such theories are conservative. It will be helpful to look at pages 106–107 of (Cameron 1990) where he describes conditions on a topological group  $G$  that ensure it corresponds to a categorical theory.

**Definition.** Given an  $\aleph_0$ -categorical theory  $T$ , let  $\mathcal{G}(T)$  be the topological group of automorphisms of its unique countable model.

**Conjecture.** *If  $F : T_1 \rightarrow T_2$  is a translation then  $F^*|_{\mathcal{G}(T_2)}$  is a continuous homomorphism from  $\mathcal{G}(T_2)$  to  $\mathcal{G}(T_1)$ .*

**Conjecture.** *There is a 2-functor  $\mathcal{G}$  from the 2-category of topological groups to the 2-category of  $\aleph_0$ -categorical theories. (This is not stated correctly yet: we should not expect all topological groups to occur in the domain. It is only the “nice” topological groups, i.e. the automorphism groups of  $\aleph_0$ -categorical theories.)*

**Conjecture.** *There is a 2-functor  $\mathcal{F}$  from the 2-category of  $\aleph_0$ -categorical theories and the 2-category of topological groups.*

## References

- Ahlbrandt, Gisela, and Martin Ziegler. 1986. “Quasi finitely axiomatizable totally categorical theories”. *Annals of Pure and Applied Logic* 30 (1): 63–82. doi:10.1016/0168-0072(86)90037-0.
- Andréka, Hajnal, Judit X Madarász, and István Németi. 2005. “Mutual definability does not imply definitional equivalence, a simple example”. *Mathematical Logic Quarterly* 51 (6): 591–597. doi:10.1002/malq.200410051.
- Barrett, Thomas William. 2020. “How to count structure”. *Noûs*. doi:10.1111/nous.12358.
- Cameron, Peter J. 1990. *Oligomorphic permutation groups*. Cambridge University Press. doi:10.1017/CB09780511549809.
- D’Arienzo, A., V. Pagano, and I. Johnson. 2020. *The 2-categorical structure of predicate theories*. arXiv: 2011.14056 [math.LO].
- Evans, David M, and Paul R Hewitt. 1990. “Counterexamples to a conjecture on relative categoricity”. *Annals of pure and applied logic* 46 (2): 201–209. doi:10.1016/0168-0072(90)90034-Y.
- Freire, Alfredo Roque, and Joel David Hamkins. 2020. “Bi-interpretation in weak set theories”. Under review, *Mathematics arXiv*. arXiv: 2001.05262 [math.LO]. <http://jdh.hamkins.org/bi-interpretation-in-weak-set-theories>.
- Halvorson, Hans. 2019. *The logic in philosophy of science*. Cambridge University Press. doi:10.1017/9781316275603.

Marker, David. 2006. *Model theory: an introduction*. Vol. 217. Springer Science & Business Media. doi:10.1007/b98860.