

Graphs Algorithms I

Breadth First Search



BFS(G, s)

$/* \ G = (V, E) \text{ and } s \in V \ */$

color[s] \leftarrow grey ; d[s] \leftarrow 0 ; p[s] \leftarrow NIL

For each $v \in V - \{s\}$ **do**

color[v] \leftarrow white

d[v] \leftarrow ∞

p[v] \leftarrow NIL

Q \leftarrow empty ; ENQ(Q, s)

$/* \ Q: \text{nodes that are discovered but not yet explored} \ */$

While Q is not empty **do**

u \leftarrow DEQ(Q)

For each $(u, v) \in E$ **do**

If color[v] = white **then do**

color[v] \leftarrow grey

d[v] \leftarrow d[u] + 1

p[v] \leftarrow u

ENQ(Q, v)

End If

End For

color[u] \leftarrow black

$/* \ \text{Explore } u \ */$

$/* \ \text{Explore edge } (u,v) \ */$

$/* \ \text{If } v \text{ is first discovered} \ */$

$/* \ \text{Done exploring } u \ */$

End While

End BFS



Breadth First Search

Proof of Correctness



Execute BFS(s) on a graph G

For every node v of G :



Execute BFS(s) on a graph G

For every node v of G:

v's **discovery** path from s : $s \rightarrow u_1 \rightarrow u_2 \rightarrow \dots \rightarrow u \rightarrow v$



Execute BFS(s) on a graph G

For every node v of G:

v's **discovery** path from s : $s \rightarrow u_1 \rightarrow u_2 \rightarrow \dots \rightarrow u \rightarrow v$

Length of discovery path : **d[v]**



Execute BFS(s) on a graph G

For every node v of G:

v's **discovery** path from s : $s \rightarrow u_1 \rightarrow u_2 \rightarrow \dots \rightarrow u \rightarrow v$

Length of discovery path : **d[v]**

v's **shortest** path from s : $s \rightarrow v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_k \rightarrow v$



Execute BFS(s) on a graph G

For every node v of G:

v's **discovery** path from s : $s \rightarrow u_1 \rightarrow u_2 \rightarrow \dots \rightarrow u \rightarrow v$

Length of discovery path : **$d[v]$**

v's **shortest** path from s : $s \rightarrow v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_k \rightarrow v$

Length of shortest path : **$\delta(s,v)$**
(Distance)



Execute BFS(s) on a graph G

For every node v of G:

v's **discovery** path from s : $s \rightarrow u_1 \rightarrow u_2 \rightarrow \dots \rightarrow u \rightarrow v$

Length of discovery path : **$d[v]$**

v's **shortest** path from s : $s \rightarrow v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_k \rightarrow v$

Length of shortest path : **$\delta(s,v)$**
(Distance)

Lemma 0: **$d[v]$** **$\delta(s,v)$**



Execute BFS(s) on a graph G

For every node v of G:

v's **discovery** path from s : $s \rightarrow u_1 \rightarrow u_2 \rightarrow \dots \rightarrow u \rightarrow v$

Length of discovery path : **$d[v]$**

v's **shortest** path from s : $s \rightarrow v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_k \rightarrow v$

Length of shortest path : **$\delta(s,v)$**
(Distance)

Lemma 0: **$d[v] \geq \delta(s,v)$**



We would like to prove the following:

Main Theorem:

After BFS(s), for every $v \in V$,

$$d[v] = \delta(s, v)$$



We would like to prove the following:

Main Theorem:

After BFS(s), for every $v \in V$,

$$d[v] = \delta(s, v)$$


That is: the **discovery** path to v is a **shortest** path to v



We first prove the following:

Lemma 1:


If u enters Q



We first prove the following:

Lemma 1:


If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then:



We first prove the following:

Lemma 1:


If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then:



We first prove the following:

Lemma 1:


If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then:



We first prove the following:

Lemma 1:

If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then:




$$d[u] = d[v]$$



We first prove the following:

Lemma 1:

If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then




We first prove the following:

Lemma 1:

If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then

$$d[u] \leq d[v]$$



$$d[u] \leq d[v]$$



We first prove the following:

Lemma 1:

If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then

$$d[u] \leq d[v]$$

Proof of Lemma 1:



We first prove the following:

Lemma 1:

If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then

$$d[u] \leq d[v]$$

Proof of Lemma 1:

Suppose, for contradiction, that Lemma 1 is false.



We first prove the following:

Lemma 1:

If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then

$$d[u] \leq d[v]$$

Proof of Lemma 1:

Suppose, for contradiction, that Lemma 1 is false.

- Let v be the **first** node that enter Q such that $d[u] > d[v]$ for some node u that entered Q before v .



We first prove the following:

Lemma 1:


If u enters Q **before** v enters Q during the execution of $\text{BFS}(s)$, then

$$d[u] \leq d[v]$$

Proof of Lemma 1:

Suppose, for contradiction, that Lemma 1 is false.


- Let v be the **first** node that enter Q such that $d[u] > d[v]$ for some node u that entered Q before v .




$$d[u] > d[v]$$



Proof of Lemma 1 contd:




Proof of Lemma 1 contd:



- $v \neq s$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
 - $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
- $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
 - $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
- $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
- $\Rightarrow d[u] =$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
 - $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
- $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
- $\Rightarrow d[u] = d[u'] + 1$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
 - $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
- $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
- $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$,




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v']$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are **distinct** nodes




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before v




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before $v \Rightarrow u'$ was discovered before v'




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before $v \Rightarrow u'$ was discovered before v'
 $\Rightarrow u'$ entered Q before v'




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before $v \Rightarrow u'$ was discovered before v'
 $\Rightarrow u'$ entered Q before v'




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before $v \Rightarrow u'$ was discovered before v'
 $\Rightarrow u'$ entered Q before v'
 $\Rightarrow d[u'] < d[v']$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 - $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 - $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before $v \Rightarrow u'$ was discovered before v'
 - $\Rightarrow u'$ entered Q before v'
 - $\Rightarrow d[u'] \leq d[v']$ By definition of v




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 - $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 - $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before $v \Rightarrow u'$ was discovered before v'
 - $\Rightarrow u'$ entered Q before v'
 - $\Rightarrow d[u'] \leq d[v']$ By definition of v
 - $\Rightarrow d[u'] + 1 \leq d[v'] + 1$




Proof of Lemma 1 contd:



- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before $v \Rightarrow u'$ was discovered before v'
 $\Rightarrow u'$ entered Q before v'
 $\Rightarrow d[u'] \leq d[v']$ By definition of v
 $\Rightarrow d[u'] + 1 \leq d[v'] + 1$
 $\Rightarrow d[u] \leq d[v]$



Proof of Lemma 1 contd:




- $v \neq s$ because no vertex u enters Q before s
- $u \neq s$ because $d[s] = 0$ and $d[v] \geq 0$
 - $\Rightarrow u$ and v entered Q during the exploration of some nodes, say u' and v' respectively
 - $\Rightarrow d[u] = d[u'] + 1$ and $d[v] = d[v'] + 1$
- Since $d[u] \neq d[v]$, $d[u'] \neq d[v'] \Rightarrow u' \neq v'$ i.e. u' and v' are distinct nodes
- u was discovered before $v \Rightarrow u'$ was discovered before v'
 - $\Rightarrow u'$ entered Q before v'
 - $\Rightarrow d[u'] \leq d[v']$ By definition of v
 - $\Rightarrow d[u'] + 1 \leq d[v'] + 1$
 - $\Rightarrow d[u] \leq d[v]$ Contradiction !



So we proved the following:

Lemma 1: If u enters Q **before** v enters Q then $d[u] \leq d[v]$



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s, v)$



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s, v)$

Proof of Main Theorem:



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$.



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s, v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s, x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s, v)$
- By Lemma 0, $d[v] > \delta(s, v)$
- Consider some **shortest** path in s to v in G




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) =$




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$
- Since u is closer to s than v , $d[u] = \delta(s,u)$ (By definition of v)




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$
- Since u is closer to s than v , $d[u] = \delta(s,u)$ (By definition of v)




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$
- Since u is closer to s than v , $d[u] = \delta(s,u)$ (By definition of v)
- So: $d[v] > \delta(s,v)$




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$
- Since u is closer to s than v , $d[u] = \delta(s,u)$ (By definition of v)
- So: $d[v] > \delta(s,v)$




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$
- Since u is closer to s than v , $d[u] = \delta(s,u)$ (By definition of v)
- So: $d[v] > \delta(s,v) = \delta(s,u) + 1$




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$
- Since u is closer to s than v , $d[u] = \delta(s,u)$ (By definition of v)
- So: $d[v] > \delta(s,v) = \delta(s,u) + 1$




Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:

Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$
- Since u is closer to s than v , $d[u] = \delta(s,u)$ (By definition of v)
- So: $d[v] > \delta(s,v) = \delta(s,u) + 1 = d[u] + 1$



Main Theorem:

After BFS(s), for every $v \in V$, $d[v] = \delta(s,v)$

Proof of Main Theorem:


Suppose, for contradiction, that there exists $x \in V$ such that $d[x] \neq \delta(s,x)$. Clearly $x \neq s$.

- Let v be the **closest** node from s such that $d[v] \neq \delta(s,v)$
- By Lemma 0, $d[v] > \delta(s,v)$
- Consider some **shortest** path in s to v in G
 - Let (u,v) be the last edge on that path
 - Clearly $\delta(s,v) = \delta(s,u) + 1$
- Since u is closer to s than v , $d[u] = \delta(s,u)$ (By definition of v)
- So: $d[v] > \delta(s,v) = \delta(s,u) + 1 = d[u] + 1$

$$d[v] > d[u] + 1$$

(*)

© 2019 by Jay Balasundaram and Sam Toueg. This document may not be posted on the internet without the written permission of the copyright owners.



Proof of Main Theorem contd:

$$d[v] > d[u] + 1 \quad (*)$$



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored.



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white

\Rightarrow When u is explored, u discovers v



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white

\Rightarrow When u is explored, u discovers v

$\Rightarrow d[v] = d[u] + 1$



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white

\Rightarrow When u is explored, u discovers v

$\Rightarrow d[v] = d[u] + 1$

Contradicting $(*)$!



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white

\Rightarrow When u is explored, u discovers v

$\Rightarrow d[v] = d[u] + 1$

Contradicting $(*)$!

Case 2. v is black



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white

\Rightarrow When u is explored, u discovers v

$\Rightarrow d[v] = d[u] + 1$

Contradicting $(*)$!

Case 2. v is black

$\Rightarrow v$ was explored before u is explored



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white

\Rightarrow When u is explored, u discovers v

$\Rightarrow d[v] = d[u] + 1$

Contradicting $(*)$!

Case 2. v is black

$\Rightarrow v$ was explored before u is explored

$\Rightarrow v$ entered Q before u enters Q



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white

\Rightarrow When u is explored, u discovers v

$\Rightarrow d[v] = d[u] + 1$

Contradicting $(*)$!

Case 2. v is black

$\Rightarrow v$ was explored before u is explored

$\Rightarrow v$ entered Q before u enters Q

\Rightarrow By Lemma 1, $d[v] \leq d[u]$



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Now consider the **color** of v **just before** u is explored. 3 possible cases:

Case 1. v is white

\Rightarrow When u is explored, u discovers v

$\Rightarrow d[v] = d[u] + 1$

Contradicting $(*)$!

Case 2. v is black

$\Rightarrow v$ was explored before u is explored

$\Rightarrow v$ entered Q before u enters Q

\Rightarrow By Lemma 1, $d[v] \leq d[u]$

Contradicting $(*)$!



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$


Case 3. v is grey (discovered but not explored)



Proof of Main Theorem contd:

$$d[v] > d[u] + 1 \quad (*)$$

Case 3. v is grey (discovered but not explored)




Proof of Main Theorem contd:

$$d[v] > d[u] + 1 \quad (*)$$

Case 3. v is grey (discovered but not explored)

\Rightarrow Some node w discovered v before u is explored




Proof of Main Theorem contd:

$$d[v] > d[u] + 1 \quad (*)$$

Case 3. v is grey (discovered but not explored)

\Rightarrow Some node w discovered v before u is explored

$\Rightarrow w$ is explored before u




Proof of Main Theorem contd:

$$d[v] > d[u] + 1 \quad (*)$$

Case 3. v is grey (discovered but not explored)

\Rightarrow Some node w discovered v before u is explored

$\Rightarrow w$ is explored before u and $d[v] = d[w] + 1$



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$


Case 3. v is grey (discovered but not explored)

\Rightarrow Some node w discovered v before u is explored

\Rightarrow w is explored before u and $d[v] = d[w] + 1$

(a)

(b)



Proof of Main Theorem contd:


$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Case 3. v is grey (discovered but not explored)

\Rightarrow Some node w discovered v before u is explored

\Rightarrow w is explored before u and $d[v] = d[w] + 1$
(a) (b)

(a) $\Rightarrow w$ enters Q before u



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Case 3. v is grey (discovered but not explored)


\Rightarrow Some node w discovered v before u is explored

\Rightarrow w is explored before u and $d[v] = d[w] + 1$
(a) (b)

(a) $\Rightarrow w$ enters Q before u

$\Rightarrow d[w] < d[u]$

By Lemma 1



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Case 3. v is grey (discovered but not explored)


\Rightarrow Some node w discovered v before u is explored

\Rightarrow w is explored before u and $d[v] = d[w] + 1$
(a) (b)

(a) $\Rightarrow w$ enters Q before u

$\Rightarrow d[w] \leq d[u]$

By Lemma 1



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Case 3. v is grey (discovered but not explored)

\Rightarrow Some node w discovered v before u is explored


\Rightarrow w is explored before u and $d[v] = d[w] + 1$
(a) (b)

(a) $\Rightarrow w$ enters Q before u

$$\Rightarrow d[w] \leq d[u]$$

By Lemma 1

$$\Rightarrow d[w] + 1 \leq d[u] + 1$$



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Case 3. v is grey (discovered but not explored)

\Rightarrow Some node w discovered v before u is explored

\Rightarrow w is explored before u and $d[v] = d[w] + 1$
(a) (b)


(a) $\Rightarrow w$ enters Q before u

$$\Rightarrow d[w] \leq d[u]$$

By Lemma 1

$$\Rightarrow d[w] + 1 \leq d[u] + 1$$

(b) $\Rightarrow d[v] \leq d[u] + 1$



Proof of Main Theorem contd:

$$\boxed{d[v] > d[u] + 1} \quad (*)$$

Case 3. v is grey (discovered but not explored)

\Rightarrow Some node w discovered v before u is explored

\Rightarrow w is explored before u and $d[v] = d[w] + 1$
(a) (b)

(a) $\Rightarrow w$ enters Q before u


$$\Rightarrow d[w] \leq d[u]$$

By Lemma 1

$$\Rightarrow d[w] + 1 \leq d[u] + 1$$

(b) $\Rightarrow d[v] \leq d[u] + 1$

Contradicting (*) !



We just proved

Theorem: After BFS(s), for every $v \in V$: $d[v] = \delta(s, v)$



We just proved

Theorem: After BFS(s), for every $v \in V$: $d[v] = \delta(s,v)$

So the BFS(s) **discovery** path from s to v is a **shortest** path from s to v in G

