
GENETICS · PREVENTION · NUTRACEUTICALS

Before the *Statin*

Detecting genetic and early cardiometabolic risk at 15–18, and testing *Citrus bergamot* as a first step

Noor Al Ahmad • Clinical Pharmacy • Evidence review & proposed RCT



−54.5% — **CHD per 1 mmol/L lower LDL-C, lifelong**
≈3× the benefit of lowering it later · Ference, JACC 2012


≈46% — **lower risk with a healthy lifestyle**
even in people with high genetic risk · Khera, NEJM 2016

−40 mg/dL — **LDL-C with bergamot**
pooled adult RCTs, low certainty · Chambari, Food Sci Nutr 2026

01 Risk starts in childhood, *not at 40*

LDL-C damage is cumulative: every year of exposure counts.

CUMULATIVE LDL-C EXPOSURE BY AGE (ILLUSTRATIVE MODEL)



Conceptual curves (LDL-C × years). Each line assumes a constant LDL-C; the earlier the slope falls, the later the threshold is reached. Ference 2012 & 2017; Jacobs (i3C), NEJM 2022.

Childhood risk → adult events
Risk factors at 3–19 y predict adult CV events (38,589 followed)

Genes are not destiny
High polygenic risk, healthy lifestyle: ≈46% fewer coronary events

A screening window exists
Universal lipid screening at 9–11 and 17–21 y (NHLBI)

Most teens don't meet statin criteria
A gap between "advice only" and drug therapy

02 Detect

Five checks that flag a high-risk teenager early.

GENETIC & EARLY RISK SCREEN · AGE 15–18

Polygenic risk score
CAD PRS from a genotyping array

Lp(a), once
Genetically set; measure at least once

Family history
Premature CVD in a 1st-degree relative

Lipids & apoB
Non-fasting LDL-C, TG, HDL-C

Metabolic profile
Waist, blood pressure, glucose / HOMA-IR, liver fat (MASLD), BMI

PATHWAY A · PROPOSED

Polygenic or metabolic risk
LDL-C below drug thresholds, no monogenic variant → **lifestyle + bergamot**, re-check every 6–12 months

PATHWAY B · GUIDELINE CARE

Monogenic FH / very high LDL
LDLR / APOB / PCSK9 variant → **statin as indicated**. Bergamot is not a substitute here.


03 Treat *early*

One supplement, Citrus bergamot, acting on several linked risk factors.

POOLED EFFECTS IN ADULT RCTS (9 TRIALS, 615 ADULTS) AND SINGLE TRIALS

BLOOD LIPIDS
−40 mg/dL LDL-C
TG −52 · HDL-C +5.6 mg/dL

BLOOD PRESSURE
−2.1 mmHg systolic
pooled, 9 RCTs



FATTY LIVER (MASLD)
−48 dB/m liver fat
vs −27 placebo · Ferro 2020

GLUCOSE & INSULIN
−12 mg/dL glucose
HOMA-IR −0.38 · pooled

WAIST & VISCERAL FAT
−1 cm waist (pooled)
visceral fat ↓ · Rondanelli 2020

VESSELS & INFLAMMATION
hs-CRP ↓
endothelial function ↑ · Fogacci 2022

Pooled values: Chambari et al., Food Sci Nutr 2026 (GRADE low to very low). Single-trial values are from adults; several used combination products.

04 The evidence so far

Randomised trials of bergamot (adults), plus the closest trial in young people.

Gliozzi 2013 · n = 77
Mixed hyperlipidaemia · BPF 1000 mg ± rosuvastatin, 30 d
↓LDL-C alone, and boosted the statin's effect

Mollace 2019 · n = 60
Type 2 diabetes + mixed hyperlipaemia · double-blind
↓LDL ↓TG ↓glucose, ↑HDL, fewer small dense LDL

Capomolla 2019 · n = 52
Metabolic syndrome, obesity · 90 days, 2 doses
−18% glucose · TG −32% · weight ↓

Rondanelli 2020 · n = 64
Overweight, mild ↑cholesterol · 12 weeks
↓VAT visceral fat, TC and LDL-C vs placebo at 30 d

Ferro 2020 · n = 102
Fatty liver · BPF + wild cardoon, 12 weeks
↓liver fat −48 vs −27 dB/m (placebo)

Fogacci 2022 · n = 60
Polygenic hypercholesterolaemia · combination, 8 weeks
↓LDL-C ↓hs-CRP, ↓fatty liver index vs placebo

Stonehouse 2025 · n = 42
Mild ↑LDL-C · sterols + bergamot + artichoke, 4 months
No effect on LDL-C vs placebo (−0.12 mmol/L, NS)

Guardamagna 2011 · age 8–16
Children with FH / FCH · red yeast rice + policosanols
−25% LDL-C; well tolerated (not bergamot)

−55 mg/dL — **LDL-C · 14 trials**
Sadeghi-Dehsahraei, Phytother Res 2022

−40 mg/dL — **LDL-C · 9 RCTs, 615 adults**
Chambari, Food Sci Nutr 2026 · GRADE low–very low

#1 — **of 10 nutraceuticals for LDL-C**
Network MA, 131 trials · Osadnik, Pharmacol Res 2022

The gap
No bergamot RCT in anyone under 18 has been published. That is what this study would test.

PROPOSED STUDY

Genetic screening + *Citrus bergamot* at 15–18: fewer risk factors before statins are needed?

1. **Screen** — Ages 15–18: PRS, Lp(a), family history, lipids, metabolic profile
2. **Select** — High polygenic or metabolic risk; monogenic FH referred to guideline care
3. **Randomise** — Bergamot polyphenol fraction vs placebo, both with lifestyle advice
4. **12 months** — Double-blind, with adherence and safety checks (liver, muscle, growth)
5. **Outcomes** — LDL-C / apoB, TG, HOMA-IR, BP, waist, liver fat, carotid IMT

REFERENCES

1. Ference BA, et al. Long-term exposure to lower LDL-C beginning early in life and CHD risk: a Mendelian randomization analysis. *J Am Coll Cardiol* 2012;60:2631–9.
2. Ference BA, et al. LDL causes atherosclerotic cardiovascular disease. EAS consensus. *Eur Heart J* 2017;38:2459–72.
3. Khera AV, et al. Genetic risk, adherence to a healthy lifestyle, and coronary disease. *N Engl J Med* 2016;375:2349–58.
4. Jacobs DR Jr, et al. Childhood cardiovascular risk factors and adult cardiovascular events. *N Engl J Med* 2022;386:1877–88.
5. NHLBI Expert Panel. Integrated guidelines for cardiovascular health and risk reduction in children and adolescents. *Pediatrics* 2011;128 Suppl 5:S213–56.
6. Kronenberg F, et al. Lipoprotein(a) in atherosclerotic CVD and aortic stenosis: EAS consensus. *Eur Heart J* 2022;43:3925–46.
7. Chambari M, et al. Bergamot supplementation and CVD risk factors: SR & MA. *Food Sci Nutr* 2026;14:e72357. doi:10.1002/fsn3.72357
8. Sadeghi-Dehsahraei H, et al. Bergamot and lipid profiles: SR & MA of RCTs. *Phytother Res* 2022;36:4409–24. doi:10.1002/ptr.7647
9. Osadnik T, et al. Network meta-analysis of nutraceuticals on lipid profile. *Pharmacol Res* 2022;183:106402. doi:10.1016/j.phrs.2022.106402
10. Gliozzi M, et al. BPF enhances rosuvastatin effect on LDL-C. *Int J Cardiol* 2013. doi:10.1016/j.ijcard.2013.08.125
11. Mollace V, et al. Lecithin formulation of BPF in T2DM with hyperlipaemia. *Endocr Metab Immune Disord Drug Targets* 2019. doi:10.2174/1871530319666181203151513
12. Capomolla AS, et al. Bergamot polyphenols in metabolic syndrome. *Nutrients* 2019;11:1271. doi:10.3390/nu11061271
13. Rondanelli M, et al. Bergamot phytosome, visceral fat and lipids. *Phytother Res* 2021;35:2045–56. doi:10.1002/ptr.6950
14. Ferro Y, et al. Bergamot and wild cardoon reduce liver steatosis. *Front Endocrinol* 2020;11:494. doi:10.3389/fendo.2020.00494
15. Fogacci F, et al. ANEMONE study, polygenic hypercholesterolaemia. *Nutrients* 2022;14:2099. doi:10.3390/nu14102099
16. Stonehouse W, et al. Nutraceutical combination and LDL-C: RCT. *Atherosclerosis* 2025;403:119177. doi:10.1016/j.atherosclerosis.2025.119177

Noor Al Ahmad
Clinical Pharmacy

Safety note: bergamot is a supplement, not an approved therapy. Teens with monogenic FH or very high LDL-C follow guideline statin care.