5.4 参考文献

目 录

5.4.1 参考文献目录.....2
5.4.2 参考文献译文及原文.....2

5.4.1 参考文献目录

文献编号	引用位置	参考文献
[1]	1.3.4.3 参考文献[1] 1.8 参考文献[1] 2.5 参考文献[1]	Yucai Wang, MD, PhD1,2; Shouhao Zhou, PhD3,4; Fang Yang, MD, PhD5,6; et al. Treatment-Related Adverse Events of PD-1 and PD-L1 Inhibitors in Clinical TrialsA Systematic Review and Meta-analysis. JAMA Oncol. April 25,2019.

5.4.2 参考文献译文及原文

[1] Yucai Wang, MD, PhD1,2; Shouhao Zhou, PhD3,4; Fang Yang, MD, PhD5,6; et al. Treatment-Related Adverse Events of PD-1 and PD-L1 Inhibitors in Clinical TrialsA Systematic Review and Meta-analysis. JAMA Oncol. April 25,2019.

➢ 译文

其他所有级别 irAEs 中，最常见的是腹泻（9.47%；95%CI，8.43%-10.58%），AST 升高（3.39%；95%CI，2.94%-3.89%），白癜风（3.26%；95%CI，2.80%-3.79%），丙氨酸氨基转移酶（ALT）增加（3.14%；95%CI，2.71%-3.62%），**肺炎（2.79%；95%CI，2.39%-3.23%）**和大肠杆菌-tis（1.24%；95%CI，0.99%-1.54%）（图 2C）。对于 3 级或更高级别的 irAE，最常见的是 AST 增加（0.75%；95%CI，0.56%-0.99%），其次是 ALT 增加（0.70%；95%CI，0.52%-0.93%），**肺炎（0.67%；95%CI，0.50%-0.89%）**，腹泻（0.59%；95%CI，0.45%-0.77%）和结肠炎（0.47；

➢ 原文

The most common other all-grade irAEs were diarrhea (9.47%; 95% CI, 8.43%-10.58%), AST increase (3.39%; 95% CI, 2.94%-3.89%), vitiligo (3.26%; 95% CI, 2.80%-3.79%), alanine aminotransferase (ALT) increase (3.14%; 95% CI, 2.71%-3.62%), pneumonitis (2.79%; 95% CI, 2.39%-3.23%), and colitis (1.24%; 95% CI, 0.99%-1.54%) (Figure 2C). For grade 3 or higher irAEs, AST increase (0.75%; 95% CI, 0.56%-0.99%) was most common, followed by ALT increase (0.70%; 95% CI, 0.52%-0.93%), pneumonitis (0.67%; 95% CI, 0.50%-0.89%), diarrhea (0.59%; 95% CI, 0.45%-0.77%), and colitis (0.47%;