Massachusetts General Hospital
Founding Member, Mass General Brigham

George A. "Doc" Lopez, MD
Laboratory for Regenerative Cell Therapy
White 536, 55 Fruit St, Boston, MA 02114

5	~~Aspirate the spent DPBS and 4~~ mL fresh DPBS to each dish. add	Volume added: 4 mL/dish # of dishes: _____
6	~~Aspirate the spent DPBS and 1 mL EDTA to each dish.~~	~~Volume added:~~ ~~_____ mL/dish~~ ~~# of dishes:~~ ~~_____~~
7	Return the dishes back to incubator for ~~8~~ minutes. 630	Time in: ~~[illegible]~~ 1:45 PM
8	During the 8 min of incubation, take out the new Matrigel-coated dishes and aspirate the Matrigel. Add 3 mL NutriStem hPSC XF medium to each dish. Note: Keep the dishes in BSC. No need to return to incubator.	Volume added: 3 mL/dish # of dishes: 2
9	During the 8 min of incubation, prepare a new 15 mL conical tube for the hPSC cell suspension. Note: if there is only one dish, this step can be skipped.	☑ NA ☑
10	After incubation, check the cells. If still attached, put them back in the incubator for 1 more min or until the cells start to detach.	Time out: 1:51 PM
11	Aspirate the EDTA from the two dishes. Take 1 mL of NutriStem hPSC XF medium (from the original aliquot) and add to one dish. Gently tap the dish to detach the colonies. If there are still colonies attached, gently pipette the colonies to create a cell suspension. Transfer the cell suspension to the 15 mL tube.	Volume added: 1 mL (~~[illegible] 2 dishes~~)
12	Take 1 mL of NutriStem hPSC XF medium (from the original aliquot) and add to the second dish. Gently tap the dish to detach the colonies. If there are still colonies attached, gently pipette the colonies to create a cell suspension. Transfer the cell suspension to the same 15 mL conical tube.	Volume added: 1 mL
13	Add 0.1 or 0.15 mL cell suspension drop by drop from the 15 mL conical tube to each Matrigel-coated dish (containing 3 mL NutriStem hPSC XF medium)	Volume added: 0.15 mL/dish # of dishes: 2
14	Gently stir to distribute cells evenly and return the dishes to the incubator.	Time in: 2:01

Comments:

Massachusetts General Hospital
Founding Member, Mass General Brigham

George A. "Doc" Lopez, MD
Laboratory for Regenerative Cell Therapy
White 536, 55 Fruit St, Boston, MA 02114

Process	hPSC passage	Date	04 Mar 2026
Time	1:07 PM	Operator	Harrison/Siang

Bill of Materials:

Cells	hPSC in culture	Cell type	PD05 mR03 hiPSC
# Passage	P6	Dish size	6 cm.
# of dishes	2	Thaw/last passage date	28Feb26
Passage ratio	1:10 (0.1 mL) ☐ 1:7 (0.15 mL) ☑	Cell condition	Normal; 75% confluent

Reagent	Lot#	Amount /container	# of container	Expiry date
Matrigel	33223001	8 mL	1	03 Feb 2026
Nutristem	250400812	40 mL	1	14 July 2026
DPBS	~~T3247188~~ AL3084602	40 mL	1	30 Jun 2027
0.5mM EDTA Or Versene	2624987	40 mL	1	20 Mar 26

Manufacturing Steps:

Step	Instruction	Data
1	Coat 6-cm dishes with 1.5 mL Matrigel solution per dish and incubate 37°C for at least 30 min.	# of dishes: 2 Time in 37°C 12:37 PM
2	Warm up the NutriStem medium in bead bath for 30 min.	☑
3	Take the two hPSC dishes out of incubator.	Time out: 1:30 PM
4	Aspirate the spent medium ~~and add 1 mL DPBS to each dish.~~	Volume added: 4 mL/dish # of dishes: