

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BCCA treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: BRAVTAX

DOCTOR'S ORDERS Htcm Wtkg	BSAm²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form	
DATE: To be given: Cycle #:	
Date of Previous Cycle:	
☐ Delay Treatment week(s) ☐ CBC & Diff, platelets day of treatment May proceed with doses as written if within 96 hours ANC greater than or equal to 1.5 x 90 x 10 ⁹ /L	x 10 ⁹ /L, Platelets <u>greater than</u>
Dose modification for:	
Proceed with treatment based on blood work from	
PREMEDICATIONS: 45 minutes prior to PACLitaxel: dexamethasone 20 mg IV in 50 mL NS over 15 minutes. 30 minutes prior to PACLitaxel: diphenhydrAMINE 50 mg IV and ranitidine 50 mg IV in 50 mL NS over 20 minutes. (compatible up to 3 hours when mixed in a bag)	
☐ Other:	
Have Hypersensitivity Reaction Tray and Protocol Available	
CHEMOTHERAPY: PACLitaxel 175 mg/m² x BSA =mg Dose Modification:mg/m² x BSA =mg IV in 250 to 500 mL (use non-DEHP bag) NS over 3 hours. (Use non DEHP tubing with 0.22 micron or smaller in-line filter)	
RETURN APPOINTMENT ORDERS	
☐ Return in three weeks for Doctor and Cycle ☐ Last Cycle. Return inweeks.	
Last Gycle. Returninweeks.	
CBC & Diff, Platelets prior to each cycle	
CBC & Diff, Platelets prior to each cycle	
CBC & Diff, Platelets prior to each cycle If clinically indicated: Bilirubin AST Other tests: Consults:	SIGNATURE: