

INTRODUCTION TO A MATROID AND ITS REPRESENTABILITY

HIDENORI SHINOHARA

ABSTRACT. This paper introduces a matroid. A matroid is a mathematical structure that abstracts the concept of linearly independence. The goal of this paper is to discuss the representability of a matroid with several examples and to introduce an important conjecture related to the representability of a matroid.

1. INTRODUCTION TO MATROID

A matroid is a mathematical structure that abstracts the concept of linearly independence. A linearly independent set of vectors has a lot of good properties. Interestingly, a lot of sets of other mathematical objects often share those properties. For example, any subset of a linearly independent set of vectors is always linearly independent. Consider a subset of edges of a graph that does not contain any cycle. Then any subset of it does not contain any cycle. The matroid theory is an attempt to mathematically formalize those properties and investigate in them.

Definition 1.1. A matroid $M = (E, \mathcal{I})$ is a pair such that E is a finite set of elements, and \mathcal{I} is a family of subsets of E with the following properties:

- $\emptyset \in \mathcal{I}$
- For any $A \in \mathcal{I}$, any subset of A is in \mathcal{I} .
- For any $A, B \in \mathcal{I}$ such that $|A| < |B|$, there always exists $x \in B - A$ such that $A \cup \{x\} \in \mathcal{I}$.

E is called the ground set, \mathcal{I} is called the independent sets. A subset of E is called independent if and only if it is in \mathcal{I} . In this paper, we will put our focus on a matroid with a finite ground set.

There are some basic matroids that are important in the following discussions. We will start by introducing a column matroid. A column matroid is constructed from a matrix over a field \mathbb{F} .

Definition 1.2. Let a matrix A with m rows over \mathbb{F} be given. A column matroid M of A is a matroid with a ground set $\{1, 2, \dots, m\}$. A subset of E is independent in M if and only if the set of the corresponding column vectors is linearly independent.

Theorem 1.3. *A column matroid is indeed a matroid.*

Proof. First, an empty set of vectors is linearly independent by definition. Suppose there exists a linearly independent set of vectors that has a linearly dependent subset. Let $A = \{a_1, \dots, a_k\} \subseteq B = \{b_1, \dots, b_n\}$ be such sets. Since A is linearly dependent, there exist constants c_1, \dots, c_k such that $c_1 a_1 + \dots + c_k a_k = 0$ and

not all c_i 's are 0. However, that implies that we can find constants d_1, \dots, d_n such that $d_1 b_1 + \dots + d_n b_n = 0$ and not all d_i 's are 0. This is a contradiction since B is supposed to be independent. Therefore, such subset must not exist, and thus all subsets of linearly independent sets are linearly independent. Now we want to prove the third property. Let $A = \{a_1, \dots, a_k\}, B = \{b_1, \dots, b_n\}$ be linearly independent sets, and assume that $k < n$. Suppose for each $i = 1, \dots, n$, $b_i \in \text{span}\{a_1, \dots, a_k\}$. It means that the span of B is a subspace of the span of A , which has smaller dimension than n . That is a contradiction. Therefore, there exists i such that $b_i \notin \text{span}\{a_1, \dots, a_k\}$. For such i , $\{a_1, \dots, a_k, b_i\}$ should be linearly independent. Since a column matroid satisfies the three properties, it is indeed a matroid. \square

Definition 1.4. A uniform matroid $U_{r,k}$ is a matroid such that $E = \{1, \dots, k\}$ and $\mathcal{I} = \{X \mid X \subseteq E, |X| \leq r\}$.

It is easy to see that a uniform matroid is indeed a matroid.

Here are some important results that will show up later in this paper.

Theorem 1.5. *All maximal independent sets have the same size.*

Proof. Let X, Y be maximal independent sets of some matroid. Suppose $|X| \neq |Y|$. Without loss of generality, $|X| < |Y|$. By the third property of a matroid, there exists $e \in Y - X$ such that $X \cup \{e\}$ is independent. It is a contradiction since X is a maximal independent set. Therefore $|X| = |Y|$. \square

This is indeed true in linear algebra. Given a matrix, any maximal independent subset of column vectors always has the same size. In linear algebra, this number is often referred to as the *dimension* of a vector space, or the *rank* of a matrix. In the matroid theory, we use the term *rank* as well.

Definition 1.6. The rank of a matroid is the size of a maximal independent set.

The rank of a column matroid is equal to the rank of the matrix since the subset of a ground set is independent if and only if the subset of column vectors is linearly independent.

Definition 1.7. Let $M = (E, \mathcal{I})$ be given. $e \in E$ is called a loop if $\{e\} \notin \mathcal{I}$.

We will conclude this chapter by introducing the notion of isomorphic matroids.

Definition 1.8. Let $M_1 = (E_1, \mathcal{I}_1), M_2 = (E_2, \mathcal{I}_2)$ be given. M_1, M_2 are isomorphic to each other if there exists a bijective mapping $\phi : E_1 \rightarrow E_2$ such that $\forall X \subseteq E_1, X \in \mathcal{I}_1 \iff \{\phi(e) : e \in X\} \in \mathcal{I}_2$.

Two isomorphic matroids have the same structure.

2. INTRODUCTION TO REPRESENTABILITY

We will start this chapter by defining the representability of a matroid.

Definition 2.1. A matroid $M = (E, \mathcal{I})$ is representable over a field \mathbb{F} if there exists a matrix A over \mathbb{F} such that the column matroid of A is isomorphic to M .

Therefore, if the matroid indeed succeeded in abstracting the concept of linearly independence, there should be some matroids that are *not* representable over some fields. If all matroids are representable over every field, it means that we are simply discussing linear algebra using different terms.

Before introducing some unrepresentable matroids, we will start by introducing a nice property of representable matroids.

Theorem 2.2. *Let $M = (E, \mathcal{I})$ be a matroid that is representable over \mathbb{F} . Let r be a rank of M and $k = |E|$. Then there exists a matrix $A \in \mathbb{F}^{r \times k}$ such that M is isomorphic to the column matroid of A .*

Proof. Let B a matrix over \mathbb{F} such that the column matroid of B is isomorphic to M . It is easy to see that the number of columns of B is k . From linear algebra, we know that elementary row operations preserve the linear independency of column vectors. Let R be the reduced row echelon form of B . Since R must have a rank of r , it only has r leading zeros. In other words, R has exactly r non-zero row vectors. Removing zero rows clearly does not affect the linear independency. Therefore, we found a matrix in $\mathbb{F}^{r \times k}$ whose column matroid is isomorphic to M . \square

This property is useful when proving that a matroid is unrepresentable over some field.

Here are some matroids that are not representable over some fields to show that not all matroids are representable over every field.

Theorem 2.3. *$U_{2,4}$ is not representable over $GF(2)$.*

A matroid is called a *binary matroid* if it is representable over $GF(2)$.

Proof. We know that $U_{2,4}$ has a rank of 2. If $U_{2,4}$ is representable over $GF(2)$, there exists a matrix $A \in GF(2)^{2 \times 4}$ such that A 's column matroid is isomorphic to $U_{2,4}$. Since $GF(2)$ only has two elements, there are only four possible column vectors of size 2. $\left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$ A must not contain $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ since it is a loop. Since A has 4 columns and there are only 3 different column vectors, we know that there are two columns in A that have the exact same column vectors. This is also a contradiction since a subset of such two elements will not be independent. Therefore, $U_{2,4}$ is not representable over $GF(2)$. \square

$U_{2,4}$ is representable over some field such as \mathbb{R} . For example, the column matroid of $\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{pmatrix}$ is isomorphic to $U_{2,4}$.

Now, we will introduce Fano matroid. Fano matroid can be constructed from Fano plane. Fano matroid is one of the examples of matroids that are representable over $GF(2)$, but not over \mathbb{R} . We will start by defining Fano matroid mathematically.

Definition 2.4. Fano matroid is a matroid with a ground set $\{1, 2, \dots, 7\}$. A set of vertexes is independent if it satisfies one of the followings:

- (1) it contains less than or equal to 2 elements,
- (2) it contains exactly 3 points and they are not on the same line.

Any set of vertexes that have more than 3 elements is dependent.

For example, $\{4, 5\}$ and $\{1, 2, 3\}$ are independent, but $\{1, 4, 2\}$, $\{1, 5, 7\}$, and $\{4, 5, 6\}$ are not independent as each of them is on one line.

Theorem 2.5. *Fano matroid is indeed a matroid.*



FIGURE 1. Fano plane

Proof. An empty set is independent since it contains less than 2 elements. Any independent set has at most three elements, so any proper subset of it has at most two elements. Therefore, any subset of independent set is always independent. The third property can be proved by checking each case. Since any set with 2 or less elements is independent, we only need to care about the case when we have a set with 2 elements and a set of three elements. Let $A = \{a_1, a_2, a_3\}$, $B = \{b_1, b_2\}$ be independent sets. It is easy to see from the figure that there must be exactly one line that goes through both b_1 and b_2 . Let x denote the third point on such line. Then, adding any element other than b_1, b_2, x to B will generate an independent set of three elements. Therefore, we just need to make sure that $A \neq \{b_1, b_2, x\}$. That cannot be the case since $\{b_1, b_2, x\}$ is dependent and A is independent. Therefore, there must be an element $a \in A - B$ such that $B \cup \{a\}$ is independent. Since Fano matroid satisfies all three properties, it is indeed a matroid. \square

Here is an interesting property of Fano matroid

Theorem 2.6. *If Fano matroid is representable over a field \mathbb{F} , $1 + 1 = 0$ in that field.*

Proof. Suppose Fano matroid is representable over a given field \mathbb{F} . Let $A \in \mathbb{F}^{3 \times 7}$ such that A 's column matroid is isomorphic to Fano matroid. Let R be a row reduced echelon form of A . Then the first three columns should be identical to I_3 since R has a rank of 3. Since $\{1, 2, 4\}$ is dependent, $R_{3,4}$ is 0. Applying the same argument to $\{2, 3, 5\}$, $\{1, 3, 6\}$, we get the following:

$$\begin{pmatrix} 1 & 0 & 0 & ? & 0 & ? & ? \\ 0 & 1 & 0 & ? & ? & 0 & ? \\ 0 & 0 & 1 & 0 & ? & ? & ? \end{pmatrix}$$

Since multiplying a non-zero constant to some column does not affect on the linearly independency, assume that the first non-zero elements of 4, 5, 6th columns are all

$$1. \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 & ? \\ 0 & 1 & 0 & ? & 1 & 0 & ? \\ 0 & 0 & 1 & 0 & ? & ? & ? \end{pmatrix}$$

Let $a = R_{2,4}, b = R_{3,5}$. $\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 & ? \\ 0 & 1 & 0 & a & 1 & 0 & ? \\ 0 & 0 & 1 & 0 & b & ? & ? \end{pmatrix}$ Since $\{4, 5, 6\}$ is dependent,

$R_{3,6}$ must be $-ab$. $\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 & ? \\ 0 & 1 & 0 & a & 1 & 0 & ? \\ 0 & 0 & 1 & 0 & b & -ab & ? \end{pmatrix}$ We need to look into the seventh

column. The seventh column actually cannot contain any zero. For example, suppose $R_{1,7} = 0$. Then, $\{2, 3, 7\}$ would be dependent. That's a contradiction. Similar arguments apply to the case of $R_{2,7} = 0, R_{3,7} = 0$. By multiplying a non-zero constant, we get:

$\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & a & 1 & 0 & ? \\ 0 & 0 & 1 & 0 & b & -ab & ? \end{pmatrix}$ Since $\{3, 4, 7\}$ is dependent, $R_{2,7} = a$.

And since $\{1, 5, 7\}$ is dependent, $R_{3,7} = b * R_{2,7} = ab$. $\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & a & 1 & 0 & a \\ 0 & 0 & 1 & 0 & b & -ab & ab \end{pmatrix}$

$\{2, 6, 7\}$ is dependent as well. By observing those three columns, it is easy to see that $-(-ab) + ab$ must be 0. Since neither of a, b is 0, $ab(1+1) = 0$ implies $1+1 = 0$. (If a is 0, 1st column and 4th column would be identical. The similar argument applies to b .) \square

This theorem is powerful, since this implies that Fano matroid is not representable over \mathbb{R} .

Corollary 2.7. *Fano matroid is not representable over \mathbb{R} .*

Proof. In \mathbb{R} , $1 + 1 \neq 0$. Therefore, Fano matroid is not representable over \mathbb{R} . \square

Theorem 2.8. *If a matroid $M = (E, \mathcal{I})$ only contains at most 3 non-loop elements, it is representable over any field \mathbb{F} .*

A matroid is called *regular* if it can be represented over any field.

Proof. I'll use rank here. \square

3. MORE DISCUSSION ON MATROID REPRESENTABILITY

This chapter will introduce a new concept, a matroid minor, which is crucial when discussing the matroid representability. Rota's conjecture will be also introduced at the end of the chapter. (I am thinking of adding a sketch of the proof by the end of the semester if I can understand the outline)

In order to introduce a matroid minor, we first need to introduce two operations on matroids. Deletion and contraction.

Definition 3.1. Let $M = (E, \mathcal{I})$, $X \subseteq E$ be given. $M \setminus X$ denotes a deletion of X in M and is defined to be $(E - X, \{I \in \mathcal{I} \mid X \cap I = \emptyset\})$

Note that $M \setminus e$ for some element $e \in E$ is equivalent to $M \setminus \{e\}$.

Theorem 3.2. *Let $M = (E, \mathcal{I})$, $X \subseteq E$ be given. $M \setminus X$ is indeed a matroid*

Proof. Let $M' = (E', \mathcal{I}') = M \setminus X$. Since $\emptyset \in \mathcal{I}$ and $X \cap \emptyset = \emptyset$, $\emptyset \in \mathcal{I}'$. Let $I \in \mathcal{I}', J \subseteq I$. Since $I \in \mathcal{I}$, $J \in \mathcal{I}$. Since $J \subseteq I$ and $I \cap X = \emptyset$, $J \cap X = \emptyset$. Therefore, $J \in \mathcal{I}'$. Let $A, B \in \mathcal{I}'$ such that $|A| < |B|$. Since $A, B \in \mathcal{I}'$, we know that $A, B \in \mathcal{I}$. Therefore, we can find $x \in B - A$ such that $(A \cup \{x\}) \in \mathcal{I}$. Since $(A \cup \{x\}) \subseteq (A \cup B)$ and $X \cap A = X \cap B = \emptyset$, $X \cap (A \cup \{x\}) = \emptyset$. Therefore, $A \cup \{x\} \in \mathcal{I}'$. Thus we have found such $x \in B - A$ that $A \cup \{x\} \in \mathcal{I}'$. Since $M' = M \setminus X$ follows the three properties, it is indeed a matroid. \square

Definition 3.3. Let $M = (E, \mathcal{I})$, $e \in E$ be given. M/e denotes contraction of M by e and $M/e = \begin{cases} M \setminus e, & \text{if } e \text{ is a loop,} \\ (E - \{e\}, \{I \in \mathcal{I} \mid e \notin I, (I \cup \{e\}) \in \mathcal{I}\}), & \text{otherwise.} \end{cases}$

Theorem 3.4. *Contraction by an element indeed generates a matroid.*

Proof. If e is a loop, M/e is obviously a matroid since we know that deletion always generates a matroid. Suppose otherwise. Let \mathcal{I}' denote the independent sets of M/e . First, $\emptyset \in \mathcal{I}, e \notin \emptyset$. Since e is not a loop, $(\emptyset \cup \{e\}) \in \mathcal{I}$. Therefore, $\emptyset \in \mathcal{I}'$. Let $I \in \mathcal{I}', J \subseteq I$. Since $I \in \mathcal{I}$, $J \in \mathcal{I}$. Since $e \notin I$, $e \notin J$. Since $(I \cup \{e\}) \in \mathcal{I}$ and $J \subseteq I$, $(J \cup \{e\}) \in \mathcal{I}$. Therefore, $J \in \mathcal{I}'$. Let $A, B \in \mathcal{I}'$ such that $|A| < |B|$. Let $A' = A \cup \{e\}, B' = B \cup \{e\}$. Since $A, B \in \mathcal{I}'$, $A', B' \in \mathcal{I}$. Since $e \notin A, e \notin B$, $|A'| < |B'|$. Let $x \in B' - A'$ such that $A' \cup \{x\} \in \mathcal{I}$. Since $B' - A' = B - A$, $x \in B - A$. For such x , we just showed that $A \cup \{e\} \cup \{x\} \in \mathcal{I}$. Also, $x \neq e$ since $e \in A'$. Therefore, $A \cup \{x\} \in \mathcal{I}'$. Hence, we have found $x \in B - A$ such that $A \cup \{x\} \in \mathcal{I}'$. Since this follows three properties given in the definition, this is indeed a matroid. Therefore, contraction by an element indeed generates a matroid. \square

Here are a few simple yet useful results about deletion.

Theorem 3.5. *Let a matroid $M = (E, \mathcal{I})$, $X \subseteq E$ be given. Let A be an independent set in $M \setminus X$. Then A is independent in M .*

Proof. The independent sets of $M \setminus X$ is $\{I \in \mathcal{I} \mid (I \cap X) = \emptyset\}$. It is easy to see that it is a subset of \mathcal{I} . Since A is in the subset of \mathcal{I} , A must be in \mathcal{I} . \square

In other words, this means that deletion never "adds" a new element to the independent sets.

Theorem 3.6. *The deletion of a loop does not change the independent sets.*

Proof. Let $M = (E, \mathcal{I})$, $e \in E, \{e\} \notin \mathcal{I}$. The independent sets of $M \setminus e$ is $\{I \in \mathcal{I} \mid (I \cap \{e\}) = \emptyset\}$. Since $\{e\}$ is a loop, no independent set can contain e . Therefore, the independent sets of $M \setminus e$ is identical to \mathcal{I} . \square

Now that we have defined contraction of a matroid by an element, we can define contraction by a subset of a ground set.

Definition 3.7. Let $M = (E, \mathcal{I})$, $X = \{x_1, \dots, x_k\} \subseteq E$. M/X is defined to be $((M/x_1)/x_2) \cdots /x_k$.

It is not obvious that this is well-defined. In other words, it is not obvious that the order of contraction does not matter. The following theorem shows that the order does not matter.

Theorem 3.8. *For any given matroid $M = (E, \mathcal{I})$, $(M/e)/f = (M/f)/e$ for any $e \neq f \in E$.*

Proof. There are a few cases.

- (1) e, f are both loops.
 $(M/e)/f = (M \setminus e)/f$ Since deletion of a loop does not change the independent sets, f is a loop in $(M \setminus e)$. Therefore, $(M/e)/f = (M \setminus e) \setminus f$. Again, deletion of f does not change the independent sets since f is a loop in (M/e) . Therefore, we have $(M/e)/f = (E - \{e, f\}, \mathcal{I})$. By symmetry, $(M/e)/f = (M/f)/e$.
- (2) One of e, f is a loop, and the other one is not.
 Without loss of generality, assume e is a loop. $(M/e)/f = (M \setminus e)/f$. Since deletion of a loop does not change the independent set, the independent set of (M/e) is \mathcal{I} . Therefore, the independent set of $(M/e)/f$ is $\mathcal{I}' = \{I \in \mathcal{I} \mid f \notin I, (I \cup \{f\}) \in \mathcal{I}\}$. On the other hand, it is easy to see that \mathcal{I}' is identical to the independent sets of (M/f) . Since contraction by an element does not add new elements to the independent sets, e is a loop in (M/f) . Since deletion by a loop does not change the independent sets, the independent sets of $(M/f)/e$ is \mathcal{I}' . Now we confirmed that $(M/e)/f$ and $(M/f)/e$ have the same independent sets. Therefore, $(M/e)/f = (M/f)/e$.
- (3) Neither of them is a loop, and $\{e, f\} \in \mathcal{I}$.
 The independent set of M/e is $\mathcal{I}' = \{I \in \mathcal{I} \mid e \notin I, (I \cup \{e\}) \in \mathcal{I}\}$. Since $\{e, f\} \in \mathcal{I}$, $\{f\} \in \mathcal{I}'$. Therefore, f is not a loop in M/e . Hence, the independent sets of $(M/e)/f$ is $\mathcal{I}'' = \{I \in \mathcal{I}' \mid f \notin I, (I \cup \{f\}) \in \mathcal{I}'\}$. \mathcal{I}'' is actually equivalent to $S = \{I \in \mathcal{I} \mid e \notin I, f \notin I, (I \cup \{e, f\}) \in \mathcal{I}\}$. We can prove $\mathcal{I}'' = S$ by starting to show that $\mathcal{I}'' \subseteq S$. Let $I \in \mathcal{I}''$. Since I is an independent set of $(M/e)/f$, we know that $e, f \notin I$. Since $(I \cup \{f\}) \in \mathcal{I}'$, we also know that $((I \cup \{f\}) \cup \{e\}) \in \mathcal{I}$. Therefore, $(I \cup \{e, f\}) \in \mathcal{I}$. Thus $I \in S$, and $\mathcal{I}'' \subseteq S$. Now, we want to show that $S \subseteq \mathcal{I}''$. Let $I \in S$. By the definition of S , we know that $e, f \notin I$. Since $(I \cup \{e, f\}) \in \mathcal{I}$, we know that $(I \cup \{e\}) \in \mathcal{I}$. Since $((I \cup \{f\}) \cup \{e\}) \in \mathcal{I}$ and $e \notin (I \cup \{f\})$, we know that $(I \cup \{f\}) \in \mathcal{I}'$. Since $f \notin I$ and $(I \cup \{f\}) \in \mathcal{I}'$, $I \in \mathcal{I}''$. Hence, $S \subseteq \mathcal{I}''$.
 Combining these two results, we know that $S = \mathcal{I}''$. By the symmetry, $(M/e)/f$ and $(M/f)/e$ have the same independent sets. Therefore $(M/e)/f = (M/f)/e$.
- (4) Neither of e, f is a loop, but $\{e, f\} \notin \mathcal{I}$.
 The independent set of M/e is $\mathcal{I}' = \{I \in \mathcal{I} \mid e \notin I, (I \cup \{e\}) \in \mathcal{I}\}$. Since $\{e, f\} \notin \mathcal{I}$, f is a loop in M/e . Therefore, $(M/e)/f = (M/e) \setminus f$. Since the deletion of a loop does not change the independent sets, the independent sets of $(M/e)/f$ is \mathcal{I}' . By applying the same argument, the independent set of $(M/f)/e$ is $\mathcal{I}'' = \{I \in \mathcal{I} \mid f \notin I, (I \cup \{f\}) \in \mathcal{I}\}$. We want to show that $\mathcal{I}' = \mathcal{I}''$. By the symmetry, it suffices to show that $\mathcal{I}' \subseteq \mathcal{I}''$. Let $I \in \mathcal{I}'$. Since $\{e, f\}$ is dependent, $f \notin I$. (Otherwise, $I \cup \{e\}$ would be dependent.) Since both $(I \cup \{e\})$ and $\{f\}$ are independent, we can grow $\{f\}$ by adding elements from $(I \cup \{e\})$ until they have the same size. Since $\{e, f\}$ is dependent, we never add e . In other words, we add every element from \mathcal{I} . It means that $I \cup \{f\}$ is independent. Therefore, $I \in \mathcal{I}''$, and thus $\mathcal{I}' \subseteq \mathcal{I}''$. By symmetry, $\mathcal{I}'' \subseteq \mathcal{I}'$. Therefore, $\mathcal{I}' = \mathcal{I}''$.

Therefore, in any case, $(M/e)/f = (M/f)/e$. □

List of topics I could discuss further:

- More properties about contraction deletion
 - $(M/e) \setminus f = (M \setminus f)/e$?
 - Therefore, any minor can be represented as $(M \setminus A)/B$

3.1. What do contraction and deletion mean in graphs and vector spaces?

3.2. Why do these matter? Because if a matroid is representable over some field \mathbb{F} , its minor is always representable over \mathbb{F} .

Theorem 3.9. *Let a matroid $M = (E, \mathcal{I})$ such that it is representable over \mathbb{F} . Any minor of M is representable over \mathbb{F} .*

Proof. prove! □

However, neither the converse nor the inverse of this theorem is always true. Any matroid has a representable minor since $U_{0,k}$ is a minor of any matroid. Also, $U_{2,4}$ is not a binary matroid, but any minor of it only contains at most 3 elements, so we know that any minor of $U_{2,4}$ is regular by the theorem. Rota's conjecture is about unrepresentable matroids any of whose minor is representable.

3.3. The introduction of Rota's conjecture.

Theorem 3.10. *For any finite field \mathbb{F} , there are only finitely many unrepresentable matroids all of whose minors are representable. In other words, there are only finitely many excluded minor.*

REFERENCES

- [1] "WHAT IS A MATROID?" Oxley
- [2] "On Matroid Representability and Minor Problems" Hlineny
- [3] "Solving Rota's Conjecture" Geelen, Gerards, Whittle
- [4] "Advanced Graph Theory, lecture 1" Rudi Pendavingh
- [5] "Matroids You Have Known" Neel, Neudauer