

MATH 602(HOMEWORK 1)

HIDENORI SHINOHARA

Exercise. 1

- Let $p \in V(I \cap J)$. For any $\sum_{i=1}^n f_i g_i \in IJ$, we have $f_i g_i \in I \cap J$ for each i . Thus $(\sum_{i=1}^n f_i g_i)(p) = 0$, so $p \in V(IJ)$. Let $p \in V(IJ)$. Let $f \in I \cap J$. Then $f^2 \in IJ$, so $(f(p))^2 = 0$. Thus $f(p) = 0$, so $p \in V(I \cap J)$. Therefore, $V(I \cap J) = V(IJ)$.
Let $p \in V(I) \cup V(J)$. Then either all polynomials in I vanish at p or all polynomials in J vanish at p . Thus all the polynomials in the intersection must vanish at p . Thus $V(I) \cup V(J) \subset V(I \cap J)$. On the other hand, let $p \in V(I \cap J) \setminus (V(I) \cup V(J))$. If no such element exists, we are done. Then every polynomial in the intersection vanishes at p . Let $f \in I$ and $g \in J$ be polynomials that do not vanish at p . Then $fg \in I \cap J$, so $(fg)(p) = 0$. However, this is impossible because $f(p) \neq 0$ and $g(p) \neq 0$. Therefore, $V(I) \cup V(J) = V(I \cap J)$.
- $p \in V(I + J)$ if and only if $\forall f \in I + J, f(p) = 0$ if and only if $\forall f \in I, f(p) = 0$ and $\forall f \in J, f(p) = 0$ if and only if $p \in V(I) \cap V(J)$.
- If every polynomial in J vanishes at a point, every polynomial in I must vanish at that point.
- If a polynomial vanishes in Y , then it must vanish in X .
- TODO

Exercise. 2

•

$$\begin{aligned}
 y \in (I_1 + I_2)^e &\iff y \in f(I_1 + I_2)B \\
 &\iff \exists x_1, x_2 \in I_1, I_2, b \in B, y = f(x_1 + x_2)b \\
 &\iff \exists x_1, x_2 \in I_1, I_2, b \in B, y = f(x_1)b + f(x_2)b \\
 &\iff y \in I_1^e + I_2^e.
 \end{aligned}$$

•

$$\begin{aligned}
 y \in (I_1 \cap I_2)^e &\implies y \in f(I_1 \cap I_2)B \\
 &\implies \exists x \in I_1 \cap I_2, b \in B, y = f(x)b \\
 &\implies (\exists x \in I_1, b \in B, y = f(x)b) \text{ and } (\exists x \in I_2, b \in B, y = f(x)b) \\
 &\implies y \in I_1^e, y \in I_2^e \\
 &\implies y \in I_1^e \cap I_2^e.
 \end{aligned}$$