## MATH 611 (DUE 10/23)

## HIDENORI SHINOHARA

## 1. SIMPLICIAL AND SINGULAR HOMOLOGY

**Exercise.** (Problem 2) Show that the  $\Delta$ -complex obtained from  $\Delta^3$  by performing the edge identifications  $[v_0, v_1] \sim [v_1, v_3]$  and  $[v_0, v_2] \sim [v_2, v_3]$  deformation retracts onto a Klein bottle. Find other pairs of identifications of edges that produce  $\Delta$ -complexes deformation retracting onto a torus, a 2-sphere, and  $\mathbb{R}\mathbf{P}^2$ .

*Proof.* Maybe something like this? Either way, I noticed that it looks like it contains  $2 \mathbb{R}P^2$ .

**Exercise.** (Problem 4) Compute the simplicial homology groups of the triangular parachute obtained from  $\Delta^2$  by identifying its three vertices to a single point.

*Proof.* Let  $v_0$  denote the only vertex,  $e_1, e_2, e_3$  denote the three edges of the parachute, and  $\sigma$  denote the face of the parachute.  $C_k = 0$  for  $k \geq 3$  because  $\Delta^2$  with the vertices identified does not contain any k-dimensional simplicies.  $C_2 = \langle \sigma \rangle$ ,  $C_1 = \langle e_1, e_2, e_3 \rangle$ ,  $C_0 = \langle v_0 \rangle$ . Let  $n \in \mathbb{N}$ .  $\partial_n$  is defined such that  $\partial_n(\sigma_\alpha) = \sum_i (-1)^i \sigma_\alpha \mid [v_0, \cdots, \hat{v_i}, \cdots, v_n]$ . Since there is only one vertex,  $\partial_n$  is the zero map.

This argument doesn't work. Check the torus example from class. It only has one vertex, but  $\partial_n$  is not the zero map for some n.

Therefore,  $H_n = \ker(\partial_n)/\operatorname{Im}(\partial_{n+1}) = C_n/\langle 0 \rangle = C_n$ . Thus

$$H_n = \begin{cases} \{0\} & (n \ge 3) \\ \langle \sigma \rangle \cong \mathbb{Z} & (n = 2) \\ \langle e_1, e_2, e_3 \rangle \cong \mathbb{Z}^3 & (n = 1) \\ \langle v_0 \rangle \cong \mathbb{Z} & (n = 0). \end{cases}$$




Figure 1. mycaption




FIGURE 2. Problem 5

## I'm not sure if this is correct.

**Exercise.** (Problem 5) Compute the simplicial homology groups of the Klein bottle using the  $\Delta$ -complex structure described at the beginning of this section.

*Proof.* We will use the notations in Figure 2.

$$C_n = \begin{cases} 0 & (n \ge 3) \\ \langle U, L \rangle & (n = 2) \\ \langle a, b, c \rangle & (n = 1) \\ \langle v \rangle & (n = 0). \end{cases}$$

 $\partial_n = 0$  for  $n \ge 3$  and n = 0.

$$\partial_2(U) = \sum_{i=0}^{2} (-1)^i \sigma \mid [0, 1, 2]$$
  
=  $\sigma \mid [1, 2] - \sigma \mid [0, 2] + \sigma \mid [0, 1]$   
=  $b - c + a$ .

$$\partial_2(L) = \sum_{i=0}^2 (-1)^i \sigma \mid [0, 1, 2]$$
  
=  $\sigma \mid [1, 2] - \sigma \mid [0, 2] + \sigma \mid [0, 1]$   
=  $a - b + c$ .