

MATH 601 (DUE 11/22)

HIDENORI SHINOHARA

CONTENTS

1. THE THEOREM ON SYMMETRIC POLYNOMIALS	1
2. Galois Theory VI	1
3. Galois Theory V(Further exercises)	3
4. Galois Theory V	3

1. THE THEOREM ON SYMMETRIC POLYNOMIALS

Exercise. (Problem 1) By substituting $u_4 = 0$, we get $u_1^2 u_2 u_3 + u_1 u_2^2 u_3 + u_1 u_2 u_3^2 = s_3 s_1$. $s_3 s_1$ with 4 variables expands to $u_1^2 u_2 u_3 + u_1^2 u_2 u_4 + u_1^2 u_3 u_4 + u_1 u_2^2 u_3 + u_1 u_2^2 u_4 + u_1 u_2 u_3^2 + 4u_1 u_2 u_3 u_4 + u_1 u_2 u_4^2 + u_1 u_3^2 u_4 + u_1 u_3 u_4^2 + u_2^2 u_3 u_4 + u_2 u_3^2 u_4 + u_2 u_3 u_4^2$. Then $s_3 s_1 - f$ where f is the original polynomial gives us $4u_1 u_2 u_3 u_4 = 4s_4$. Therefore, $f = s_3 s_1 - 4s_4$.

Exercise. (Problem 2) We are given that $|M - xI| = x^3 - ax^2 + bx - c$. This implies that $|M - (-x)I| = -x^3 - ax^2 - bx - c$. Since the determinant function preserves multiplication, $|M - xI||M - (-x)I| = |M^2 - x^2I|$. This implies $|M^2 - x^2I| = -x^6 + (a^2 - 2b)x^4 + (-b^2 + 2ac)x^2 + c^2$. Therefore, the characteristic polynomial of M is $-x^3 + (a^2 - 2b)x^2 + (-b^2 + 2ac)x + c^2$.

$|xI - M|$ is what I need to calculate.

2. GALOIS THEORY VI

Exercise. (Problem 3)

- $\{(123), (132), e\}$ is clearly a subgroup of the stabilizer group S_v of v . Since $(12) \notin S_v$, $3 \leq |S_v| \leq 5$. By Lagrange's Theorem, $S_v = \langle (123) \rangle$.
- By (i), $S_3 v$ contains only $[S_3 : S_v] = 2$ elements. Thus $v' = (12) \cdot v = u_2 u_1^2 + u_1 u_3^2 + u_3 u_2^2$.
- By substituting $u_3 = 0$ for $v + v'$, we get $u_1 u_2^2 + u_2 u_1^2 = s_1 s_2$. Then $v + v' - s_1 s_2 = -3u_1 u_2 u_3 = -3s_3$. Therefore, $v + v' = s_1 s_2 - 3s_3$.
- We will use the fundamental theorem of Galois Theory. $F(v) = K^{\langle (123) \rangle}$, so $|\langle (123) \rangle| = 3 = [K : F(v)]$. Moreover, $|\langle \text{Gal}(K/F) \rangle| = [K : F]$. Therefore, $[F(v) : F] = [K : F]/[K : F(v)] = |\text{Gal}(K/F)|/3$.
- Calculation shows that $vv' = 9s_3^2 + s_3 s_1^3 - 6s_3 s_1 s_2 + s_2^3$. By substituting $s_1 = 0, s_2 = p, s_3 = q$, we get $9q^2 + p^3$.
- Since A_3 is the only proper transitive subgroup of S_3 , $\text{Gal}(K/F) = S_3$ if and only if $\sigma \in \text{Gal}(K/F)$ where σ corresponds to the permutation (12) . (i.e., $u_1 \mapsto u_2, u_2 \mapsto u_1$.)

v, v' are not fixed by σ , so $v, v' \notin F$ if $\text{Gal}(K/F) = S_3$. v, v' are fixed by every permutation if $\text{Gal}(K/F) = A_3$ because it is generated by σ' that corresponds to (123). Therefore, we can conclude that $\text{Gal}(K/F) \neq S_3$ if and only if $v, v' \in F$. $v, v' \in F$ if and only if $(y-v)(y-v')$ factors in F . Therefore, $h(y) = y^2 - (v+v')y + vv' = y^2 + 3qy + (9q^2 + p^3)$ is the desired polynomial.

(vii) The discriminant is $(3q)^2 - 4(9q^2 + p^3) = 9q^2 - 36q^2 - 4p^3 = -27q^2 - 4p^3$.

Exercise. (Problem 4)

- (i) The discriminant can be expressed as $-4s_1^3s_3 + s_1^2s_2^2 + 18s_1s_2s_3 - 4s_2^3 - 27s_3^2$. By substituting $s_1 = 1, s_2 = -2, s_3 = -1$, we get 49.

```
from sympy.polys.polyfuncs import symmetrize
from sympy import *
```

```
u1, u2, u3 = symbols('u1_u2_u3')
```

```
u = [u1, u2, u3]
```

```
discriminant = 1
```

```
for i in range(3):
```

```
    for j in range(i + 1, 3):
```

```
        discriminant *= (u[i] - u[j]) * (u[i] + u[j])
```

```
print(latex(symmetrize(discriminant, formal = True)[0]))
```

Since the discriminant is a square, the Galois group is isomorphic to A_3 .

Exercise. (Problem 5)

(i)

(ii) $x^4 + x + 1$ is irreducible because

- It does not have a linear factor by the rational root theorem.
- If it factors into two rational quadratic polynomials, they will factor into two monic integer quadratic polynomials, namely, $x^2 + ax + b$ and $x^2 - ax + 1/b$ based on the coefficients. This implies $b = \pm 1$. Since the coefficient of x is 1, $-ab + a/b = 1$, but this implies $b \neq \pm 1$.

We will use the discussion presented in the Galois Theory IV handout. By (i), the discriminant is 229, so $h(y) = y^2 - 229$. Also, $g(y) = y^3 - 4y - 1$ since $a = b = 0, c = -1, d = 1$. Therefore, both $h(y)$ and $g(y)$ are irreducible, so the Galois group is S_4 .

- (iii) It does not have a linear factor by the rational root theorem. Based on coefficients, if it factors into quadratic polynomials, it will be $(x^2 + ax + b)(x^2 - ax + c)$ for some $a, b, c \in \mathbb{Z}$ by Gauss' lemma. This gives $bc = 12$ and $-ab + ac = -8$, so $a(c - 12/c) = -8$. This is a quadratic polynomial in c with the discriminant $64 - 48a$. This must be a square for c to exist. By checking each possible value of a , we get $64 - 48 \cdot -8 = 448, 64 - 48 \cdot -4 = 256, 64 - 48 \cdot -2 = 160, 64 - 48 \cdot -1 = 112, 64 - 48 \cdot 1 = 16$. (For other a , $64 - 48a < 0$.) Thus the only two possible values are $a = 1, -4$. $a = 1$ gives $c - b = -8$ and $bc = 12$, which we can confirm to be impossible by




FIGURE 1. Subgroups

examining the divisors of 12. Similarly, $a = -4$ gives $c - b = 2$ and $bc = 12$ and this is impossible to satisfy. Therefore, $x^4 - 8x + 12$ is irreducible over \mathbb{Q} .

- (iv) Again, we will use the discussion presented in the Galois Theory IV handout. By calculating the discriminant, we have $h(y) = h(y) = y^2 - 331776$ and $g(y) = y^3 - 48y - 64$. $h(y)$ factors as $576^2 = 331776$. $g(y)$ does not factor by the rational root theorem. Therefore, the Galois group is A_4 .

3. GALOIS THEORY V(FURTHER EXERCISES)

Exercise. (Problem 3)

- (i)

4. GALOIS THEORY V

Exercise. (Problem 1)

- (i) D_4 is a subgroup of S_4 and the Galois Theory V handout states that S_4 is solvable and any subgroup of a finite solvable group is solvable.
- (ii) If S_5 is solvable, $A_5 \leq S_5$ is solvable. However, as stated in the handout, A_5 is not solvable.

Exercise. (Problem 2)

- (i) Let $a = (12345)$ and $b = (2354)$. Since $|a| = 5$ and $|b| = 4$, $20 \mid |G|$. On the other hand, every element can be written as $a^i b^j$ because of the relation $ab = (1243) = ba^3$. Since every element can be written as $a^i b^j$ where $0 \leq i \leq 4$ and $0 \leq j \leq 3$, G contains exactly 20 elements.
- (ii) G has a subgroup $G_1 = \{(1), (12)(35), (12345), (13)(45), (13524), (14)(23), (14253), (15)(24), (15432), (25)(34)\}$ of index 2. G_1 has a subgroup $G_2 = \{(1), (12345), (13524), (14253), (15432)\}$ of index 2. G_2 is abelian, so we can pick $G_3 = \{(1)\}$. Then $G_3 \subset G_2 \subset G_1 \subset G$ is a filtration.

Exercise. (Problem 3) Figure 1 shows all the subgroups of G where $a = (1325), b = (1435), c = (1243), d = (1254), e = (2354)$. By the fundamental theorem of Galois theory,

there is a bijective correspondence between intermediate groups and intermediate fields, and the inclusion is reversed. The degree of extension corresponds to the index. By Lagrange's theorem, it is clear that, between any two groups connected by an edge in the figure, there cannot be any proper subgroups between them.

The number of subgroups of order 4 and 5 must be 5 and 1 by Sylow's theorem. By writing out all the 20 elements

$$(1), (12)(35), (12345), (1243), (1254), (13)(45), (1325), (1342), (13524), (14)(23), \\ (14253), (1435), (1452), (15)(24), (1523), (1534), (15432), (2354), (2453), (25)(34),$$

we conclude that the number of elements of order 2 is 5. Thus the number of subgroups of order 2 is 5. Finally, $G/\langle(12345)\rangle$ is isomorphic to $\mathbb{Z}/4$. Therefore, there can be only one subgroup of order 10 between G and $\langle(12345)\rangle$. Conversely, since every subgroup of order 10 must contain a subgroup of order 5, $\langle(12345), a^2\rangle$ is the only subgroup of order 10.