

MATH 611 (DUE 11/20)

HIDENORI SHINOHARA

Exercise. (Problem 1)

Exercise. (Problem 28 (a)) Let A, B be the Mobius strip and a torus with a small neighborhood around them so the strip and torus are contained in A and B . For any $n \geq 3$, the exact sequence $H_n(A \cap B) \rightarrow H_n(A) \oplus H_n(B) \rightarrow H_n(X) \rightarrow H_{n-1}(A \cap B)$ implies that $H_n(X) \cong H_n(A) \oplus H_n(B) = 0 \oplus 0 = 0$ because the intersection $A \cap B$ is homotopic to S^1 , so $H_n(A \cap B) = H_{n-1}(A \cap B) = 0$. $H_0(X) =$

We will examine the LES

$$\tilde{H}_2(A \cap B) \rightarrow \tilde{H}_2(A) \oplus \tilde{H}_2(B) \xrightarrow{f_1} \tilde{H}_2(X) \xrightarrow{f_2} \tilde{H}_1(A \cap B) \xrightarrow{f_3} \tilde{H}_1(A) \oplus \tilde{H}_1(B) \xrightarrow{f_4} \tilde{H}_1(X) \rightarrow \tilde{H}_0(A \cap B).$$

- Since $\tilde{H}_2(A \cap B) = 0$, so f_1 is injective.
- $\tilde{H}_1(A \cap B) = \mathbb{Z}$, and $f_3(1) = (2, (1, 0))$ because the intersection goes around the mobius strip twice while it only goes around the torus once. Then f_3 is injective, so $\text{Im}(f_2) = \ker(f_3) = 0$. This implies that $(f_1) = \ker(f_2) = H_2$, so f_1 is surjective.

Therefore, f_1 is bijective, so $H_2(X) = \tilde{H}_2(X) = \tilde{H}_2(A) \oplus \tilde{H}_2(B) = 0 \oplus \mathbb{Z} = \mathbb{Z}$.

Finally, f_4 's surjectivity implies that

$$\begin{aligned} \tilde{H}_1(X) &\cong \tilde{H}_1(A) \oplus \tilde{H}_1(B) / \ker(f_4) \\ &= \mathbb{Z} \oplus \mathbb{Z}^2 / \langle (2, (1, 0)) \rangle \\ &\cong \langle a, b, c \rangle / \langle 2a + b \rangle \\ &\cong \langle a, b, c \mid 2a + b \rangle \\ &\cong \langle a, -2a, c \rangle \\ &\cong \langle a, c \rangle = \mathbb{Z} \oplus \mathbb{Z}. \end{aligned}$$

Thus $H_1(X) = \mathbb{Z} \oplus \mathbb{Z}$.

Exercise. (Problem 28 (b))

Exercise. (Problem 29)