

# MATH 601 (DUE 10/23)

HIDENORI SHINOHARA

## CONTENTS

### 1. Field Extension

1

#### 1. FIELD EXTENSION

**Exercise.** (Problem 1) Let  $p$  be a prime number. Let  $K = \mathbb{Z}/p\mathbb{Z}(t)$  be the fraction field of  $\mathbb{Z}/p\mathbb{Z}[t]$ .

- (i) What is the characteristic of  $K$ ?
- (ii) What is the characteristic of any extension field of  $K$ ?
- (iii) Show that the Frobenius endomorphism,  $F : K \rightarrow K$  is not a ring isomorphism.
- (iv) Let  $f(x) = x^p - t \in K[x]$ . Prove that  $f(x)$  is irreducible.
- (v) Prove that  $f(x)$  is not a separable polynomial.
- (vi) Construct an explicit field extension  $K \subset L$  such that  $f(x) \in L[x]$  has a factor of positive degree  $< p$ .
- (vii) With  $f$  and  $L$  above find all the roots of  $f(x)$  in  $L$  and determine their multiplicities.

*Proof.*

- (i) We will write  $k \cdot 1$  to denote  $1 + 1 + \cdots + 1$  ( $k$  times). Since  $p \cdot 1 = 0$  in  $K$ , the characteristic of  $K$  is at most  $p$ . Let  $k$  denote the characteristic of  $K$ . Let  $i : \mathbb{Z}/p\mathbb{Z} \rightarrow (\mathbb{Z}/p\mathbb{Z})[t], i' : \mathbb{Z}/p\mathbb{Z}[t] \rightarrow K$  be inclusions. Then  $i' \circ i : \mathbb{Z}/p\mathbb{Z} \rightarrow K$  is an injective ring homomorphism.  $k \cdot 1 \neq 0$  in  $\mathbb{Z}/p\mathbb{Z}$ . Thus  $(i' \circ i)(k \cdot 1) = k \cdot (i' \circ i)(1) = k' \cdot 1 = 0$ . Since  $i' \circ i$  is injective, this implies  $k \cdot 1 = 0$ . Therefore,  $k \geq p$ , so  $k$  must be equal to  $p$ .

□

**Exercise.** (Problem 2) Let  $F$  be a field of characteristic 0. Let  $f(x) \in F[x]$  be an irreducible polynomial. Then  $f(x)$  is separable.

*Proof.* Let  $f(x) = \sum_{i=0}^n a_i x^i \in F[x]$  be an irreducible polynomial with  $a_n \neq 0$ . Since  $f(x)$  is irreducible,  $f(x)$  is neither a unit nor 0. Since  $F$  is a field, all polynomials of degree 0 are units. Thus  $\deg(f(x)) = n \geq 1$ . It suffices to show that  $\text{GCD}(f(x), f'(x)) = F^*$  by Lemma 3.2. Let  $g(x) \in F[x]$  be given such that  $g(x) \mid f(x), g(x) \mid f'(x)$ . Since  $f(x)$  is irreducible, either  $g(x)$  is a unit or there exists a unit  $u \in F^*$  such that  $g(x) = uf(x)$ . Suppose  $g(x)$  is not a unit. Since  $g(x) \mid f'(x)$ ,  $f'(x) = h(x)g(x) = uh(x)f(x)$  for some  $h(x) \in F[x]$ . Thus  $\deg(f'(x)) = \deg(uh(x)) + \deg(f(x))$ .

- $f'(x) = \sum_{i=1}^n i a_i x^{i-1}, n \geq 1$  and  $a_n \neq 0$ . Since  $F$  is a field of characteristic 0,  $na_n \neq 0$ . Therefore,  $\deg(f'(x)) = n - 1$ .

- $\deg(gh(x)) \geq 0$ .
- $\deg(f(x)) = n$ .

However, this implies that  $n - 1 \geq 0 + n = n$ . This is a contradiction, so  $g(x)$  must be a unit. Therefore,  $\text{GCD}(f(x), f'(x)) = F^*$ .  $\square$

**Exercise.** (Problem 3) Let  $F$  be a field. Let  $f(x) \in F[x]$  be an irreducible polynomial which is not separable. Show that  $f'(x) = 0 \in F[x]$ .

*Proof.* Suppose  $f(x)$  is irreducible. Then  $f(x) \neq 0$  and  $f(x)$  is not a unit by definition. Thus  $\deg(f(x)) \geq 1$ .

Since  $f(x)$  is not separable, there exists a non-unit  $g(x) \in F[x]$  such that  $g(x) \mid f(x)$  and  $g(x) \mid f'(x)$  by Lemma 3.2 from the Field Extension handout. Since  $f(x)$  is irreducible and  $g(x)$  is not a unit,  $f(x)$  is the product of  $g(x)$  and a unit. This implies that  $\deg(f(x)) = \deg(g(x))$ .

Since  $g(x) \mid f'(x)$ ,  $f'(x) = h(x)g(x)$ . If  $f'(x) = 0$ , we are done. Suppose otherwise. Then  $\deg(f'(x)) = \deg(h(x)) + \deg(g(x)) = \deg(h(x)) + \deg(f(x)) \geq \deg(f(x))$ . However, by the definition of the  $'$  operator,  $\deg(f'(x)) < \deg(f(x))$ . This is a contradiction, so  $f'(x) = 0$ .  $\square$

**Exercise.** (Problem 4) Let  $F$  be a field of prime characteristic  $p$ . Let  $f(x) = \sum_{i=0}^n a_i x^i \in F[x]$  be an irreducible polynomial. Give a necessary and sufficient criterion for  $f(x)$  to be inseparable in terms of the coefficients  $a_i$ .

*Proof.* We claim that  $\forall i, (i \notin p\mathbb{Z} \implies a_i = 0)$  is a necessary and sufficient criterion.

- Suppose  $f(x)$  is inseparable. By Lemma 5.5 from the Field Extension handout,  $f'(x) = 0$ . If  $f'(x) = 0$ , then  $ia_i = 0$  for each  $i$ . Since  $p$  is a prime,  $a_i$  must be 0 if  $i \notin p\mathbb{Z}$ .
- Suppose  $\forall i, (i \notin p\mathbb{Z} \implies a_i = 0)$ . Then  $f'(x) = 0$ , so  $f(x) \mid f(x), f(x) \mid f'(x)$  and  $f(x)$  is not a unit since  $f(x)$  is irreducible. Therefore,  $\text{GCD}(f(x), f'(x)) \neq F^\times$ , so  $f$  is inseparable by Lemma 3.2.

Hence,  $\forall i, (i \notin p\mathbb{Z} \implies a_i = 0)$  is a necessary and sufficient criterion.  $\square$

**Exercise.** (Problem 5) What is the characteristic of the ring  $\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$ ?

*Proof.* Define  $\phi : \mathbb{Z} \rightarrow \mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$  such that  $\phi(k) = (k, 0, 0)$ . Then  $\phi$  is injective, so the characteristic is 0.  $\square$

**Exercise.** (Problem 6) Let  $K$  be a finite field of characteristic  $p$ . Let  $a, b \in K^*$  be two elements which have the same order in this finite group. Show that  $\mathbb{Z}/p[a] = \mathbb{Z}/p[b]$  as subfields of  $K$ .

*Proof.*

Can I just say they both have the same number of elements and use the lemma from class?

$\square$