

MATH 633(HOMEWORK 7)

HIDENORI SHINOHARA

Exercise. (1) Suppose f is locally bijective. For every $p \in U$, f is locally injective. Therefore, f' is nonzero in the neighborhood around p . In other words, f' is nonzero on U .

The other direction

Exercise. (10) Let $\sigma(z) = -i(z+1)/(z-1)$. Then σ sends the unit disk to the upper half-plane with ∞ since $\sigma(a+bi) = (-2b - (a^2 + b^2 - 1)i)/((a-1)^2 + b^2)$. On the other hand, $\sigma^{-1} : z \mapsto (z-i)/(z+i)$ sends the upper half plane with ∞ to the unit disk because $|a + (b-1)i| \leq |a + (b+1)i|$ if $b \geq 0$. Therefore, σ is a bijection between the unit disk and $H \cup \{\infty\}$. $F \circ \sigma$ sends the unit disk to the unit disk, and $F(\sigma(0)) = 0$. By Lemma 2.1, $|(F \circ \sigma)(w)| \leq |w|$ for every $w \in D$. Then for every $z \in \mathbb{H}$, $\sigma^{-1}(z) \in D$. Then $|F(z)| = |(F \circ \sigma)(\sigma^{-1}(z))| \leq |\sigma^{-1}(z)| = |(z-i)/(z+i)|$, which is the desired result.

Exercise. (12(a)) Let $a \neq b$ be two fixed points. Let $\sigma(z) = (z-a)/(1-\bar{a}z)$. Then σ sends a to 0 and maps D to D bijectively. Let $g = \sigma \circ f \circ \sigma^{-1}$. g has two fixed points, 0 and $\sigma(b)$. By applying Lemma 2.1, g is a rotation. However, g fixes $\sigma(b) \neq 0$, so g must be the identity map. Then f must be the identity.

Exercise. (12(b)) The map $\sigma : z \mapsto (z-i)/(z+i)$ maps the upper half-plane to the unit disk bijectively. Then $\sigma \circ f \circ \sigma^{-1}$ where $f(z) = z+1$ is a holomorphic bijection on f that has no fixed point because f has no fixed point.

Exercise. (16(a)) The composition of mobius transformations corresponds to the multiplication of the corresponding matrices. Thus it suffices to calculate

$$\begin{aligned} \begin{bmatrix} -1 & i \\ 1 & i \end{bmatrix}^{-1} \begin{bmatrix} e^{i\theta} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & i \\ 1 & i \end{bmatrix} &= \frac{1}{2} \begin{bmatrix} e^{i\theta} + 1 & -i(e^{i\theta} - 1) \\ i(e^{i\theta} - 1) & e^{i\theta} + 1 \end{bmatrix} \\ &\rightarrow \frac{1}{2} \begin{bmatrix} 1 & -i(e^{i\theta} - 1)/(e^{i\theta} + 1) \\ i(e^{i\theta} - 1)/(e^{i\theta} + 1) & 1 \end{bmatrix} \\ &\rightarrow \frac{1}{2} \begin{bmatrix} 1 & -i(i \tan(\theta/2)) \\ i(i \tan(\theta/2)) & 1 \end{bmatrix} \\ &\rightarrow \frac{1}{2} \begin{bmatrix} 1 & \tan(\theta/2) \\ -\tan(\theta/2) & 1 \end{bmatrix}. \end{aligned}$$

Thus the answer is the mobius transformation associated to the last matrix.

Exercise. (16(b)) Let $\alpha = a + bi$.

$$\begin{aligned} \begin{bmatrix} -1 & i \\ 1 & i \end{bmatrix}^{-1} \begin{bmatrix} -1 & \alpha \\ -\bar{\alpha} & 1 \end{bmatrix} \begin{bmatrix} -1 & i \\ 1 & i \end{bmatrix} &= \frac{1}{2} \begin{bmatrix} \bar{\alpha} - \alpha & -i(\alpha + \bar{\alpha} - 2) \\ -i(\alpha + \bar{\alpha} + 2) & \alpha - \bar{\alpha} \end{bmatrix} \\ &\rightarrow \begin{bmatrix} b & a-1 \\ a+1 & -b \end{bmatrix}. \end{aligned}$$

After multiplying $1/(1 - a^2 - b^2)$ to every term, we obtain a matrix associated to the desired mobius transformation.