## MATH 602 HOMEWORK 2

## HIDENORI SHINOHARA

**Exercise.** (Problem 1) We will assume that the problem meant to say "su with  $s \in S \setminus \{0\}$ " because it would be trivial otherwise. Choose  $a_{n-1}, \dots, a_0 \in R$  such that  $u^n + a_{n-1}u^{n-1} + \dots + a_1u^1 + a_0 = 0$ . If  $a_0 = 0$ , then  $u(u^{n-1} + a_{n-1}u^{n-2} + \dots + a_1) = 0$ . Since we are dealing with integral domains, this implies  $u^{n-1} + a_{n-1}u^{n-2} + \dots + a_1 = 0$ . By repeating this process, we obtain a monic polynomial with coefficients in R and a nonzero constant term that u satisfies.

Therefore, we may assume  $a_0 \neq 0$ . Then  $u(a_1 + a_2u + \cdots + a_{n-1}u^{n-2} + u^{n-1}) = -a_0 \in R$ . Since  $a_0 \neq 0$ ,  $a_1 + a_2u + \cdots + a_{n-1}u^{n-2} + u^{n-1}$  is a nonzero element in S. Hence, we showed that some multiple of u lives in R.

**Exercise.** (Problem 2) Let  $R = \mathbb{Z}$  and  $S = 2\mathbb{Z}$ .  $R \setminus S$  is not even an ideal because  $0 \notin R \setminus S$ . Thus  $R \setminus S$  is not a prime ideal.

Exercise. (Problem 3)

## Solve this!

**Exercise.** (Problem 4) Let  $p \in \operatorname{Spec}(R)$  such that  $I \subset p$ . Define  $p/I = \{x+I \mid x \in p\} \subset R/I$ . By the third isomorphism theorem of rings, p/I is an ideal of R/I. Let  $x+I, y+I \in R/I$  and suppose  $(x+I)(y+I) \in p/I$ . Then xy+I=z+I for some  $z \in p$ . Thus  $xy-z \in I \subset p$  and  $z \in p$ . Thus  $xy \in p$ , so  $x \in p$  or  $y \in p$ . This implies  $x+I \in p/I$  or  $y+I \in p/I$ , so  $p/I \in \operatorname{Spec}(R/I)$ .

On the other hand, let  $A/I \subset \operatorname{Spec}(R/I)$  be given. By the third isomorphism theorem of rings, every ideal of R/I must be of the form A/I where A is an ideal of R that contains I. Let  $x, y \in R$  and suppose  $xy \in A$ . Then  $xy + I \in A/I$ , so  $(x + I)(y + I) \in A/I$ . Without loss of generality,  $x + I \in A/I$ . Then x + I = a + I for some  $a \in A$ . Thus  $x - a \in I \subset A$  and  $a \in A$ , so  $x \in A$ . Therefore, A is a prime ideal of R containing I.

**Exercise.** (Problem 5) By the second isomorphism of rings,  $R \cap q$  is an ideal in R. Let  $x, y \in R$ . Suppose  $xy \in R \cap q$ . Then  $xy \in q$ , so  $x \in q$  or  $y \in q$ . Without loss of generality,  $x \in q$ . Then  $x \in R \cap q$ . Thus  $R \cap q$  is a prime ideal of R.