

MATH 612 (HOMEWORK 2)

HIDENORI SHINOHARA

Exercise. (Exercise 1) Fix G and let $\alpha : H \rightarrow H'$ be given. Let $0 \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H \rightarrow 0, 0 \rightarrow G_1 \xrightarrow{g_1} G_0 \xrightarrow{g_0} H \rightarrow 0$ be free resolutions. By Lemma 3.1(a), we obtain two homomorphisms $\alpha_1 : F_1 \rightarrow G_1, \alpha_0 : F_0 \rightarrow G_0$ which commutes with f_i, g_i, α . Then we obtain two chain complexes

$$\begin{aligned} 0 \leftarrow \text{Hom}(F_1, G) \xleftarrow{f_1^*} \text{Hom}(F_0, G) \xleftarrow{f_0^*} \text{Hom}(H, G) \leftarrow 0 \\ 0 \leftarrow \text{Hom}(F_1, G') \xleftarrow{f_1^*} \text{Hom}(F_0, G') \xleftarrow{f_0^*} \text{Hom}(H, G') \leftarrow 0. \end{aligned}$$

with induced maps $\alpha_1^*, \alpha_0^*, \alpha^*$ forming a chain map from the chain complex on the bottom to the one on the top. Then α_1^* induces a map from $\text{Ext}(H', G) \rightarrow \text{Ext}(H, G)$.

Fix H and let $f : G \rightarrow G'$ be given. Let $0 \rightarrow F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} H \rightarrow 0$ be a free resolution of H . We obtain two cochain complexes where f_* is a chain map from the top one to the bottom one.

$$\begin{aligned} 0 \leftarrow \text{Hom}(F_1, G) \xleftarrow{f_1^*} \text{Hom}(F_0, G) \xleftarrow{f_0^*} \text{Hom}(H, G) \leftarrow 0 \\ 0 \leftarrow \text{Hom}(F_1, G') \xleftarrow{f_1^*} \text{Hom}(F_0, G') \xleftarrow{f_0^*} \text{Hom}(H, G') \leftarrow 0. \end{aligned}$$

f_* indeed makes the diagram commute because for any $\sigma \in \text{Hom}(H, G)$,

$$\begin{aligned} f_*(f_0^*(\sigma)) &= f_*(\sigma \circ f_0) \\ &= f \circ (\sigma \circ f_0) \\ &= (f \circ \sigma) \circ f_0 \\ &= f_0^*(f \circ \sigma) \\ &= f_0^*(f_*(\sigma)). \end{aligned}$$

Similarly, $f_*(f_1^*(\sigma)) = f_1^*(f_*(\sigma))$ for every $\sigma \in \text{Hom}(F_0, G)$. Since a chain map induces a homomorphism on cohomology groups, f induces a map from $\text{Ext}(H, G) \rightarrow \text{Ext}(H, G')$.