


Salivary glands




Drake: Gray's Anatomy for Students, 2nd Edition.
Copyright © 2009 by Churchill Livingstone, an imprint of Elsevier, Inc. All rights reserved.

Major salivary glands:


- Parotid
- Submandibular
- Sublingual

Minor salivary glands:

They are 800-1000 glands located throughout the oral cavity within the submucosa.




Submandibular gland




- located in the *submandibular trigone*
- the *facial artery* is embedded to its medial/internal side
- the *facial vein* attaches to its external side
- the *lingual nerve* is presented at its superior side
- it is supplied by the lingual artery and innervated by the submandibular ganglion via lingual nerve

Submandibular gland




- the *submandibular duct* arises from its medial side and enters to the lateral lingual groove

Submandibular gland




- the *submandibular ducts* open onto the surface of the sublingual caruncles on either side of the lingual frenulum

Lateral lingual sulcus




- V-shaped groove
- medial wall: hyoglossus
- lateral wall: mylohyoid
- contents from superior to inferior: lingual nerve, submandibular duct, hypoglossal nerve

Sublingual gland




- located on the *upper side of the mylohyoid anterior to the submandibular gland*
- their multiple small ducts empty into the submandibular duct as well as directly onto the floor of the mouth
- it is supplied by the sublingual artery and innervated by the submandibular ganglion via lingual nerve

Sublingual gland




- located on the *upper side of the mylohyoid anterior to the submandibular gland*
- their multiple small ducts empty into the submandibular gland as well as directly onto the floor of the mouth
- it is supplied by the sublingual artery and innervated by the submandibular ganglion via lingual nerve

Parotid gland



- it is located in the *parotid nest*
- covered by the *parotid fascia*
- *parotid duct* travels underneath the zygomatic arch and at the anterior edge of the masseter turns to medial and pierces the buccinator then enters to the *vestibule of the oral cavity at the level of second molar tooth*

Parotid gland



- supplied by the *external carotid artery* and drained by the *external jugular vein*
- the *facial* and *auriculotemporal nerves* run through the gland
- the *auriculotemporal nerve* provides secretomotor (parasympathetic) innervation for it