



Jejunum and ileum




- **intraperitoneal** structures
- their total length: 3-5 m
- their form *intestinal loops*
- only the terminal ileum has relatively constant position
- *there is no macroscopic landmark between the two parts*
- *circular mucosal folds internally*


Ileocecal valve (Bauhin's valve)



The partly digested food, the chyme, flows into the cecum through the ileocecal valve. It prevents the reflux from the cecum.



Radix mesenterii (root of mesentery)



Mesentery: peritoneal duplication suspending the jejunum and ileum.

Root of mesentery: The attachment of the mesentery *to the posterior abdominal wall, extending from the duodenojejunal flexure to the ileocecal junction.*

Neurovasculature of the duodenum

Arterial supply:

- *superior anterior pancreaticoduodenal artery*
- *superior posterior pancreaticoduodenal artery*
- *inferior pancreaticoduodenal artery*

Venous drainage:

into the hepatic portal vein


Lymphatic drainage:


pyloric, hepatic and superior mesenteric lymph nodes (into the celiac lymph nodes)

Innervation:

parasympathetic: vagus nerve

sympathetic: splanchnic nerves





Neurovasculature of the jejunum and ileum

- jejunal arteries
- ileal arteries
- ileocolic artery

Venous drainage:

into the hepatic portal vein via superior mesenteric vein


Lymphatic drainage:

superior mesenteric lymph nodes

Innervation:

parasympathetic: vagus nerve


sympathetic: splanchnic nerves



Neurovasculature of the jejunum and ileum

Jejunum:

- 1-2 series of arcades from the jejunal arteries
- the vessels enter to the intestinal wall at the **mesenteric side**



Ileum:

- 2-3 series of arcades from the ileal arteries
- the vessels enter to the intestinal wall at the **mesenteric side**

Meckel's diverticulum



- *slight bulge in the terminal ileum that results from the failure of the vitelline duct to obliterate*
- *it is considered the most common structural congenital anomaly of the gastrointestinal tract (2% of the population)*
- *usually, it is asymptomatic; most patients with a complication are thought to present in the first two years of life*

Perforated Meckel's diverticulum, Case courtesy of Dr Eric F Greif, Radiopaedia.org, rID: 34130