



# Auscultation



Auscultation is a method used to listen to sounds of the body during a physical examination



Anterior View

# HEART, SCELETOTOPIA

contour:

- apex: left 5th intercostal space, 9 cm far from the midline
- superior vena cava entrance: right 3rd rib (cartilage) sternal end
- right edge of the sulcus coronarius: on the right 6th rib (cartilage), 2 cm far from its sternal end
- left edge of the sulcus coronarius: on the left 3rd rib (cartilage), 3 cm far from its sternal end

absolute cardiac dullness: beside the sternum, on the left side, between the 4th and 6th ribs, an area similar in size to the palm of a child

relative cardiac dullness (percussion of the heart):

- upper margin is in the 3rd intercostal space
- left margin is 1 cm medial from the midclavicular line
- right margin doesn't go beyond the sternum
- lower margin is at the level of the 5th rib

ostium:

- ostium of truncus pulmonalis: left 3rd rib cartilage sternal end
- ostium of aorta: behind and right to the former
- tricuspid valve: between the sternal ends of the 5th rib cartilages, in the middle
- bicuspid valve: left 4th rib cartilage sternal end

auscultation:

- valve of truncus pulmonalis: left 2nd intercostal space sternal end
- valve of aorta: right 2nd intercostal space sternal end
- tricuspidal valve: right to the lower edge of the sternum
- bicuspidal valve: on the line between the projection of the apex of the heart and the left 4th rib cartilage sternal end

# AUSCULTATION POINTS

