

and then observing that the canonical function of Y writes as $k(x) = \lambda e^{Ax-B|x|}$, implying $\tau_+^{-A} = \tau_-^A = \lambda \delta_B$. The Dirac integration to which (5.25) reduces, produces

$$k_\rho(x) = \lambda e^{(A^2-B^2)x/2} = \lambda e^{-\theta_+ \theta_- x/2}. \quad (5.27)$$

Therefore, the bilateral gamma process can be represented in law as $Y_t = W_{T_t} + (\theta_- - \theta_+)t/2$ where $\mathcal{L}(T_1) \sim \Gamma(\lambda, \theta_+ \theta_-/2)$. This has been known and used at least since Madan et al. (1998).

Example 10. *Subordinated representation of CTS_α processes.* Madan and Yor (2008) through a series of propositions covering a substantial part of the article, explain how to represent a $\text{CTS}_\alpha(\mathbb{R})$ process as a subordinated Brownian motion with drift. Based on the discussion so far, the main Proposition 2 therein trivialize. First of all notice that the assumptions in Theorem 6 are met also in the case of $\text{CTS}_\alpha(\mathbb{R})$ processes with symmetric spherical part. Namely, assuming $\lambda_+ = \lambda_-$ and letting again $A = (\theta_- - \theta_+)/2$, $B = (\theta_- + \theta_-)/2$, leads to $k(x) = x^{-\alpha} e^{Ax-B|x|}$, so that using the Thorin measure found in Subsection 4.3 we deduce the required density relationship $\tau_+^{-A}(y) = \tau_-^A(y) = \lambda(y-B)^{\alpha-1}(\Gamma(\alpha))^{-1} \mathbf{1}_{\{y \geq B\}}$. It then follows from Corollary 6, and with the substitution $w = (s-B)\sqrt{x}$, that

$$\begin{aligned} k_\rho(x) &= \lambda \frac{e^{xA^2/2}}{\Gamma(\alpha)} \int_B^\infty e^{-\frac{s^2x}{2}} (s-B)^{\alpha-1} ds = \lambda \frac{e^{(A^2-B^2)x/2}}{\Gamma(\alpha)x^{\alpha/2}} \int_0^\infty e^{-\frac{w^2}{2}-wB\sqrt{x}} w^{\alpha-1} dw \\ &= \lambda \frac{e^{-\theta_+ \theta_- x/2}}{x^{\alpha/2}} H_{-\alpha} \left(\frac{\theta_+ + \theta_-}{2} \sqrt{x} \right) \end{aligned} \quad (5.28)$$

where H_a , $a < 0$, is the Hermite function, given by

$$H_a(z) = \frac{1}{\Gamma(-a)} \int_0^\infty e^{-x^2/2-xz} x^{-a-1} dx, \quad z > 0. \quad (5.29)$$

The corresponding Lévy density coincides with that determined in Madan and Yor (2008), Proposition 2.

Example 11. *Subordinated representation of generalized- z processes.* Let $\mu \in \text{GZD}_G(\mathbb{R})$ with Lévy measure of the form (4.20), i.e. $c(\pm 1) = c_\pm > 0$, $\sigma(\pm) = \sigma > 0$ and $\lambda(du) = \lambda(\delta_1(du) + \delta_{-1}(du))$, $\lambda > 0$. We can rewrite again $k(x) = e^{Ax/\sigma-B|x|/\sigma}/(1-e^{-|x|/\sigma})$, $A = (c_- - c_+)/2$, $B = (c_- + c_+)/2$ so we are under the assumption of Theorem 6. Based on (4.22) and (5.25), integrating term by term the series results in

$$\begin{aligned} k_\rho(x) &= \lambda e^{\frac{A^2x}{2\sigma^2}} \sum_{k=0}^\infty \exp\left(-x \frac{(k+B)^2}{2\sigma^2}\right) = \lambda \sum_{k=0}^\infty \exp\left(-x \frac{k^2 + 2Bk + B^2 - A^2}{2\sigma^2}\right) \\ &= \lambda \sum_{k=0}^\infty \exp\left(-x \frac{(k+c_+)(k+c_-)}{2\sigma^2}\right). \end{aligned} \quad (5.30)$$

The subordinator is then of the form of an infinite GGC convolution of gamma processes $G^k = (G_t^k)_{t \geq 0}$, $k \geq 0$, $\mathcal{L}(G_1^k) \sim \Gamma\left(\lambda, \frac{(k+c_+)(k+c_-)}{2\sigma^2}\right)$ whose limit in law is not otherwise known.

and then observing that the canonical function of Y writes as $k(x) = \lambda e^{Ax-B|x|}$, implying $\tau_+^{-A} = \tau_-^A = \lambda \delta_B$. The Dirac integration to which (5.25) reduces, produces

$$k_\rho(x) = \lambda e^{(A^2-B^2)x/2} = \lambda e^{-\theta_+ \theta_- x/2}. \quad (5.27)$$

Therefore, the bilateral gamma process can be represented in law as $Y_t = W_{T_t} + (\theta_- - \theta_+)t/2$ where $\mathcal{L}(T_1) \sim \Gamma(\lambda, \theta_+ \theta_-/2)$. This has been known and used at least since Madan et al. (1998).

Example 10. *Subordinated representation of CTS_α processes.* Madan and Yor (2008) through a series of propositions covering a substantial part of the article, explain how to represent a $\text{CTS}_\alpha(\mathbb{R})$ process as a subordinated Brownian motion with drift. Based on the discussion so far, the main Proposition 2 therein trivialize. First of all notice that the assumptions in Theorem 6 are met also in the case of $\text{CTS}_\alpha(\mathbb{R})$ processes with symmetric spherical part. Namely, assuming $\lambda_+ = \lambda_-$ and letting again $A = (\theta_- - \theta_+)/2$, $B = (\theta_- + \theta_-)/2$, leads to $k(x) = x^{-\alpha} e^{Ax-B|x|}$, so that using the Thorin measure found in Subsection 4.3 we deduce the required density relationship $\tau_+^{-A}(y) = \tau_-^A(y) = \lambda(y-B)^{\alpha-1}(\Gamma(\alpha))^{-1} \mathbf{1}_{\{y \geq B\}}$. It then follows from Corollary 6, and with the substitution $w = (s-B)\sqrt{x}$, that

$$\begin{aligned} k_\rho(x) &= \lambda \frac{e^{xA^2/2}}{\Gamma(\alpha)} \int_B^\infty e^{-\frac{s^2x}{2}} (s-B)^{\alpha-1} ds = \lambda \frac{e^{(A^2-B^2)x/2}}{\Gamma(\alpha)x^{\alpha/2}} \int_0^\infty e^{-\frac{w^2}{2}-wB\sqrt{x}} w^{\alpha-1} dw \\ &= \lambda \frac{e^{-\theta_+ \theta_- x/2}}{x^{\alpha/2}} H_{-\alpha} \left(\frac{\theta_+ + \theta_-}{2} \sqrt{x} \right) \end{aligned} \quad (5.28)$$

where H_a , $a < 0$, is the Hermite function, given by

$$H_a(z) = \frac{1}{\Gamma(-a)} \int_0^\infty e^{-x^2/2-xz} x^{-a-1} dx, \quad z > 0. \quad (5.29)$$

The corresponding Lévy density coincides with that determined in Madan and Yor (2008), Proposition 2.

Example 11. *Subordinated representation of generalized- z processes.* Let $\mu \in \text{GZD}_G(\mathbb{R})$ with Lévy measure of the form (4.20), i.e. $c(\pm 1) = c_\pm > 0$, $\sigma(\pm) = \sigma > 0$ and $\lambda(du) = \lambda(\delta_1(du) + \delta_{-1}(du))$, $\lambda > 0$. We can rewrite again $k(x) = e^{Ax/\sigma-B|x|/\sigma}/(1-e^{-|x|/\sigma})$, $A = (c_- - c_+)/2$, $B = (c_- + c_+)/2$ so we are under the assumption of Theorem 6. Based on (4.22) and (5.25), integrating term by term the series results in

$$\begin{aligned} k_\rho(x) &= \lambda e^{\frac{A^2x}{2\sigma^2}} \sum_{k=0}^\infty \exp\left(-x \frac{(k+B)^2}{2\sigma^2}\right) = \lambda \sum_{k=0}^\infty \exp\left(-x \frac{k^2 + 2Bk + B^2 - A^2}{2\sigma^2}\right) \\ &= \lambda \sum_{k=0}^\infty \exp\left(-x \frac{(k+c_+)(k+c_-)}{2\sigma^2}\right). \end{aligned} \quad (5.30)$$

The subordinator is then of the form of an infinite GGC convolution of gamma processes $G^k = (G_t^k)_{t \geq 0}$, $k \geq 0$, $\mathcal{L}(G_1^k) \sim \Gamma\left(\lambda, \frac{(k+c_+)(k+c_-)}{2\sigma^2}\right)$ whose limit in law is not otherwise known.

and then observing that the canonical function of Y writes as $k(x) = \lambda e^{Ax-B|x|}$, implying $\tau_+^{-A} = \tau_-^A = \lambda \delta_B$. The Dirac integration to which (5.25) reduces, produces

$$k_\rho(x) = \lambda e^{(A^2-B^2)x/2} = \lambda e^{-\theta_+ \theta_- x/2}. \quad (5.27)$$

Therefore, the bilateral gamma process can be represented in law as $Y_t = W_{T_t} + (\theta_- - \theta_+)t/2$ where $\mathcal{L}(T_1) \sim \Gamma(\lambda, \theta_+ \theta_-/2)$. This has been known and used at least since Madan et al. (1998).

Example 10. Subordinated representation of CTS_α processes. Madan and Yor (2008) through a series of propositions covering a substantial part of the article, explain how to represent a $\text{CTS}_\alpha(\mathbb{R})$ process as a subordinated Brownian motion with drift. Based on the discussion so far, the main Proposition 2 therein trivialize. First of all notice that the assumptions in Theorem 6 are met also in the case of $\text{CTS}_\alpha(\mathbb{R})$ processes with symmetric spherical part. Namely, assuming $\lambda_+ = \lambda_-$ and letting again $A = (\theta_- - \theta_+)/2$, $B = (\theta_- + \theta_+)/2$, leads to $k(x) = x^{-\alpha} e^{Ax-B|x|}$, so that using the Thorin measure found in Subsection 4.3 we deduce the required density relationship $\tau_+^{-A}(y) = \tau_-^A(y) = \lambda(y-B)^{\alpha-1}(\Gamma(\alpha))^{-1} \mathbf{1}_{\{y \geq B\}}$. It then follows from Corollary 6, and with the substitution $w = (s-B)\sqrt{x}$, that

$$\begin{aligned} k_\rho(x) &= \lambda \frac{e^{xA^2/2}}{\Gamma(\alpha)} \int_B^\infty e^{-\frac{s^2x}{2}} (s-B)^{\alpha-1} ds = \lambda \frac{e^{(A^2-B^2)x/2}}{\Gamma(\alpha)x^{\alpha/2}} \int_0^\infty e^{-\frac{w^2}{2}-wB\sqrt{x}} w^{\alpha-1} dw \\ &= \lambda \frac{e^{-\theta_+ \theta_- x/2}}{x^{\alpha/2}} H_{-\alpha} \left(\frac{\theta_+ + \theta_-}{2} \sqrt{x} \right) \end{aligned} \quad (5.28)$$

where H_a , $a < 0$, is the Hermite function, given by

$$H_a(z) = \frac{1}{\Gamma(-a)} \int_0^\infty e^{-x^2/2-xz} x^{-a-1} dx, \quad z > 0. \quad (5.29)$$

The corresponding Lévy density coincides with that determined in Madan and Yor (2008),

Proposition 2.

Example 11. Subordinated representation of generalized- z processes. Let $\mu \in \text{GZD}_G(\mathbb{R})$ with Lévy measure of the form (4.20), i.e. $c(\pm 1) = c_\pm > 0$, $\sigma(\pm) = \sigma > 0$ and $\lambda(du) = \lambda(\delta_1(du) + \delta_{-1}(du))$, $\lambda > 0$. We can rewrite again $k(x) = e^{Ax/\sigma-B|x|/\sigma}/(1-e^{-|x|/\sigma})$, $A = (c_- - c_+)/2$, $B = (c_- + c_+)/2$ so we are under the assumption of Theorem 6. Based on (4.22) and (5.25), integrating term by term the series results in

$$\begin{aligned} k_\rho(x) &= \lambda e^{\frac{A^2x}{2\sigma^2}} \sum_{k=0}^\infty \exp\left(-x \frac{(k+B)^2}{2\sigma^2}\right) = \lambda \sum_{k=0}^\infty \exp\left(-x \frac{k^2 + 2Bk + B^2 - A^2}{2\sigma^2}\right) \\ &= \lambda \sum_{k=0}^\infty \exp\left(-x \frac{(k+c_+)(k+c_-)}{2\sigma^2}\right). \end{aligned} \quad (5.30)$$

The subordinator is then of the form of an infinite GGC convolution of gamma processes $G^k = (G_t^k)_{t \geq 0}$, $k \geq 0$, $\mathcal{L}(G_1^k) \sim \Gamma\left(\lambda, \frac{(k+c_+)(k+c_-)}{2\sigma^2}\right)$ whose limit in law is not otherwise known.