

Objectives

- Objectives for today:
- Introducing specific vocabulary.
 - Quick revision of quadratic function.
 - Factorising Quadratics.
 - Proving Vieta's formulas.
 - Carrying out gained knowledge by working out some word problems.

Quick Revision

Forms of Quadratic Function

- $f(x) = ax^2 + bx + c$ is called the **standard form**.
- $f(x) = a(x - x_1)(x - x_2)$ is called the **factored form**, where x_1 and x_2 are the roots of the quadratic function.
- $f(x) = a(x - h)^2 + k$ is called the **vertex form**.

Delta Δ

Δ determines tells us how many solutions quadratic equation have:

number of solutions =

2

when $\Delta > 0$

1

when $\Delta = 0$

0

when $\Delta < 0$

The Quadratic Formula

$$x = \frac{-b \pm \sqrt{\Delta}}{2a}$$

Graph of Quadratic Function




Figure: Graph of $f(x) = ax^2|_{\{0,1,0.3,1.0,3.0\}}$

Factorising a Quadratic

Factorising a quadratic means putting it into two brackets, and is useful if you're trying to draw a graph of a quadratic solve a quadratic equation. It's pretty easy if $a = 1$ (in $ax^2 + bx + c$ form), but can be a real pain otherwise.

In order to factorise a quadratic you should follow steps outlined below:

- ➊ Rearrange the equation into the standard $ax^2 + bx + c$ form.
- ➋ Write down two brackets: $(x \quad)(x \quad)$
- ➌ Find two numbers that multiply to give 'c' and add or subtract to give 'b' (ignoring signs).
- ➍ Put the numbers in brackets and choose their signs.

Myth of Delta Δ

It's commonly believed that in order to work out roots of a quadratic function you must count Δ and use other previously established formulas. However this is untrue since factorising in many cases is as good or even better than simply counting Δ .

Example of Factorisation

Solve $x^2 + 4x - 21 = 0$ by factorising.

$$x^2 + 4x - 21 = (x \quad)(x \quad)$$

- 1 and 21 multiply to give 21 - and add or subtract to give 22 and 20.
3 and 7 multiply to give 21 - and add or subtract to give 10 and 4.

$$x^2 + 4x + 21 = (x + 7)(x - 3)$$

And solving the equation:

$$(x + 7)(x - 3) = 0$$

we get

$$x = -7, \quad x = 3$$

Factorising- Tasks

1. Factorise $x^2 - x - 12$.
2. Solve $x^2 - 8 = 2x$ by factorising.

Proof of Vieta's Formulas

Let's prove that:

$$x_1 + x_2 = \frac{-b}{a}$$

When Δ is positive we have two roots:

$$x_1 = \frac{-b - \sqrt{\Delta}}{2a}, \quad x_2 = \frac{-b + \sqrt{\Delta}}{2a}$$

Substituting for x_1 and x_2 respectively, we receive:

$$\begin{aligned} x_1 + x_2 &= \frac{-b - \sqrt{\Delta}}{2a} + \frac{-b + \sqrt{\Delta}}{2a} = \\ &= \frac{(-b - \sqrt{\Delta}) + (-b + \sqrt{\Delta})}{2a} = \frac{-2b}{2a} = \frac{-b}{a} \end{aligned}$$

The same we could do with another pattern, which state that $x_1x_2 = \frac{c}{a}$, but proving this is going to be your task in next section.

Vieta's Formulas- Task

1. Prove that

$$x_1x_2 = \frac{c}{a}$$

Glossary

| verb | noun | meaning |
|------------|----------------|----------------|
| add | addition | + |
| subtract | subtraction | − |
| multiply | multiplication | · |
| divide | division | ÷ |
| solve | solution | getting answer |
| substitute | substitution | $t = x^2$ |

[Table](#): Word Formation

Some Necessary and Useful Vocabulary

- (n.) sign $\rightarrow +$ or $-$
- (n.) equation $\rightarrow something = 0$
- (n.) factor \rightarrow two multiplied factors give result
- (v.) factorise \rightarrow putting into brackets
- (n.) coefficient \rightarrow a constant number i.e. a, b, c in a pattern $ax^2 + bx + c$
- (n.) quadratic function $\rightarrow f(x) = ax^2 + bx + c$
- (n.) root $\rightarrow \sqrt{sth}$ or solution of quadratic equation
- (n.) formula = pattern