

# Large scale Finite Element solvers for the large eddy simulation of incompressible turbulent flows

*author:*

ORIOL COLOMÉS GENÉ

*supervisor:*

SANTIAGO BADIA

Departament d'Enginyeria Civil i Ambiental

February 24, 2016



**Escola de Camins**

Escola Tècnica Superior d'Enginyeria de Camins, Canals i Ports  
UPC BARCELONATECH

1. Motivation
2. Residual-based VMS
3. Mixed FE VMS
4. Segregated Runge-Kutta
5. Segregated VMS
6. Conclusions

# 1. Motivation

## 2. Residual-based VMS

## 3. Mixed FE VMS

## 4. Segregated Runge-Kutta

## 5. Segregated VMS

## 6. Conclusions

# Motivation

## Thesis motivation

Highly scalable Finite Element (FE) framework for Large Eddy Simulations (LES) of incompressible turbulent flows

# Motivation

## Thesis motivation

Highly scalable Finite Element (FE) framework for Large Eddy Simulations (LES) of incompressible turbulent flows

## How to get there?

# Motivation

## Thesis motivation

Highly scalable Finite Element (FE) framework for Large Eddy Simulations (LES) of incompressible turbulent flows

## How to get there?

1. **Variational MultiScale** (VMS) methods as LES models.

# Motivation

## Thesis motivation

Highly scalable Finite Element (FE) framework for Large Eddy Simulations (LES) of incompressible turbulent flows

## How to get there?

1. Variational MultiScale (VMS) methods as LES models.
2. Time integration schemes with **velocity-pressure segregation**.

# Motivation

## Thesis motivation

Highly scalable Finite Element (FE) framework for Large Eddy Simulations (LES) of incompressible turbulent flows

## How to get there?

1. Variational MultiScale (VMS) methods as LES models.
2. Time integration schemes with velocity-pressure segregation.
3. Highly scalable algorithms based on **Domain Decomposition (DD)** and **block preconditioners**.



# Motivation

## Step by step...

- Residual-based VMS as LES models.

# Motivation

## Step by step...

- Residual-based VMS as LES models.
- Mixed FE formulations LES.

# Motivation

## Step by step...

- Residual-based VMS as LES models.
- Mixed FE formulations LES.
- High-order FE methods.

# Motivation

## Step by step...

- Residual-based VMS as LES models.
- Mixed FE formulations LES.
- High-order FE methods.
- High-order time integration schemes.

# Motivation

## Step by step...

- Residual-based VMS as LES models.
- Mixed FE formulations LES.
- High-order FE methods.
- High-order time integration schemes.
- Adaptive time stepping techniques.

# Motivation

## Step by step...

- Residual-based VMS as LES models.
- Mixed FE formulations LES.
- High-order FE methods.
- High-order time integration schemes.
- Adaptive time stepping techniques.
- Velocity-pressure segregation.

# Motivation

## Step by step...

- Residual-based VMS as LES models.
- Mixed FE formulations LES.
- High-order FE methods.
- High-order time integration schemes.
- Adaptive time stepping techniques.
- Velocity-pressure segregation.
- Scalable solvers.

# Motivation

## Step by step...

- Residual-based VMS as LES models.
- Mixed FE formulations LES.
- High-order FE methods.
- High-order time integration schemes.
- Adaptive time stepping techniques.
- Velocity-pressure segregation.
- Scalable solvers.
- Application.



# Motivation

## Step by step...

- Residual-based VMS as LES models.
- Mixed FE formulations LES.
- High-order FE methods.
- High-order time integration schemes.
- Adaptive time stepping techniques.
- Velocity-pressure segregation.
- Scalable solvers.
- Application.

## 1. Motivation


## 2. Residual-based VMS

Formulation

Energy statements

Numerical experiments

Conclusions



Rb-VMS

## 3. Mixed FE VMS

## 4. Segregated Runge-Kutta

## 5. Segregated VMS

## 6. Conclusions

# Implicit LES

**ILES:** only numerical dissipation (for stabilization) acts as turbulent model

- Not based on filtering of the Navier-Stokes equations
- Rely on numerical artifacts, no modification at the continuous level



# Incomp. Navier Stokes equations

Find  $\mathbf{u}$  and  $p$  defined in  $\Omega$

$$\begin{aligned}\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p - \nu \nabla^2 \mathbf{u} &= \mathbf{f} \\ \nabla \cdot \mathbf{u} &= 0\end{aligned}$$

with appropriate boundary conditions on  $\Gamma$ .

# Incomp. Navier Stokes equations

Find  $\mathbf{u}$  and  $p$  defined in  $\Omega$

$$\begin{aligned}\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p - \nu \nabla^2 \mathbf{u} &= \mathbf{f} \\ \nabla \cdot \mathbf{u} &= 0\end{aligned}$$

with appropriate boundary conditions on  $\Gamma$ .

The weak problem is:  $\forall \mathbf{v} \in \mathcal{V}_0$  and  $\forall q \in \mathcal{Q}_0$ , find  $\mathbf{u} \in \mathcal{V}$  and  $p \in \mathcal{Q}$  such that

$$\begin{aligned}(\mathbf{v}, \partial_t \mathbf{u})_{\Omega} + (\nabla \mathbf{v}, \nu \nabla \mathbf{u})_{\Omega} + b(\mathbf{u}, \mathbf{u}, \mathbf{v}) - (\nabla \cdot \mathbf{v}, p)_{\Omega} &= \langle \mathbf{v}, \mathbf{f} \rangle_{\Omega} \\ (q, \nabla \cdot \mathbf{u})_{\Omega} &= 0\end{aligned}$$

# Incomp. Navier Stokes equations

Find  $\mathbf{u}$  and  $p$  defined in  $\Omega$

$$\begin{aligned}\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p - \nu \nabla^2 \mathbf{u} &= \mathbf{f} \\ \nabla \cdot \mathbf{u} &= 0\end{aligned}$$

with appropriate boundary conditions on  $\Gamma$ .

The weak problem is:  $\forall \mathbf{v} \in \mathcal{V}_0$  and  $\forall q \in \mathcal{Q}_0$ , find  $\mathbf{u} \in \mathcal{V}$  and  $p \in \mathcal{Q}$  such that

$$\begin{aligned}(\mathbf{v}, \partial_t \mathbf{u})_\Omega + (\nabla \mathbf{v}, \nu \nabla \mathbf{u})_\Omega + b(\mathbf{u}, \mathbf{u}, \mathbf{v}) - (\nabla \cdot \mathbf{v}, p)_\Omega &= \langle \mathbf{v}, \mathbf{f} \rangle_\Omega \\ (q, \nabla \cdot \mathbf{u})_\Omega &= 0\end{aligned}$$

where

$$b(\mathbf{a}, \mathbf{u}, \mathbf{v}) = \langle \mathbf{v}, \mathbf{a} \cdot \nabla \mathbf{u} \rangle_\Omega$$

# Incomp. Navier Stokes equations

Find  $\mathbf{u}$  and  $p$  defined in  $\Omega$

$$\begin{aligned}\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p - \nu \nabla^2 \mathbf{u} &= \mathbf{f} \\ \nabla \cdot \mathbf{u} &= 0\end{aligned}$$

with appropriate boundary conditions on  $\Gamma$ .

The weak problem is:  $\forall \mathbf{v} \in \mathcal{V}_0$  and  $\forall q \in \mathcal{Q}_0$ , find  $\mathbf{u} \in \mathcal{V}$  and  $p \in \mathcal{Q}$  such that

$$\begin{aligned}(\mathbf{v}, \partial_t \mathbf{u})_{\Omega} + (\nabla \mathbf{v}, \nu \nabla \mathbf{u})_{\Omega} + b(\mathbf{u}, \mathbf{u}, \mathbf{v}) - (\nabla \cdot \mathbf{v}, p)_{\Omega} &= \langle \mathbf{v}, \mathbf{f} \rangle_{\Omega} \\ (q, \nabla \cdot \mathbf{u})_{\Omega} &= 0\end{aligned}$$

where

$$b(\mathbf{a}, \mathbf{u}, \mathbf{v}) = \frac{1}{2} \langle \mathbf{v}, \mathbf{a} \cdot \nabla \mathbf{u} \rangle_{\Omega} - \frac{1}{2} \langle \mathbf{a} \cdot \nabla \mathbf{v}, \mathbf{u} \rangle_{\Omega} + \frac{1}{2} \langle \mathbf{v}, \mathbf{n} \cdot \mathbf{a} \mathbf{u} \rangle_{\Gamma}$$

# VMS decomposition (Hughes 1995)

A decomposition of spaces  $\mathcal{V}$  and  $\mathcal{Q}$  given by

$$\mathcal{V} = \mathcal{V}_h \oplus \tilde{\mathcal{V}}, \quad \mathcal{Q} = \mathcal{Q}_h \oplus \tilde{\mathcal{Q}}$$



# VMS decomposition (Hughes 1995)

A decomposition of spaces  $\mathcal{V}$  and  $\mathcal{Q}$  given by

$$\mathcal{V} = \mathcal{V}_h \oplus \tilde{\mathcal{V}}, \quad \mathcal{Q} = \mathcal{Q}_h \oplus \tilde{\mathcal{Q}}$$

is applied to the function and test spaces

$$\mathbf{u} = \mathbf{u}_h + \tilde{\mathbf{u}}, \quad p = p_h + \tilde{p}$$

$$\mathbf{v} = \mathbf{v}_h + \tilde{\mathbf{v}}, \quad q = q_h + \tilde{q}$$

# VMS decomposition (Hughes 1995)

A decomposition of spaces  $\mathcal{V}$  and  $\mathcal{Q}$  given by

$$\mathcal{V} = \mathcal{V}_h \oplus \tilde{\mathcal{V}}, \quad \mathcal{Q} = \mathcal{Q}_h \oplus \tilde{\mathcal{Q}}$$

is applied to the function and test spaces

$$\mathbf{u} = \mathbf{u}_h + \tilde{\mathbf{u}}, \quad p = p_h + \tilde{p}$$

$$\mathbf{v} = \mathbf{v}_h + \tilde{\mathbf{v}}, \quad q = q_h + \tilde{q}$$

We keep all the (eight) contributions from the splitting of the convective term

$$\mathbf{u} \cdot \nabla \mathbf{u} = \mathbf{u}_h \cdot \nabla \mathbf{u}_h + \tilde{\mathbf{u}} \cdot \nabla \mathbf{u}_h + \mathbf{u}_h \cdot \nabla \tilde{\mathbf{u}} + \tilde{\mathbf{u}} \cdot \nabla \tilde{\mathbf{u}}$$

# VMS decomposition (Hughes 1995)

A decomposition of spaces  $\mathcal{V}$  and  $\mathcal{Q}$  given by

$$\mathcal{V} = \mathcal{V}_h \oplus \tilde{\mathcal{V}}, \quad \mathcal{Q} = \mathcal{Q}_h \oplus \tilde{\mathcal{Q}}$$

is applied to the function and test spaces

$$\mathbf{u} = \mathbf{u}_h + \tilde{\mathbf{u}}, \quad p = p_h + \tilde{p}$$

$$\mathbf{v} = \mathbf{v}_h + \tilde{\mathbf{v}}, \quad q = q_h + \tilde{q}$$

We keep all the (eight) contributions from the splitting of the convective term

$$\mathbf{u} \cdot \nabla \mathbf{u} = \mathbf{u}_h \cdot \nabla \mathbf{u}_h + \tilde{\mathbf{u}} \cdot \nabla \mathbf{u}_h + \mathbf{u}_h \cdot \nabla \tilde{\mathbf{u}} + \tilde{\mathbf{u}} \cdot \nabla \tilde{\mathbf{u}}$$

and all the (four) contributions from the temporal term

$$\partial_t \mathbf{u} = \partial_t \mathbf{u}_h + \partial_t \tilde{\mathbf{u}}$$

# Semidiscrete problem

## FEM equations

$$B((\mathbf{u}_h, p_h); (\tilde{\mathbf{u}}, \tilde{p}); (\mathbf{v}_h, q_h)) = L(\mathbf{v}_h, q_h)$$

## SGS equations

$$B((\tilde{\mathbf{u}}, \tilde{p}); (\mathbf{u}_h, p_h); (\tilde{\mathbf{v}}, \tilde{q})) = L(\tilde{\mathbf{v}}, \tilde{q})$$

# Semidiscrete problem

## FEM equations

$$\begin{aligned}
 &(\mathbf{v}_h, \partial_t \mathbf{u}_h)_\Omega + b(\mathbf{a}, \mathbf{u}_h, \mathbf{v}_h) + (\nabla \mathbf{v}_h, \nu \nabla \mathbf{u}_h)_\Omega - (\nabla \cdot \mathbf{v}_h, p_h)_\Omega \\
 &+ (\mathbf{v}_h, \partial_t \tilde{\mathbf{u}})_\Omega + (\mathcal{L}^* \mathbf{v}_h, \tilde{\mathbf{u}})_{\Omega^h} - (\nabla \cdot \mathbf{v}_h, \tilde{p})_{\Omega^h} = \langle \mathbf{v}_h, \mathbf{f} \rangle_\Omega \\
 &(q_h, \nabla \cdot \mathbf{u}_h)_\Omega - (\nabla q_h, \tilde{\mathbf{u}})_{\Omega^h} = 0
 \end{aligned}$$

## SGS equations

$$\partial_t \tilde{\mathbf{u}} + \tau_m^{-1} \tilde{\mathbf{u}} = \mathcal{P} \mathbf{R}_m$$

$$\tau_c^{-1} \tilde{p} = \mathcal{P} R_c$$

$$\tau_m = \left( \frac{c_1 \nu}{h^2} + \frac{c_2 |\mathbf{a}|}{h} \right)^{-1}, \quad \tau_c = \frac{h^2}{c_1 \tau_m}$$

$$\mathbf{a} = \mathbf{u}_h + \tilde{\mathbf{u}}$$

# Semidiscrete problem

## FEM equations

$$\begin{aligned}
 &(\mathbf{v}_h, \partial_t \mathbf{u}_h)_\Omega + b(\mathbf{a}, \mathbf{u}_h, \mathbf{v}_h) + (\nabla \mathbf{v}_h, \nu \nabla \mathbf{u}_h)_\Omega - (\nabla \cdot \mathbf{v}_h, p_h)_\Omega \\
 &+ (\mathbf{v}_h, \partial_t \tilde{\mathbf{u}})_\Omega + (\mathcal{L}^* \mathbf{v}_h, \tilde{\mathbf{u}})_{\Omega^h} - (\nabla \cdot \mathbf{v}_h, \tilde{p})_{\Omega^h} = \langle \mathbf{v}_h, \mathbf{f} \rangle_\Omega \\
 &(q_h, \nabla \cdot \mathbf{u}_h)_\Omega - (\nabla q_h, \tilde{\mathbf{u}})_{\Omega^h} = 0
 \end{aligned}$$

## SGS equations

$$\partial_t \tilde{\mathbf{u}} + \tau_m^{-1} \tilde{\mathbf{u}} = \mathcal{P} \mathbf{R}_m$$

$$\tau_c^{-1} \tilde{p} = \mathcal{P} R_c$$

$$\mathbf{R}_m := \mathbf{f} - \partial_t \mathbf{u}_h - \mathcal{L} \mathbf{u}_h - \nabla p_h, \quad R_c := -\nabla \cdot \mathbf{u}_h$$

# Semidiscrete problem

## FEM equations

$$\begin{aligned}
 &(\mathbf{v}_h, \partial_t \mathbf{u}_h)_\Omega + b(\mathbf{a}, \mathbf{u}_h, \mathbf{v}_h) + (\nabla \mathbf{v}_h, \nu \nabla \mathbf{u}_h)_\Omega - (\nabla \cdot \mathbf{v}_h, p_h)_\Omega \\
 &+ (\mathbf{v}_h, \partial_t \tilde{\mathbf{u}})_\Omega + (\mathcal{L}^* \mathbf{v}_h, \tilde{\mathbf{u}})_{\Omega^h} - (\nabla \cdot \mathbf{v}_h, \tilde{p})_{\Omega^h} = \langle \mathbf{v}_h, \mathbf{f} \rangle_\Omega \\
 &(q_h, \nabla \cdot \mathbf{u}_h)_\Omega - (\nabla q_h, \tilde{\mathbf{u}})_{\Omega^h} = 0
 \end{aligned}$$

## SGS equations

$$\partial_t \tilde{\mathbf{u}} + \tau_m^{-1} \tilde{\mathbf{u}} = \mathcal{P} \mathbf{R}_m$$

$$\tau_c^{-1} \tilde{p} = \mathcal{P} R_c$$

$$\mathcal{P} = I \quad (\text{ASGS}), \quad \mathcal{P} = P_h^\perp = I - P_h \quad (\text{OSS})$$

# Semidiscrete problem

## FEM equations

$$\begin{aligned}
 &(\mathbf{v}_h, \partial_t \mathbf{u}_h)_\Omega + b(\mathbf{a}, \mathbf{u}_h, \mathbf{v}_h) + (\nabla \mathbf{v}_h, \nu \nabla \mathbf{u}_h)_\Omega - (\nabla \cdot \mathbf{v}_h, p_h)_\Omega \\
 &+ (\mathbf{v}_h, \partial_t \tilde{\mathbf{u}})_\Omega + (\mathcal{L}^* \mathbf{v}_h, \tilde{\mathbf{u}})_{\Omega^h} - (\nabla \cdot \mathbf{v}_h, \tilde{p})_{\Omega^h} = \langle \mathbf{v}_h, \mathbf{f} \rangle_\Omega \\
 &(q_h, \nabla \cdot \mathbf{u}_h)_\Omega - (\nabla q_h, \tilde{\mathbf{u}})_{\Omega^h} = 0
 \end{aligned}$$

## SGS equations

$$\partial_t \tilde{\mathbf{u}} + \tau_m^{-1} \tilde{\mathbf{u}} = \mathcal{P} \mathbf{R}_m$$

$$\tau_c^{-1} \tilde{p} = \mathcal{P} R_c$$

$$\mathcal{P} = I \quad (\text{ASGS}), \quad \mathcal{P} = P_h^\perp = I - P_h \quad (\text{OSS})$$

$$\mathbf{a} = \mathbf{u}_h + \tilde{\mathbf{u}}$$



# Summary

	Sgs space	Sgs dynamics	Advection
1	ASGS	Static	Linear
2	ASGS	Dynamic	Linear
3	ASGS	Dynamic	Nonlinear
4	OSS	Static	Linear
5	OSS	Dynamic	Linear
6	OSS	Dynamic	Nonlinear

- 1 It is the most standard method (SUPG for linear elements) up to the choice of the stabilization parameters. Unknown stability properties.
- 4 Strictly pointwise positive for linear elements (no backscatter).
- 5 Convergent to weak solutions of NS equations (Badia & Gutierrez 2012).

# Energy statements

**FE counterpart:**

$$B((\mathbf{u}_h, p_h); (\tilde{\mathbf{u}}, \tilde{p}); (\mathbf{u}_h, p_h)) = L(\mathbf{u}_h, p_h)$$

**SGS counterpart:**

$$B((\tilde{\mathbf{u}}, \tilde{p}); (\mathbf{u}_h, p_h); (\tilde{\mathbf{u}}, \tilde{p})) = L(\tilde{\mathbf{u}}, \tilde{p})$$

**TOTAL:**

$$\begin{aligned} & B((\mathbf{u}_h, p_h); (\tilde{\mathbf{u}}, \tilde{p}); (\mathbf{u}_h, p_h)) \\ & + B((\tilde{\mathbf{u}}, \tilde{p}); (\mathbf{u}_h, p_h); (\tilde{\mathbf{u}}, \tilde{p})) = L(\mathbf{u}_h, p_h) + L(\tilde{\mathbf{u}}, \tilde{p}) \end{aligned}$$

# Energy statements

## FE counterpart:

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

## SGS counterpart:

$$\begin{aligned} & \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

## TOTAL:

# Energy statements

**FE counterpart:**

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

**SGS counterpart:**

$$\begin{aligned} & \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

**TOTAL:**

$$\frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2$$

# Energy statements

## FE counterpart:

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

## SGS counterpart:

$$\begin{aligned} & \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

## TOTAL:

$$\frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \nu \|\nabla \mathbf{u}_h\|^2$$

# Energy statements

## FE counterpart:

$$\begin{aligned} \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

## SGS counterpart:

$$\begin{aligned} \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

## TOTAL:

$$\frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \nu \|\nabla \mathbf{u}_h\|^2$$

# Energy statements

## FE counterpart:

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

## SGS counterpart:

$$\begin{aligned} & \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

## TOTAL:

$$\frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2$$

# Energy statements

## FE counterpart:

$$\begin{aligned} \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

## SGS counterpart:

$$\begin{aligned} \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

## TOTAL:

$$\begin{aligned} \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) \\ - (\nabla \cdot \mathbf{u}_h, \tilde{p}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) \end{aligned}$$



# Energy statements

## FE counterpart:

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

## SGS counterpart:

$$\begin{aligned} & \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

## TOTAL:

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) \\ & - (\nabla \cdot \mathbf{u}_h, \tilde{p}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle + \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

# Energy statements

**FE counterpart:**

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

**SGS counterpart:**

$$\begin{aligned} & \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

**TOTAL: Static subscales**

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) \\ & - (\nabla \cdot \mathbf{u}_h, \tilde{p}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle + \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

# Energy statements

**FE counterpart:**

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

**SGS counterpart:**

$$\begin{aligned} & \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

**TOTAL: Dynamic subscales - ASGS**

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h + \tilde{\mathbf{u}}\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & - 2(\nu \Delta \mathbf{u}_h, \tilde{\mathbf{u}}) \\ & = \langle \mathbf{f}, \mathbf{u}_h \rangle + \langle \mathbf{f}, \tilde{\mathbf{u}} \rangle \end{aligned}$$

# Energy statements

**FE counterpart:**

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + b(\mathbf{a}, \mathbf{u}_h, \mathbf{u}_h) \\ & + (\partial_t \tilde{\mathbf{u}}, \mathbf{u}_h) + (\mathcal{L}_a^*(\mathbf{u}_h, p_h), \tilde{\mathbf{u}}) - (\nabla \cdot \mathbf{u}_h, \tilde{p}) = \langle \mathbf{f}, \mathbf{u}_h \rangle \end{aligned}$$

**SGS counterpart:**

$$\begin{aligned} & \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & + (\mathcal{P}(\partial_t \mathbf{u}_h), \tilde{\mathbf{u}}) + (\mathcal{P}(\mathcal{L}_a^*(\mathbf{u}_h, p_h)), \tilde{\mathbf{u}}) - (\mathcal{P}(\nabla \cdot \mathbf{u}_h), \tilde{p}) = \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

**TOTAL: Dynamic subscales - OSS**

$$\begin{aligned} & \frac{1}{2} d_t \|\mathbf{u}_h\|^2 + \frac{1}{2} d_t \|\tilde{\mathbf{u}}\|^2 + \nu \|\nabla \mathbf{u}_h\|^2 + \tau_m^{-1} \|\tilde{\mathbf{u}}\|^2 + \tau_c^{-1} \|\tilde{p}\|^2 \\ & - 2(\nu \Delta \mathbf{u}_h, \tilde{\mathbf{u}}) \\ & = \langle \mathbf{f}, \mathbf{u}_h \rangle + \langle \mathcal{P}(\mathbf{f}), \tilde{\mathbf{u}} \rangle \end{aligned}$$

## 1. Motivation

## 2. Residual-based VMS


## 3. Mixed FE VMS

Formulation

Block-preconditioning

Numerical experiments

Conclusions



## 4. Segregated Runge-Kutta

## 5. Segregated VMS

## 6. Conclusions

## 1. Motivation


## 2. Residual-based VMS

## 3. Mixed FE VMS

4. Segregated Runge-Kutta  
Formulation  
Numerical experiments  
Conclusions

## 5. Segregated VMS

## 6. Conclusions



## 1. Motivation

## 2. Residual-based VMS

## 3. Mixed FE VMS

## 4. Segregated Runge-Kutta

## 5. Segregated VMS


Formulation

Block-preconditioning

Numerical experiments

Conclusions

## 6. Conclusions



1. Motivation
2. Residual-based VMS
3. Mixed FE VMS
4. Segregated Runge-Kutta
5. Segregated VMS
6. Conclusions



# Outline

- Line 1.

# Outline

- Line 1.
- Line 2.

Less formal

# Outline

- Line 1.
- Line 2.  
Less formal
- Line 3.  
Less formal, different color.

# Blocks

## Standard Block

This is a standard block.

## Example Block

This is an example block.

## Alert Block

This is an alert block.

# Questions?