

Time-series Analysis Notes

Nong Minh Hieu¹

¹ School of Physical and Mathematical Sciences, Nanyang Technological University (NTU - Singapore)

Contents

1	Introduction	2
1.1	Time-series definitions	2
1.2	Trend model	2
1.3	Trend + Seasonality model	3
1.4	Modelling seasonality using dummy variables	4
1.5	Autocorrelation & autocorrelation function (ACF)	5
2	Stationary time-series	6
2.1	Strict vs. Weak stationarity	6
A	List of Definitions	7
B	Important Theorems	7
C	Important Corollaries	7
D	Important Propositions	7
E	References	8

1 Introduction

1.1 Time-series definitions

Definition 1.1 (Time series).

A time-series is a sequence of observations of time-indexed random variables $(X_t)_{t \geq 0}$. Typically, A time-series consists of three components.

- Trend (T_t) .
- Cycle (C_t) .
- Seasonal (S_t) .

Finally, there is a random error (residual) term (R_t) . Note that for the sake of this course, we will hardly consider the cycle (C_t) term.

Definition 1.2 (Multiplicative & Additive models).

The components of a time-series can be combined either **additively** or **multiplicatively**:

- **Additive model** : $X_t = T_t + C_t + S_t + R_t$.
- **Multiplicative model** : $X_t = T_t \times C_t \times S_t \times R_t$.

1.2 Trend model

Definition 1.3 (Trend model).

A trend model is defined as:

$$X_t = T_t + R_t$$

Remark : Consider the linear trend:

$$T_t = \beta_0 + \beta_1 t, \quad \beta_1 \neq 0$$

We can then estimate the time-series $(X_t)_{t \geq 0}$ using the following methods:

- **Method 1** : Least square estimate the trend component T_t by minimizing the sum squared error

$$SSE = \sum_{t \geq 0} (X_t - T_t)^2$$

- **Method 2** : Eliminate the trend component via differencing operator.

Definition 1.4 (Differencing operator).

We define the first difference operator ∇ by:

$$\nabla X_t = X_t - X_{t-1} = (I - B)X_t$$

Where I is the identity operator and B is the backward shift operator:

$$BX_t = X_{t-1}, \quad B^i X_t = X_{t-i}$$

The i^{th} order differencing operator is:

$$\nabla^i X_t = \nabla(\nabla^{i-1} X_t)$$

Note that $\nabla^0 X_t = I$.

Proposition 1.1: Binomial formula for $\nabla^n X_t$

Given a time-series $(X_t)_{t \geq 0}$, we can write the differenced data to the n^{th} order as followed:

$$\nabla^n X_t = \sum_{k=0}^n \binom{n}{k} (-1)^k B^k X_t = \sum_{k=0}^n \binom{n}{k} (-1)^k X_{t-k}$$

As a result, we can write:

$$X_t = \nabla^n X_t - \sum_{k=1}^n \binom{n}{k} (-1)^k X_{t-k}$$

Proof (Proposition 1.1).

Using the Binomial theorem, we have:

$$\begin{aligned} \nabla^n X_t &= (I - B)^n X_t \\ &= \sum_{k=0}^n \binom{n}{k} (I)^{n-k} (-B)^k X_t \quad (\text{Binomial theorem}) \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k B^k X_t \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k X_{t-k} \\ &= X_t + \sum_{k=1}^n \binom{n}{k} (-1)^k X_{t-k} \\ \Rightarrow X_t &= \nabla^n X_t - \sum_{k=1}^n \binom{n}{k} (-1)^k X_{t-k} \end{aligned}$$

□.

1.3 Trend + Seasonality model

Definition 1.5 (Trend + Seasonality model).

The model that displays both trend and seasonality is defined as:

$$X_t = T_t + S_t + R_t$$

There are two types of seasonality:

- **Constant seasonality** : degree of seasonality does not depend on time.
- **Varying (increasing) seasonality** : degree of seasonality depends on time.

Remark : When the time-series displays increasing seasonality, it is a common practice to transform the data so that the series display constant seasonality.

Definition 1.6 (Box-Cox transformation). _____

For some $\lambda > 0$, the Box-Cox transformation is defined as:

$$z_t = \frac{x_t^\lambda - 1}{\lambda}$$

Proposition 1.2: Box-Cox as $\lambda \rightarrow 0$

As $\lambda \rightarrow 0$, $z_t \rightarrow \log(x_t)$.

Proof (Proposition 1.2). _____

We can write:

$$\begin{aligned} \lim_{\lambda \rightarrow 0} e^{z_t} &= \lim_{\lambda \rightarrow 0} \lim_{n \rightarrow \infty} \left(1 + \frac{z_t}{n} \right)^n \\ &= \lim_{\lambda \rightarrow 0} \left(1 + \lambda z_t \right)^{1/\lambda} \quad \left(\text{Letting } n = \frac{1}{\lambda} \right) \\ &= \lim_{\lambda \rightarrow 0} \left(1 + (x_t^\lambda - 1) \right)^{1/\lambda} = x_t \\ \Rightarrow \lim_{\lambda \rightarrow 0} z_t &= \log(x_t) \end{aligned}$$

□.

1.4 Modelling seasonality using dummy variables

Definition 1.7 (Dummy variables for seasonality). _____

The seasonal factor (supposing that there are L seasons) expressed using dummy variables can be written as:

$$S_t = \beta_1^{(s)} D_1(t) + \cdots + \beta_{L-1}^{(s)} D_{L-1}(t)$$

Where we have:

$$D_k(t) = \begin{cases} 1, & \text{if } t \text{ belongs to season } k \\ 0, & \text{otherwise} \end{cases}$$

We can write the time-series whose seasonality is modelled using dummy variables as:

$$X_t = T_t + \sum_{k=1}^{L-1} \beta_k^{(s)} D_k(t) + R_t$$

Remark : Take a time-series with linear trend and seasonality modelled using dummy variables as an example. Suppose that we have 4 seasons, the time-series will take the form:

$$X_t = \underbrace{\beta_0 + \beta_1 t}_{T_t} + \underbrace{\beta_1^{(s)} D_1(t) + \beta_2^{(s)} D_2(t) + \beta_3^{(s)} D_3(t)}_{S_t} + R_t$$

We can estimate the parameters $\beta = (\beta_0, \beta_1, \beta_1^{(s)}, \beta_2^{(s)}, \beta_3^{(s)})$ the same way we do for linear regression models:

$$\hat{\beta} = (X^T X)^{-1} (X^T y), \quad X \in \mathbb{R}^{T \times 5}$$

Where T is the length of the time-series.

1.5 Autocorrelation & autocorrelation function (ACF)

Definition 1.8 (Autocovariance & Autocorrelation).

Given a time-series $(X_t)_{t \geq 0}$, the autocovariance $\gamma(h)$ and autocorrelation $\rho(h)$ are defined as followed:

$$\begin{aligned} \gamma(h) &= Cov(X_{t+h}, X_t) \\ \rho(h) &= \frac{\gamma(h)}{Var(X_t)} = \frac{\gamma(h)}{\gamma(0)} \end{aligned}$$

When we are given a set of observations $(x_t)_{t=1}^n$, we can estimate γ, ρ by using their sample counterparts:

$$\begin{aligned} \hat{\gamma}(h) &= \frac{1}{n} \sum_{t=1}^{n-h} (x_{t+h} - \bar{x})(x_t - \bar{x}) \\ \hat{\rho}(h) &= \frac{\hat{\gamma}(h)}{\hat{\gamma}(0)} = \frac{\sum_{t=1}^{n-h} (x_{t+h} - \bar{x})(x_t - \bar{x})}{\sum_{t=1}^n (x_t - \bar{x})^2} \end{aligned}$$

Definition 1.9 (Autocorrelation function (ACF)).

The autocorrelation function, or correlogram is the plot of sample autocorrelation coefficients $\hat{\rho}(h)$ against $h = 0, 1, 2, \dots$.

2 Stationary time-series

2.1 Strict vs. Weak stationarity

Definition 2.1 (Mean, covariance and variance functions).

Given a time-series $(X_t)_{t \in T}$ (where T is a time domain), we have the following functions:

- Mean : $\mu(t) = \mathbb{E}[X_t]$, $t \in T$.
- Covariance : $\gamma(t, s) = \text{Cov}(X_t, X_s)$, $t, s \in T$.
- Variance : $\sigma^2(t) = \gamma(t, t) = \text{Var}(X_t)$, $t \in T$.

Definition 2.2 (Strict stationarity).

A time-series is said to be **strictly stationary** if, $\forall n \in \mathbb{Z}_+, h \in \mathbb{N}$, we have:

$$(X_1, \dots, X_n) \stackrel{d}{\simeq} (X_{1+h}, \dots, X_{n+h})$$

Meaning the two tuples are equal in distribution.

Remark : Note that in practice, strict stationarity is often hard to achieve. Hence, we have some relaxed conditions for stationarity.

Definition 2.3 (Weak stationarity).

A time-series $(X_t)_{t \geq 0}$ is said to be **weakly stationary** if the following two conditions are met:

- $\mu(t)$ does not depend on t .
- $\gamma(t, h)$ does not depend on t for all $h \in \mathbb{N}$.

A List of Definitions

1.1	Definition (Time series)	2
1.2	Definition (Multiplicative & Additive models)	2
1.3	Definition (Trend model)	2
1.4	Definition (Differencing operator)	2
1.5	Definition (Trend + Seasonality model)	3
1.6	Definition (Box-Cox transformation)	4
1.7	Definition (Dummy variables for seasonality)	4
1.8	Definition (Autocovariance & Autocorrelation)	5
1.9	Definition (Autocorrelation function (ACF))	5
2.1	Definition (Mean, covariance and variance functions)	6
2.2	Definition (Strict stationarity)	6
2.3	Definition (Weak stationarity)	6

B Important Theorems

C Important Corollaries

D Important Propositions

1.1	Binomial formula for $\nabla^n X_t$	3
1.2	Box-Cox as $\lambda \rightarrow 0$	4

E References

References

- [1] Rick Durrett. *Probability: Theory and Examples*. 4th. USA: Cambridge University Press, 2010. ISBN: 0521765390.
- [2] Erhan undefinedinlar. *Probability and Stochastics*. Springer New York, 2011. ISBN: 9780387878591. DOI: [10.1007/978-0-387-87859-1](https://doi.org/10.1007/978-0-387-87859-1). URL: <http://dx.doi.org/10.1007/978-0-387-87859-1>.
- [3] Wikipedia. *Vitali set* — *Wikipedia, The Free Encyclopedia*. <http://en.wikipedia.org/w/index.php?title=Vitali%20set&oldid=1187241923>. [Online; accessed 24-December-2023]. 2023.