

High Dimensional Probability Notes

Nong Minh Hieu¹

¹ School of Physical and Mathematical Sciences, Nanyang Technological University (NTU - Singapore)

Contents

1	Random variables	2
1.1	Basic inequalities	2
1.2	Limit Theorems	4
B	List of Definitions	5
B	Important Theorems	5
C	Important Corollaries	5
D	Important Propositions	5
E	References	6

1 Random variables

1.1 Basic inequalities

First, we revisit the definition of a random variable as well as some basic inequalities that we learned in introductory statistics.

Definition 1.1.

Random variable Let $(\Omega, \Sigma, \mathbb{P})$ be a probability space. A random variable X is defined as a mapping from the sample space Ω to \mathbb{R} :

$$X : \Omega \rightarrow \mathbb{R} \quad (1)$$

Σ is the σ -algebra containing the possible events (collection of subsets of Ω) and \mathbb{P} is a probability measure that assigns events with probabilities:

$$\mathbb{P} : \Sigma \rightarrow [0, 1] \quad (2)$$

For a given probability space $(\Omega, \Sigma, \mathbb{P})$ and a random variable $X : \Omega \rightarrow \mathbb{R}$, we will use the following basic notations throughout this note:

- $\|X\|_{L^p}$ - The p^{th} root of the p^{th} moment of the random variable X .

$$\|X\|_{L^p} = (\mathbb{E}|X|^p)^{1/p}, \quad p \in (0, \infty) \quad (3)$$

$$\|X\|_{L^\infty} = \text{ess sup } |X| \quad (4)$$

- $L^p(\Omega, \Sigma, \mathbb{P})$ - The space of random variables X satisfying:

$$L^p(\Omega, \Sigma, \mathbb{P}) = \left\{ X : \Omega \rightarrow \mathbb{R} \mid \|X\|_{L^p} < \infty \right\} \quad (5)$$

Some basic inequalities and identities:

- **1. Jensen's Inequality** - For a random variable X and a convex function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$, we have:

$$\varphi(\mathbb{E}X) \leq \mathbb{E}\varphi(X) \quad (6)$$

- **2. Monotonicity of L^p norm** - For a random variable X :

$$\|X\|_{L^p} \leq \|X\|_{L^q}, \quad 0 \leq p \leq q \leq \infty \quad (7)$$

- **3. Minkowski's Inequality** - For $1 \leq p \leq \infty$ and two random variables X, Y in $L^p(\Omega, \Sigma, \mathbb{P})$ space:

$$\|X + Y\|_{L^p} \leq \|X\|_{L^p} + \|Y\|_{L^p} \quad (8)$$

- **4. Holder's Inequality** - For $p, q \in [1, \infty]$ such that $1/p + 1/q = 1$. Then, for random variables $X \in L^p(\Omega, \Sigma, \mathbb{P})$ and $Y \in L^q(\Omega, \Sigma, \mathbb{P})$, we have:

$$|\mathbb{E}XY| \leq \|X\|_{L^p} \cdot \|Y\|_{L^q} \quad (9)$$

- **5. Markov's Inequality** - For a non-negative random variable X and $t > 0$, we have:

$$\mathbb{P}(X \geq t) \leq \frac{\mathbb{E}X}{t} \quad (10)$$

- **6. Chebyshev's Inequality** - For a random variable X with mean μ and variance σ^2 . Then, for any $t > 0$, we have:

$$\mathbb{P}(|X - \mu| \geq t) \leq \frac{\sigma^2}{t^2} \quad (11)$$

- **7. Integral Identity** - Let X be a non-negative random variable, we have:

$$\mathbb{E}X = \int_0^\infty \mathbb{P}(X > t) dt \quad (12)$$

Exercises

Exercise 1.1.1: Generalized Integral Identity

Let X be a random variable (not necessarily non-negative). Prove the following identity:

$$\mathbb{E}X = \int_0^\infty \mathbb{P}(X > t) dt - \int_{-\infty}^0 \mathbb{P}(X < t) dt \quad (13)$$

Solution (Exercise 1.1.1). _____

For $x \in \mathbb{R}$, using the basic integral identity, we have:

$$|x| = \int_0^\infty \mathbf{1}\{t < |x|\} dt$$

We consider the following cases:

- When $x < 0 \implies x = -|x|$:

$$x = - \int_0^\infty \mathbf{1}\{t < |x|\} dt = - \int_0^\infty \mathbf{1}\{t < -x\} dt = - \int_0^\infty \mathbf{1}\{-t > x\} dt = - \int_{-\infty}^0 \mathbf{1}\{t > x\} dt$$

- When $x \geq 0 \implies x = |x|$:

$$x = \int_0^\infty \mathbf{1}\{t < |x|\} dt = \int_0^\infty \mathbf{1}\{t < x\} dt$$

Therefore, for $x \in \mathbb{R}$, we can write:

$$x = \int_0^\infty \mathbf{1}\{t < x\} dt - \int_{-\infty}^0 \mathbf{1}\{t > x\} dt$$

Therefore, for a random variable X not necessarily non-negative, we have:

$$\begin{aligned} \mathbb{E}X &= \mathbb{E} \left[\int_0^\infty \mathbf{1}\{t < X\} dt - \int_{-\infty}^0 \mathbf{1}\{t > X\} dt \right] \\ &= \mathbb{E} \int_0^\infty \mathbf{1}\{t < X\} dt - \mathbb{E} \int_{-\infty}^0 \mathbf{1}\{t > X\} dt \\ &= \int_0^\infty \mathbb{E} \mathbf{1}\{t < X\} dt - \int_{-\infty}^0 \mathbb{E} \mathbf{1}\{t > X\} dt \\ &= \int_0^\infty \mathbb{P}(t < X) dt - \int_{-\infty}^0 \mathbb{P}(t > X) dt \end{aligned}$$

□.

Exercise 1.1.2: p^{th} -moments via tails

Let X be a random variable and $p \in (0, \infty)$. Show that:

$$\mathbb{E}|X|^p = \int_0^\infty pt^{p-1}\mathbb{P}(|X| > t)dt \quad (14)$$

Solution (Exercise 1.1.2).

Let X be a random variable that is not necessarily non-negative. Using the integral identity, we have:

$$\mathbb{E}|X|^p = \int_0^\infty \mathbb{P}(u < |X|^p)du$$

Let $t^p = u \implies pt^{p-1}dt = du$. Since we integrate u from $0 \rightarrow \infty$, we also integrate t from $0 \rightarrow \infty$ when changing the variables. Hence, we have:

$$\mathbb{E}|X|^p = \int_0^\infty \mathbb{P}(t^p < |X|^p)pt^{p-1}dt = \int_0^\infty \mathbb{P}(t < |X|)pt^{p-1}dt$$

Hence, we obtained the desired identity. □.

1.2 Limit Theorems

B List of Definitions

1.1 Definition	2
--------------------------	---

B Important Theorems

C Important Corollaries

D Important Propositions

E References