# High Dimensional Probability Notes

## Nong Minh $\mathrm{Hieu^1}$

 $^{1}$  School of Physical and Mathematical Sciences, Nanyang Technological University (NTU - Singapore)

## Contents

1	Random variables	2
	1.1 Basic inequalities	2
A	List of Definitions	3
В	Important Theorems	3
$\mathbf{C}$	Important Corollaries	3
D	Important Propositions	3
${f E}$	References	4

#### 1 Random variables

#### 1.1 Basic inequalities

First, we revisit the definition of a random variable as well as some basic inequalities that we learned in introductory statistics.

**Definition 1.1** (Random variable). \_

Let  $(\Omega, \Sigma, \mathbb{P})$  be a probability space. A random variable X is defined as a mapping from the sample space  $\Omega$  to  $\mathbb{R}$ :

$$X: \Omega \to \mathbb{R} \tag{1}$$

 $\Sigma$  is the  $\sigma$ -algebra containing the possible events (collection of subsets of  $\Omega$ ) and  $\mathbb{P}$  is a probability measure that assigns events with probabilities:

$$\mathbb{P}: \Sigma \to [0, 1] \tag{2}$$

For a given probability space  $(\Omega, \Sigma, \mathbb{P})$  and a random variable  $X : \Omega \to \mathbb{R}$ , we will use the following basic notations throughout this note:

•  $||X||_{L^p}$  - The  $p^{th}$  root of the  $p^{th}$  moment of the random variable X.

$$||X||_{L^p} = (\mathbb{E}|X|^p)^{1/p}, \ p \in (0, \infty)$$
 (3)

$$||X||_{L^{\infty}} = \operatorname{ess\,sup}|X| \tag{4}$$

•  $L^p(\Omega, \Sigma, \mathbb{P})$  - The space of random variables X satisfying:

$$L^{p}(\Omega, \Sigma, \mathbb{P}) = \left\{ X : \Omega \to \mathbb{R} \middle| \|X\|_{L^{p}} < \infty \right\}$$
 (5)

Some basic inequalities and identities:

• 1. Jensen's Inequality - For a random variable X and a convex function  $\varphi : \mathbb{R} \to \mathbb{R}$ , we have:

$$\varphi(\mathbb{E}X) \le \mathbb{E}\varphi(X) \tag{6}$$

• 2. Monotonicity of  $L^p$  norm - For a random variable X:

$$||X||_{L^p} \le ||X||_{L^q}, \ 0 \le p \le q \le \infty \tag{7}$$

• 3. Minkowski's Inequality - For  $1 \le p \le \infty$  and two random variables  $X,Y \in L^p(\Omega,\Sigma,\mathbb{P})$ :

$$||X + Y||_{L^p} \le ||X||_{L^p} + ||Y||_{L^p} \tag{8}$$

• 4. Holder's Inequality - For  $p, q \in [1, \infty]$  such that 1/p + 1/q = 1. Then, for  $X \in L^p(\Omega, \Sigma, \mathbb{P})$  and  $Y \in L^q(\Omega, \Sigma, \mathbb{P})$ , we have:

$$|\mathbb{E}XY| \le ||X||_{L^p} \cdot ||Y||_{L^q} \tag{9}$$

• 5. Integral identity - Let X be a non-negative random variable, we have:

$$\mathbb{E}X = \int_0^\infty \mathbb{P}(X > t)dt \tag{10}$$

$\mathbf{A}$	List of Definitions	
1	1 Definition (Random variable)	2
В	Important Theorems	
$\mathbf{C}$	Important Corollaries	
D	Important Propositions	

### E References

#### References

- [1] Rick Durrett. *Probability: Theory and Examples.* 4th. USA: Cambridge University Press, 2010. ISBN: 0521765390.
- [2] Erhan undefinedinlar. *Probability and Stochastics*. Springer New York, 2011. ISBN: 9780387878591. DOI: 10.1007/978-0-387-87859-1. URL: http://dx.doi.org/10.1007/978-0-387-87859-1.
- [3] Wikipedia. Vitali set Wikipedia, The Free Encyclopedia. http://en.wikipedia.org/w/index.php?title=Vitali%20set&oldid=1187241923. [Online; accessed 24-December-2023]. 2023.