

# High Dimensional Probability Notes

Nong Minh Hieu<sup>1</sup>

<sup>1</sup> School of Physical and Mathematical Sciences, Nanyang Technological University (NTU - Singapore)

## Contents

<b>1</b>	<b>Random variables</b>	<b>2</b>
1.1	Basic Inequalities . . . . .	2
1.2	Limit Theorems . . . . .	4
1.2.1	Weak Law of Large Numbers . . . . .	4
1.2.2	Strong Law of Large Numbers . . . . .	6
1.2.3	Central Limit Theorem . . . . .	8
1.3	Convergence of Random Variables . . . . .	8
1.3.1	Convergence in Distribution . . . . .	8
1.3.2	Convergence in Probability . . . . .	9
1.3.3	Convergence in $L^p$ norm . . . . .	10
1.3.4	Almost-sure Convergence . . . . .	10
<b>2</b>	<b>Statistical Inference</b>	<b>12</b>
2.1	Sufficiency & Likelihood . . . . .	12
2.1.1	Sufficiency . . . . .	12
2.1.2	Likelihood . . . . .	12
2.2	Point Estimation . . . . .	12
2.2.1	Bias, Variance, Consistency and MSE . . . . .	12
2.2.2	Sufficient Statistics & Rao-Blackwell Theorem . . . . .	12
2.2.3	Estimator Variance & Cramer-Rao Lower Bound . . . . .	12
2.2.4	Maximum Likelihood Estimation (MLE) . . . . .	12
<b>B</b>	<b>List of Definitions</b>	<b>13</b>
<b>B</b>	<b>Important Theorems</b>	<b>13</b>
<b>C</b>	<b>Important Propositions</b>	<b>13</b>
<b>D</b>	<b>References</b>	<b>14</b>

# 1 Random variables

## 1.1 Basic Inequalities

First, we revisit the definition of a random variable as well as some basic inequalities that we learned in introductory statistics.

**Definition 1.1** (Random variable).

Let  $(\Omega, \Sigma, \mathbb{P})$  be a probability space. A random variable  $X$  is defined as a mapping from the sample space  $\Omega$  to  $\mathbb{R}$ :

$$X : \Omega \rightarrow \mathbb{R} \quad (1)$$

$\Sigma$  is the  $\sigma$ -algebra containing the possible events (collection of subsets of  $\Omega$ ) and  $\mathbb{P}$  is a probability measure that assigns events with probabilities:

$$\mathbb{P} : \Sigma \rightarrow [0, 1] \quad (2)$$

For a given probability space  $(\Omega, \Sigma, \mathbb{P})$  and a random variable  $X : \Omega \rightarrow \mathbb{R}$ , we will use the following basic notations throughout this note:

- $\|X\|_{L^p}$  - The  $p^{th}$  root of the  $p^{th}$  moment of the random variable  $X$ .

$$\|X\|_{L^p} = (\mathbb{E}|X|^p)^{1/p}, \quad p \in (0, \infty) \quad (3)$$

$$\|X\|_{L^\infty} = \text{ess sup } |X| \quad (4)$$

- $L^p(\Omega, \Sigma, \mathbb{P})$  - The space of random variables  $X$  satisfying:

$$L^p(\Omega, \Sigma, \mathbb{P}) = \left\{ X : \Omega \rightarrow \mathbb{R} \mid \|X\|_{L^p} < \infty \right\} \quad (5)$$

Some basic inequalities and identities:

- **1. Jensen's Inequality** - For a random variable  $X$  and a convex function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ , we have:

$$\varphi(\mathbb{E}X) \leq \mathbb{E}\varphi(X) \quad (6)$$

- **2. Monotonicity of  $L^p$  norm** - For a random variable  $X$ :

$$\|X\|_{L^p} \leq \|X\|_{L^q}, \quad 0 \leq p \leq q \leq \infty. \quad (7)$$

- **3. Minkowski's Inequality** - For  $1 \leq p \leq \infty$  and two random variables  $X, Y$  in  $L^p(\Omega, \Sigma, \mathbb{P})$  space:

$$\|X + Y\|_{L^p} \leq \|X\|_{L^p} + \|Y\|_{L^p}. \quad (8)$$

- **4. Holder's Inequality** - For  $p, q \in [1, \infty]$  such that  $1/p + 1/q = 1$ . Then, for random variables  $X \in L^p(\Omega, \Sigma, \mathbb{P})$  and  $Y \in L^q(\Omega, \Sigma, \mathbb{P})$ , we have:

$$|\mathbb{E}XY| \leq \|X\|_{L^p} \cdot \|Y\|_{L^q}. \quad (9)$$

- **5. Markov's Inequality** - For a non-negative random variable  $X$  and  $t > 0$ , we have:

$$\mathbb{P}(X \geq t) \leq \frac{\mathbb{E}X}{t}. \quad (10)$$

We can also generalize Markov's Inequality for  $p^{th}$  moment:

$$\mathbb{P}(|X| \geq t) \leq \frac{\mathbb{E}[|X|^p]}{t^p}, \quad \forall t > 0, p \in [2, \infty). \quad (11)$$

- **6. Chebyshev's Inequality** - For a random variable  $X$  with mean  $\mu$  and variance  $\sigma^2$ . Then, for any  $t > 0$ , we have:

$$\mathbb{P}(|X - \mu| \geq t) \leq \frac{\sigma^2}{t^2}. \quad (12)$$

- **7. Integral Identity** - Let  $X$  be a non-negative random variable, we have:

$$\mathbb{E}X = \int_0^\infty \mathbb{P}(X > t) dt. \quad (13)$$

## Exercises

### Exercise 1.1.1: Generalized Integral Identity

Let  $X$  be a random variable (not necessarily non-negative). Prove the following identity:

$$\mathbb{E}X = \int_0^\infty \mathbb{P}(X > t) dt - \int_{-\infty}^0 \mathbb{P}(X < t) dt. \quad (14)$$

**Solution** (Exercise 1.1.1).

For  $x \in \mathbb{R}$ , using the basic integral identity, we have:

$$|x| = \int_0^\infty \mathbf{1}\{t < |x|\} dt$$

We consider the following cases:

- When  $x < 0 \implies x = -|x|$ :

$$x = - \int_0^\infty \mathbf{1}\{t < |x|\} dt = - \int_0^\infty \mathbf{1}\{t < -x\} dt = - \int_0^\infty \mathbf{1}\{-t > x\} dt = - \int_{-\infty}^0 \mathbf{1}\{t > x\} dt.$$

- When  $x \geq 0 \implies x = |x|$ :

$$x = \int_0^\infty \mathbf{1}\{t < |x|\} dt = \int_0^\infty \mathbf{1}\{t < x\} dt.$$

Therefore, for  $x \in \mathbb{R}$ , we can write:

$$x = \int_0^\infty \mathbf{1}\{t < x\} dt - \int_{-\infty}^0 \mathbf{1}\{t > x\} dt.$$

Therefore, for a random variable  $X$  not necessarily non-negative, we have:

$$\begin{aligned} \mathbb{E}X &= \mathbb{E} \left[ \int_0^\infty \mathbf{1}\{t < X\} dt - \int_{-\infty}^0 \mathbf{1}\{t > X\} dt \right] \\ &= \mathbb{E} \int_0^\infty \mathbf{1}\{t < X\} dt - \mathbb{E} \int_{-\infty}^0 \mathbf{1}\{t > X\} dt \\ &= \int_0^\infty \mathbb{E} \mathbf{1}\{t < X\} dt - \int_{-\infty}^0 \mathbb{E} \mathbf{1}\{t > X\} dt \\ &= \int_0^\infty \mathbb{P}(t < X) dt - \int_{-\infty}^0 \mathbb{P}(t > X) dt. \end{aligned}$$

□.

**Exercise 1.1.2:  $p^{th}$ -moments via tails**

Let  $X$  be a random variable and  $p \in (0, \infty)$ . Show that:

$$\mathbb{E}|X|^p = \int_0^\infty pt^{p-1}\mathbb{P}(|X| > t)dt. \quad (15)$$

**Solution** (Exercise 1.1.2). \_\_\_\_\_

Let  $X$  be a random variable that is not necessarily non-negative. Using the integral identity, we have:

$$\mathbb{E}|X|^p = \int_0^\infty \mathbb{P}(u < |X|^p)du.$$

Let  $t^p = u \implies pt^{p-1}dt = du$ . Since we integrate  $u$  from  $0 \rightarrow \infty$ , we also integrate  $t$  from  $0 \rightarrow \infty$  when changing the variables. Hence, we have:

$$\mathbb{E}|X|^p = \int_0^\infty \mathbb{P}(t^p < |X|^p)pt^{p-1}dt = \int_0^\infty \mathbb{P}(t < |X|)pt^{p-1}dt.$$

Hence, we obtained the desired identity. □.

**1.2 Limit Theorems****1.2.1 Weak Law of Large Numbers****Theorem 1.1: Weak Law of Large Numbers (WLLN)**

Let  $X_1, \dots, X_N$  be *i.i.d* random variables with mean  $\mu$ . Consider the sum:

$$S_N = X_1 + \dots + X_N$$

Then, the sample mean **converges to  $\mu$  in probability** ( $S_N/N \xrightarrow{p} \mu$ ):

$$\lim_{N \rightarrow \infty} \mathbb{P}\left(|S_N/N - \mu| > \epsilon\right) = 0, \quad \forall \epsilon > 0 \quad (16)$$

**Proof** (Weak Law of Large Numbers (WLLN)). \_\_\_\_\_

We split the proof into two sections corresponding to the assumptions of finite variance and non-finite variance.

1. **Finite variance case:** Suppose that  $\text{Var}X_i = \sigma^2 < \infty$  for all  $1 \leq i \leq N$ . Let  $\bar{X} = S_N/N$ . Then,  $\bar{X}$  is a random variable with the following mean and variance:

$$\mathbb{E}\bar{X} = \mu \quad \text{and} \quad \text{Var}\bar{X} = \frac{\sigma^2}{N}.$$

Hence, by the Chebyshev's inequality, we have:

$$\mathbb{P}\left(|S_N/N - \mu| > \epsilon\right) = \mathbb{P}\left(|\bar{X} - \mu| > \epsilon\right) \leq \frac{\sigma^2}{N\epsilon^2}.$$

Therefore, we have:

$$\lim_{N \rightarrow \infty} \mathbb{P}\left(|S_N/N - \mu| > \epsilon\right) \leq \lim_{N \rightarrow \infty} \frac{\sigma^2}{N\epsilon^2} = 0.$$

Hence, we have  $\lim_{N \rightarrow \infty} \mathbb{P}\left(|S_N/N - \mu| > \epsilon\right) = 0$  and we obtained **(WLLN)**.

2. **Non-finite variance case:** In this case, we rely on the Levy Continuity Theorem (**LCT**), which relies on the convergence of the characteristic function. For  $n \geq 1$ , define the sequence of random variable  $Y_n = S_n/n$ . Hence, we have:

$$\begin{aligned}\varphi_{Y_n}(t) &= \varphi_{S_n/n}(t) \\ &= \varphi_{S_n}(t/n) \\ &= \prod_{i=1}^n \varphi_{X_i}(t/n) = \left[ \varphi_X(t/n) \right]^n,\end{aligned}$$

Where  $X = X_1 = \dots = X_n$ . By Taylor's expansion, we have:

$$\varphi_X(t/n) = 1 + \frac{it\mathbb{E}[X]}{n} + \mathcal{O}(1/n^2) = 1 + \frac{it\mu}{n} + \mathcal{O}(1/n^2).$$

Hence, we have:

$$\lim_{n \rightarrow \infty} \varphi_{Y_n}(t) = \lim_{n \rightarrow \infty} \left( 1 + \frac{it\mu}{n} + \mathcal{O}(1/n^2) \right)^n = e^{it\mu}.$$

Therefore, by (**LCT**), we have  $Y_n \xrightarrow{p} \mu$ .

**Remark 1.1** (Taylor expansion of Moment Generating and Characteristic Functions). —————  
Given a random variable  $X$ . For reference, the following are the Taylor expansions of the Moment Generating Function  $M_X(t)$  and the Characteristic Function  $\varphi_X(t)$ :

$$\begin{aligned}M_X(t) &= \mathbb{E}[e^{tX}] = 1 + \sum_{n=1}^{\infty} \frac{t^n}{n!} \mathbb{E}[X^n], \\ \varphi_X(t) &= \mathbb{E}[e^{itX}] = 1 + \sum_{n=1}^{\infty} \frac{(it)^n}{n!} \mathbb{E}[X^n].\end{aligned}\tag{17}$$

For the sake of my laziness, here are the Taylor expansion for the first three terms of both the MGF and the CF:

$$\begin{aligned}M_X(t) &= 1 + t\mathbb{E}[X] + \frac{t^2}{2}\mathbb{E}[X^2] + \mathcal{O}(t^3), \\ \varphi_X(t) &= 1 + it\mathbb{E}[X] - \frac{t^2}{2}\mathbb{E}[X^2] + \mathcal{O}(t^3).\end{aligned}\tag{18}$$

□.

### Theorem 1.2: Levy Continuity Theorem (**LCT**)

Let  $X_1, X_2, \dots$  be *i.i.d* random variables. Then:

$$\forall t \in \mathbb{R} : \lim_{n \rightarrow \infty} \varphi_{X_n}(t) = \varphi_X(t) \iff X_n \xrightarrow{d} X,\tag{19}$$

for some random variable  $X$ . In a special case where  $X = c$  for some  $c \in \mathbb{R}$ , we have:

$$\forall t \in \mathbb{R} : \lim_{n \rightarrow \infty} \varphi_{X_n}(t) = e^{itc} \iff X_n \xrightarrow{p} c.\tag{20}$$

**Proof** (Levy Continuity Theorem (**LCT**)). —————  
The proof for (**LCT**) can be found in Gut 2004, Section 9.1, Theorem 9.1 and Corollary 9.1 □.

### 1.2.2 Strong Law of Large Numbers

#### Theorem 1.3: Strong Law of Large Numbers (SLLN)

Let  $X_1, \dots, X_N$  be *i.i.d* random variables with mean  $\mu$ . Consider the sum:

$$S_N = X_1 + \dots + X_N$$

Then, the sample mean **converges to  $\mu$  almost surely** ( $S_N/N \xrightarrow{a.s} \mu$ ):

$$\mathbb{P}\left(\limsup_{N \rightarrow \infty} |S_N/N - \mu| > \epsilon\right) = 0, \quad \forall \epsilon > 0 \quad (21)$$

**Proof** (Strong Law of Large Numbers (SLLN)).

For the sake of simplicity, we will present the proof for (SLLN) with an additional assumption that  $\mathbb{E}[|X_n|^4] < \infty, \forall n \geq 1$ . The proof for the general case of (SLLN) (also called the Kolmogorov Strong Law) can be found in Gut 2004, Section 6, Theorem 6.1. For convenience, we assume the following:

1.  $\mathbb{E}[|X_n|^4] = K < \infty$ .
2.  $\mathbb{E}[X_n] = 0$ . For non-zero mean case, we can set  $Y_n = X_n - \mu$  and repeat the same arguments made below.

We aim to prove that  $\mathbb{P}\left(\limsup_{N \rightarrow \infty} |S_N/N| > \epsilon\right) = 0$  for any  $\epsilon > 0$ . Firstly, use the Multinomial formula to expand  $\mathbb{E}[S_n]$ . The expansion will contain the terms in the following forms:

$$X_i^2, X_i^3 X_j, X_i^2 X_j^2, X_i^2 X_j X_k, X_i X_j X_k X_\ell,$$

where  $i, j, k, \ell$  are distinct indices. By independence, we have:

$$\mathbb{E}[X_i^3 X_j] = \mathbb{E}[X_i^2 X_j X_k] = \mathbb{E}[X_i X_j X_k X_\ell] = 0.$$

As a result, we have the following remaining terms by the Multinomial formula:

$$\begin{aligned} \mathbb{E}[S_n^4] &= \sum_{i=1}^n \mathbb{E}[X_i^4] + \binom{4}{2} \sum_{1 \leq i < j \leq n} \mathbb{E}[X_i^2 X_j^2] \\ &= \sum_{i=1}^n \mathbb{E}[X_i^4] + 6 \underbrace{\sum_{1 \leq i < j \leq n} \mathbb{E}[X_i^2 X_j^2]}_{n(n-1)/2 \text{ terms}} \\ &= nK + 3n(n-1)\mathbb{E}[X_i^2 X_j^2]. \end{aligned}$$

By independence, we have  $\mathbb{E}[X_i^2 X_j^2] = \mathbb{E}[X_i^2] \mathbb{E}[X_j^2]$  and for any  $1 \leq i \leq n$ . Furthermore, we have  $\mathbb{E}[X_i^2] = \text{Var}(X_i) + \mu^2 = \sigma^2 + \mu^2$ . Therefore:

$$\mathbb{E}[S_n^4] = nK + 3n(n-1)(\sigma^2 + \mu^2) < nK + 3n^2(\sigma^2 + \mu^2).$$

Applying Markov's Inequality with the fourth moment, we have:

$$\begin{aligned} \mathbb{P}(|S_n/n| \geq \epsilon) &= \mathbb{P}(|S_n| \geq n\epsilon) \\ &\leq \frac{\mathbb{E}[S_n^4]}{n^4 \epsilon^4} \\ &< \frac{K}{n^3 \epsilon^4} + \frac{3(\sigma^2 + \mu^2)}{n^2}. \end{aligned}$$

Therefore, we have:

$$\sum_{n=1}^{\infty} \mathbb{P}(|S_n/n| \geq \epsilon) < \frac{K}{\epsilon^4} \sum_{n=1}^{\infty} n^{-3} + 3(\sigma^2 + \mu^2) \sum_{n=1}^{\infty} n^{-2} < \infty \quad (22)$$

Finally, by the Borel-Cantelli Lemma (**BCL**), we have:

$$\mathbb{P}\left(\limsup_{n \rightarrow \infty} |S_n/n| \geq \epsilon\right) = 0, \quad \forall \epsilon > 0.$$

□.

#### Theorem 1.4: Borel-Cantelli Lemma (**BCL**)

**1. First Borel-Cantelli Lemma:** Given a probability space  $(X, \mathcal{S}, \mathbb{P})$  and a sequence  $\{A_n\}_{n=1}^{\infty} \subset \mathcal{S}$ . If  $\sum_{n=1}^{\infty} \mathbb{P}(A_n) < \infty$ , we have:

$$\mathbb{P}\left(\limsup_{n \rightarrow \infty} A_n\right) = 0. \quad (23)$$

**2. Second Borel-Cantelli Lemma:** On the other hand, if  $\sum_{n=1}^{\infty} \mathbb{P}(E_n) = \infty$ , we have:

$$\mathbb{P}\left(\limsup_{n \rightarrow \infty} A_n\right) = 1. \quad (24)$$

**Proof** (Borel-Cantelli Lemma (**BCL**)).

We focus on proving the first Borel-Cantelli lemma. We define another sequence of  $\mathcal{S}$ -measurable sets  $\{B_n\}_{n=1}^{\infty}$  such that:

$$B_n = \bigcup_{k=n}^{\infty} A_k.$$

Hence, we have  $B_{\ell+1} \subset B_{\ell}$  for every  $\ell \geq 1$ . In other words,  $B_n$  is a decreasing sequence of  $\mathcal{S}$ -measurable sets. By continuity of measure, we have:

$$\begin{aligned} \mathbb{P}\left(\lim_{n \rightarrow \infty} B_n\right) &= \lim_{n \rightarrow \infty} \mathbb{P}(B_n) \\ &= \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \mathbb{P}(A_k) \quad (\text{By additivity}) \\ &= \sum_{i=1}^{\infty} \mathbb{P}(A_i) - \lim_{n \rightarrow \infty} \sum_{k=1}^n \mathbb{P}(A_k) \\ &= 0. \end{aligned}$$

Furthermore, we have:

$$\mathbb{P}\left(\lim_{n \rightarrow \infty} B_n\right) = \mathbb{P}\left(\lim_{n \rightarrow \infty} \bigcup_{k=n}^{\infty} A_k\right) = \mathbb{P}\left(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k\right) = \mathbb{P}\left(\limsup_{n \rightarrow \infty} A_n\right).$$

Hence proved the first Borel-Cantelli Lemma. To prove the second Borel-Cantelli Lemma, we prove the following:

$$\begin{aligned} 1 - \mathbb{P}\left(\limsup_{n \rightarrow \infty} A_n\right) &= \mathbb{P}\left(\left\{\limsup_{n \rightarrow \infty} A_n\right\}^c\right) \\ &= \mathbb{P}\left(\liminf_{n \rightarrow \infty} A_n^c\right) = 0. \end{aligned}$$

□.

### 1.2.3 Central Limit Theorem

#### Theorem 1.5: Central Limit Theorem (CLT)

Let  $X_1, \dots, X_n$  be a sequence of *i.i.d* random variables with expected value  $\mu$  and finite variance  $\sigma^2$ . Then, we have:

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty, \quad (25)$$

where  $\bar{X}_n = S_n/n$  and  $\mathcal{N}(0, 1)$  is the standard normal distribution.

**Proof** (Central Limit Theorem (CLT)).

We prove this via the Characteristic Function. Let  $\bar{Z}_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}}$ , notice that:

$$\bar{Z}_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{X_i - \mu}{\sigma},$$

Let  $Z_i = X_i - \mu$  for  $1 \leq i \leq n$  and suppose  $Z = Z_1 = \dots = Z_n$ , we have:

$$\begin{aligned} \varphi_{\bar{Z}_n}(t) &= \varphi_{\sum_{i=1}^n Z_i} \left( \frac{t}{\sqrt{n}} \right) = \left[ \varphi_Z \left( \frac{t}{\sqrt{n}} \right) \right]^n \\ &= \left[ 1 + \frac{it\mathbb{E}[Z]}{\sqrt{n}} - \frac{t^2}{2n} \mathbb{E}[Z^2] + \mathcal{O}(1/n) \right]^n \quad (\text{Taylor's Expansion}) \\ &= \left[ 1 - \frac{t^2}{2n} + \mathcal{O}(1/n) \right]^n. \end{aligned}$$

The final equality comes from the fact that  $\mathbb{E}[Z] = 0$  and  $\mathbb{E}[Z^2] = \mathbb{E}[Z]^2 + \text{Var}(Z) = 1$ . Finally, we have:

$$\lim_{n \rightarrow \infty} \varphi_{\bar{Z}_n}(t) = \lim_{n \rightarrow \infty} \left[ 1 - \frac{t^2}{2n} + \mathcal{O}(1/n) \right]^n = e^{-t^2/2}.$$

Since  $e^{-t^2/2}$  is the Characteristic Function of the standard normal distribution, by (LCT), we have  $\bar{Z}_n \xrightarrow{d} \mathcal{N}(0, 1)$ .  $\square$ .

## 1.3 Convergence of Random Variables

In this section, we revise the modes of convergence in random variables.

### 1.3.1 Convergence in Distribution

**Definition 1.2** (Convergence in Distribution).

Given a sequence of real-valued random variables  $X_1, X_2, \dots$  with CDFs  $F_1, F_2, \dots$ . We say that the sequence converges in distribution to a random variable  $X$  with CDF  $F$ , denoted  $X_n \xrightarrow{d} X$  if:

$$\lim_{n \rightarrow \infty} F_n(x) = F(x), \quad (26)$$

for all  $x \in \mathbb{R}$  at which  $F$  is continuous. Convergence in distribution can also be referred to as weak convergence in measure theory.



### 1.3.2 Convergence in Probability

**Definition 1.3** (Convergence in Probability). 

---

Given a sequence of real-valued random variables  $X_1, X_2, \dots$ . We say that the sequence converges in probability to a random variable  $X$ , denoted  $X_n \xrightarrow{p} X$  if:

$$\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X| \geq \epsilon) = 0, \quad \forall \epsilon > 0. \quad (27)$$

We also refer to convergence in probability as convergence in measure in measure theory.

**Proposition 1.1:**  $X_n \xrightarrow{p} X \implies X_n \xrightarrow{d} X$

Let  $X$  and the sequence  $X_1, X_2, \dots$  be real-valued random variables. If  $X_n \xrightarrow{p} X$ , then  $X_n \xrightarrow{d} X$ .

**Proof** (Proposition 1.1). 

---

We first prove the following claim: Let  $X, Y$  be random variables,  $a \in \mathbb{R}$  and  $\epsilon > 0$ , the inequality  $\mathbb{P}(Y \leq a) \leq \mathbb{P}(X \leq a + \epsilon) + \mathbb{P}(|Y - X| \geq \epsilon)$  holds. We have:

$$\begin{aligned} \mathbb{P}(Y \leq a) &= \mathbb{P}(Y \leq a, X \leq a + \epsilon) + \mathbb{P}(Y \leq a, X \geq a + \epsilon) \\ &\leq \mathbb{P}(X \leq a + \epsilon) + \mathbb{P}(Y - X \leq a - X, a - X \leq -\epsilon) \\ &\leq \mathbb{P}(X \leq a + \epsilon) + \mathbb{P}(Y - X \leq -\epsilon) \\ &\leq \mathbb{P}(X \leq a + \epsilon) + \mathbb{P}(Y - X \leq -\epsilon) + \mathbb{P}(Y - X \geq \epsilon) \\ &= \mathbb{P}(X \leq a + \epsilon) + \mathbb{P}(|Y - X| \geq \epsilon). \end{aligned}$$

Using the above inequality, we have:

$$\mathbb{P}(X \leq a - \epsilon) - \mathbb{P}(|X_n - X| \geq \epsilon) \leq \mathbb{P}(X_n \leq a) \leq \mathbb{P}(X \leq a + \epsilon) + \mathbb{P}(|X_n - X| \geq \epsilon).$$

Taking limits as  $n \rightarrow \infty$  from both sides, we have:

$$F_X(a - \epsilon) \leq \lim_{n \rightarrow \infty} F_{X_n}(a) \leq F_X(a + \epsilon).$$

Taking  $\epsilon \rightarrow 0^+$ , we have  $\lim_{n \rightarrow \infty} F_{X_n}(a) = F_X(a)$ . □.

**Proposition 1.2:**  $X_n \xrightarrow{d} c \iff X_n \xrightarrow{p} c$

Let  $c \in \mathbb{R}$  be a constant and  $X_1, X_2, \dots$  be a sequence of real-valued random variables. Then,  $X_n \xrightarrow{d} c \iff X_n \xrightarrow{p} c$ .

**Proof** (Proposition 1.2, Pishro-Nik 2014). 

---

Since  $X_n \xrightarrow{d} c$ , we immediately have the following:

$$\begin{aligned} \lim_{n \rightarrow \infty} F_{X_n}(c - \epsilon) &= 0, \\ \lim_{n \rightarrow \infty} F_{X_n}(c + \epsilon/2) &= 1. \end{aligned}$$

Then, for any  $\epsilon > 0$ , we have:

$$\begin{aligned}
\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - c| \geq \epsilon) &= \lim_{n \rightarrow \infty} \mathbb{P}[\mathbb{P}(X_n \leq c - \epsilon) + \mathbb{P}(X_n \geq c + \epsilon)] \\
&= \underbrace{\lim_{n \rightarrow \infty} F_{X_n}(c - \epsilon)}_{=0} + \lim_{n \rightarrow \infty} \mathbb{P}(X_n \geq c + \epsilon) \\
&\leq \lim_{n \rightarrow \infty} \mathbb{P}(X_n \geq c + \epsilon/2) \\
&= 1 - \underbrace{\lim_{n \rightarrow \infty} F_{X_n}(c + \epsilon/2)}_{=1} \\
&= 0.
\end{aligned}$$

From the above, we have  $\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - c| \geq \epsilon) = 0$  and  $X_n \xrightarrow{P} c$ .  $\square$ .

### 1.3.3 Convergence in $L^p$ norm

**Definition 1.4** (Convergence in  $L^p$  norm). 

---

Given a sequence of random variables  $X_1, X_2, \dots$  and a real number  $p \in [1, \infty)$ . We say that the sequence converges in  $L^p$  norm to a random variable  $X$ , denoted as  $X_n \xrightarrow{L^p} X$  if:

$$\lim_{n \rightarrow \infty} \mathbb{E}|X_n - X|^p = 0. \quad (28)$$

**Proposition 1.3:**  $X_n \xrightarrow{L^p} X \implies X_n \xrightarrow{P} X$

Let  $p \geq 1$  and  $X_1, X_2, \dots$  be a sequence of real-valued random variables. Let  $X$  be a random variable, then,  $X_n \xrightarrow{L^p} X \implies X_n \xrightarrow{P} X$ .

**Proof** (Proposition 1.3). 

---

Let  $\epsilon > 0$ , we have:

$$\begin{aligned}
\mathbb{P}(|X_n - X| \geq \epsilon) &= \mathbb{P}(|X_n - X|^p \geq \epsilon^p) \quad (p \geq 1) \\
&\leq \frac{\mathbb{E}|X_n - X|^p}{\epsilon^p}. \quad (\text{Markov's Inequality})
\end{aligned}$$

Taking the limits from both sides, we have  $\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X| \geq \epsilon) = 0$  and  $X_n \xrightarrow{P} X$ .  $\square$ .

### 1.3.4 Almost-sure Convergence

**Definition 1.5** (Convergence almost-surely). 

---

Let  $X_1, X_2, \dots$  be a sequence of real-valued random variables that map from a sample space  $\Omega$ . Let  $X$  also be a real-valued random variable. We say that  $X_n$  converges almost surely to  $X$ , denoted as  $X_n \xrightarrow{a.s.} X$ , if:

$$\mathbb{P}\left(\limsup_{n \rightarrow \infty} E_n\right) = 0 \quad \text{where} \quad E_n = \left\{\omega \in \Omega : |X_n(\omega) - X(\omega)| \geq \epsilon\right\}.$$

**Remark 1.2** (Consequence of **(BCL)**). 

---

$$\sum_{n=1}^{\infty} \mathbb{P}(E_n) < \infty \implies X_n \xrightarrow{a.s.} X. \quad (29)$$

**Proposition 1.4:**  $X_n \xrightarrow{a.s.} X \implies X_n \xrightarrow{p} X$

Let  $X_1, X_2, \dots$  be a sequence of real-valued random variables and also let  $X$  be a real valued random variables. If  $X_n \xrightarrow{a.s.} X$  then  $X_n \xrightarrow{p} X$ .

**Proof** (Proposition 1.4). 

---

Let  $f_n : \Omega \rightarrow \mathbb{R}_+$  be a sequence of nonnegative Borel-measurable functions such that  $f_n(\omega) = |X_n(\omega) - X(\omega)|$ . By Fatou's Lemma (reverse), we have:

$$\begin{aligned} \underbrace{\mathbb{P}\left(\limsup_{n \rightarrow \infty} \{\omega \in \Omega : |X_n(\omega) - X(\omega)| \geq \epsilon\}\right)}_{=0} &= \int f_n d\mathbb{P} \\ &\geq \limsup_{n \rightarrow \infty} \mathbb{P}(|X_n - X| \geq \epsilon) \\ &\geq \lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X| \geq \epsilon). \end{aligned}$$

Hence, we have  $\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X| \geq \epsilon) = 0$  and  $X_n \xrightarrow{p} X$ .  $\square$ .

**Theorem 1.6: Continuous Mapping Theorem (CMT)**

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous function and  $X_1, X_2, \dots$  be a sequence of real-valued random variables. Then, the following statements hold true:

1.  $X_n \xrightarrow{d} X \implies f(X_n) \xrightarrow{d} f(X)$ .
2.  $X_n \xrightarrow{p} X \implies f(X_n) \xrightarrow{p} f(X)$ .
3.  $X_n \xrightarrow{a.s.} X \implies f(X_n) \xrightarrow{a.s.} f(X)$ .

**Proof** (Continuous Mapping Theorem **(CMT)**). 

---

Since almost-sure convergence implies the other two modes of convergence, we only have to handle the almost-sure convergence case. Since  $f$  is continuous, for any  $\omega \in \Omega$  such that  $X_n(\omega) \rightarrow X(\omega)$ , we have  $f(X_n(\omega)) \rightarrow f(X(\omega))$ . Therefore, we have:

$$\{\omega \in \Omega : X_n(\omega) \rightarrow X(\omega)\} \subseteq \{\omega \in \Omega : f(X_n(\omega)) \rightarrow f(X(\omega))\}.$$

Therefore, we have:

$$\begin{aligned} &\mathbb{P}\left(\limsup_{n \rightarrow \infty} \left\{\omega \in \Omega : |f(X_n(\omega)) - f(X(\omega))| \leq \epsilon\right\}\right) \\ &\geq \mathbb{P}\left(\limsup_{n \rightarrow \infty} \left\{\omega \in \Omega : |X_n(\omega) - X(\omega)| \leq \epsilon\right\}\right) = 1, \end{aligned}$$

for all  $\epsilon > 0$ . Therefore, we have  $f(X_n) \xrightarrow{a.s.} f(X)$ .  $\square$ .

## 2 Statistical Inference

### 2.1 Sufficiency & Likelihood

#### 2.1.1 Sufficiency

#### 2.1.2 Likelihood

### 2.2 Point Estimation

#### 2.2.1 Bias, Variance, Consistency and MSE

#### 2.2.2 Sufficient Statistics & Rao-Blackwell Theorem

**Theorem 2.1: Rao-Blackwell Theorem (RB)**

#### 2.2.3 Estimator Variance & Cramer-Rao Lower Bound

**Theorem 2.2: Cramer-Rao Lower Bound (CRLB)**

#### 2.2.4 Maximum Likelihood Estimation (MLE)

## B List of Definitions

1.1	Definition (Random variable)	2
1.2	Definition (Convergence in Distribution)	8
1.3	Definition (Convergence in Probability)	9
1.4	Definition (Convergence in $L^p$ norm)	10
1.5	Definition (Convergence almost-surely)	10

## B Important Theorems

1.1	Weak Law of Large Numbers ( <b>WLLN</b> )	4
1.2	Levy Continuity Theorem ( <b>LCT</b> )	5
1.3	Strong Law of Large Numbers ( <b>SLLN</b> )	6
1.4	Borel-Cantelli Lemma ( <b>BCL</b> )	7
1.5	Central Limit Theorem ( <b>CLT</b> )	8
1.6	Continuous Mapping Theorem ( <b>CMT</b> )	11
2.1	Rao-Blackwell Theorem ( <b>RB</b> )	12
2.2	Cramer-Rao Lower Bound ( <b>CRLB</b> )	12

## C Important Propositions

1.1	$X_n \xrightarrow{p} X \implies X_n \xrightarrow{d} X$	9
1.2	$X_n \xrightarrow{d} c \iff X_n \xrightarrow{p} c$	9
1.3	$X_n \xrightarrow{L^p} X \implies X_n \xrightarrow{p} X$	10
1.4	$X_n \xrightarrow{a.s.} X \implies X_n \xrightarrow{p} X$	11

## D References

### References

- Durrett, Rick (2010). *Probability: Theory and Examples*. 4th. USA: Cambridge University Press. ISBN: 0521765390.
- Gut, Allan (2004). *A Graduate Course in Probability*. Graduate Text in Mathematics.
- Pishro-Nik, Hossein (2014). *Introduction to Probability, Statistics and Random Processes*. Kappa Research, LLC.
- undefinedinar, Erhan (2011). *Probability and Stochastics*. Springer New York. ISBN: 9780387878591. URL: <http://dx.doi.org/10.1007/978-0-387-87859-1>.
- Wikipedia (2023). *Vitali set* — *Wikipedia, The Free Encyclopedia*. <http://en.wikipedia.org/w/index.php?title=Vitali%20set&oldid=1187241923>. [Online; accessed 24-December-2023].